

**STABILIZATION OF HYPOALLERGENIC,  
HYPERDIGESTIBLE PREVIOUSLY REDUCED PROTEINS**

**CROSS-REFERENCE**

This application is a continuation-in-part of application Serial No. 09/238,379, which is a continuation-in-part of application Serial No. 08/953,703 filed October 17, 1997 (now U.S. Patent 5,952,034 issued September 14, 1999), which is a continuation-in-part of application Serial No. 08/326,976 filed October 21, 1994 (now U.S. Letters Patent No. 5,792,506 issued August 11, 1998), which is a continuation-in-part of application Serial No. 08/211,673 filed November 21, 1994 ( now U.S. Patent No. 6,113,951 issued September 5, 2000) which is a national stage filing of International Application No. PCT/US92/08595 filed October 8, 1992, which is a continuation-in-part of application Serial No. 07/935,002 filed August 25, 1992 (now abandoned), which is a continuation-in-part of application Serial No. 07/776,109 filed October 12, 1991 (now abandoned).

Reference is also made to the following related applications and patents:

- (1) application Serial No. 08/483,930, filed June 7, 1995 (now U.S. Patent No. 6,114,504 issued September 5, 2000), which is a divisional of application Serial No. 08/211,673; (2) application Serial No. 09/046,780 filed March 23, 1998, which is a divisional of application Serial No. 08/326,976; (3) application Serial No. 09/296,927 filed March 22, 1999, which is a

divisional of application Serial No. 08/953,703; (4) application Serial No. 09/447,615 filed November 23, 1999, which is a divisional of application Serial No. 08/211,673; (5) application Serial No. 09/448,109 filed November 23, 1999, which is a divisional of application Serial No. 08/211,673; and (6) application Serial No. 09/448,111 filed November 23, 1999, which is a divisional of application Serial No. 08/211,673.


### FIELD OF THE INVENTION

The present invention relates to the use of thiol redox proteins to reduce seed proteins such as cereal proteins, and to reduce enzyme inhibitor proteins, venom toxin proteins, pollen proteins and the intramolecular disulfide bonds of certain other proteins. More particularly, the invention involves use of thioredoxin and glutaredoxin to reduce gliadins, glutenins, albumins and globulins to improve the characteristics of dough and baked goods and create new doughs and to reduce cystine containing proteins such as amylase and trypsin inhibitors so as to improve the quality of feed and cereal products. Additionally, the invention involves the isolation of a novel protein that inhibits pullulanase and the reduction of that novel protein by thiol redox proteins. The invention further involves the reduction by thioredoxin of 2S albumin proteins characteristic of oil-storing seeds. Also, the invention involves inactivating snake neurotoxins and certain insect and scorpion venom toxins *in vitro* and treating the corresponding toxicities in individuals. The invention also involves using thioredoxin and dithiothreitol to decrease the allergenicity of food and pollen allergens and to increase the proteolysis of food and pollen proteins and the digestibility of food and pollens. The invention also relates to pollen proteins which are reduced by lipoic acid or by reduced thiol-redox proteins or by thioredoxin in combination with lipoic acid for use in immunotherapy. The invention further involves use of thiol-redox proteins and lipoic acid to treat and prevent allergies and allergic symptoms. Also the invention relates to the stabilization by physiological

disulfides such as cystamine and oxidized glutathione of previously reduced proteins for purposes of mitigating allergenicity and increasing pepsin digestibility of the proteins.

### BACKGROUND OF THE INVENTION



5 Chloroplasts contain a ferredoxin/thioredoxin system comprised of ferredoxin, ferredoxin-thioredoxin reductase and thioredoxins f and m that links light to the regulation of enzymes of photosynthesis (Buchanan, B.B. (1991) "Regulation of CO<sub>2</sub> assimilation in oxygenic photosynthesis: The ferredoxin/thioredoxin system. Perspective on its discovery, present status  
10 and future development", *Arch. Biochem. Biophys.* 288:1-9; Scheibe, R. (1991), "Redox-modulation of chloroplast enzymes. A common principle for individual control", *Plant Physiol.* 96:1-3). Several studies have shown that plants also contain a system, analogous to the one established for animals and most microorganisms, in which thioredoxin (h-type) is reduced by NADPH  
15 and the enzyme, NADP-thioredoxin reductase (NTR) according to the following:



(Florencio F.J. *et al.* (1988), *Arch. Biochem. Biophys.* 266:496-507;  
20 Johnson, T.C. *et al.* (1987), *Plant Physiol.* 85:446-451; Suske, G. *et al.* (1979), *Z. Naturforsch. C.* 34:214-221). Current evidence suggests that the NADP/thioredoxin system is widely distributed in plant tissues and is housed in the mitochondria, endoplasmic reticulum and cytosol (Bodenstein-Lang, J. *et al.* (1989), *FEBS Lett.* 258:22-26; Marcus, F. *et al.* (1991), *Arch.*  
25 *Biochem. Biophys.* 287:195-198).

Thioredoxin h is also known to reductively activate cytosolic enzyme of carbohydrate metabolism, pyrophosphate fructose-6-P, 1-phosphotransferase or PFP (Kiss, F. *et al.* (1991), *Arch. Biochem. Biophys.* 287:337-340).

The seed is the only tissue for which the NADP/thioredoxin system has been ascribed physiological activity in plants. Also, thioredoxin h has been shown to reduce thionins in the laboratory (Johnson, T.C. *et al.* (1987), *Plant Physiol.* 85:446-451). Thionins are soluble cereal seed proteins, rich in cystine. In the Johnson, *et al.* investigation, wheat purothionin was experimentally reduced by NADPH via NADP-thioredoxin reductase (NTR) and thioredoxin h according to Eqs. 2 and 3.



Cereal seeds such as wheat, rye, barley, corn, millet, sorghum and rice contain four major seed protein groups. These four groups are the albumins, globulins, gliadins and the glutenins or corresponding proteins. The thionins belong to the albumin group or faction. Presently, wheat and rye are the only two cereals from which gluten or dough has been formed. Gluten is a tenacious elastic and rubbery protein complex that gives cohesiveness to dough. Gluten is composed mostly of the gliadin and glutenin proteins. It is formed when rye or wheat dough is washed with water. It is the gluten that gives bread dough its elastic type quality. Flour from other major crop cereals barley, corn, sorghum, oat, millet and rice and also from the plant, soybean do not yield a gluten-like network under the conditions used for wheat and rye.

Glutenins and gliadins are cystine containing seed storage proteins and are insoluble. Storage proteins are proteins in the seed which are broken down during germination and used by the germinating seedling to grow and develop. Prolamines are the storage proteins in grains other than wheat that correspond to gliadins while the glutelins are the storage proteins in grains other than wheat that correspond to glutenins. The wheat storage proteins account for up to 80% of the total seed protein (Kasarda, D.D. *et al.* (1976), *Adv. Cer. Sci. Tech.* 1:158-236; and Osborne, T.B. *et al.* (1893), *Amer. Chem. J.* 15:392-471). Glutenins and gliadins are considered important in the formation of dough and therefore the quality of bread. It has been shown from *in vitro* experiments that the solubility of seed storage proteins is increased on reduction (Shewry, P.R. *et al.* (1985), *Adv. Cer. Sci. Tech.* 7:1-83). However, previously, reduction of glutenins and gliadins was thought to lower dough quality rather than to improve it (Dahle, L.K. *et al.* (1966), *Cereal Chem.* 43:682-688). This is probably because the non-specific reduction with chemical reducing agents caused the weakening of the dough.

The "Straight Dough" and the "Pre-Ferment" methods are two major conventional methods for the manufacture of dough and subsequent yeast raised bread products.

For the Straight Dough method, all of the flour, water or other liquid, and other dough ingredients which may include, but are not limited to yeast, grains, salt, shortening, sugar, yeast nutrients, dough conditioners, and preservatives are blended to form a dough and are mixed to partial or full development. The resulting dough may be allowed to ferment for a period of time depending upon specific process or desired end-product characteristics.

The next step in the process is the mechanical or manual division of the dough into appropriate size pieces of sufficient weight to ensure achieving the targeted net weight after baking, cooling, and slicing. The dough pieces are often then rounded and allowed to rest (Intermediate Proof) for varying  
5 lengths of time. This allows the dough to "relax" prior to sheeting and molding preparations. The time generally ranges from 5-15 minutes, but may vary considerably depending on specific processing requirements and formulations. The dough pieces are then mechanically or manually formed into an appropriate shape are then usually given a final "proof" prior to  
10 baking. The dough pieces are then baked at various times, temperatures, and steam conditions in order to achieve the desired end product.

In the Pre-Ferment method, yeast is combined with other ingredients and allowed to ferment for varying lengths of time prior to final mixing of the bread or roll dough. Baker's terms for these systems include "Water Brew",  
15 "Liquid Ferment", "Liquid Sponge", and "Sponge/Dough". A percentage of flour ranging from 0-100% is combined with the other ingredients which may include but are not limited to water, yeast, yeast nutrients and dough conditioners and allowed to ferment under controlled or ambient conditions for a period of time. Typical times range from 1-5 hours. The ferment may  
20 then be used as is, or chilled and stored in bulk tanks or troughs for later use. The remaining ingredients are added (flour, characterizing ingredients, additional additives, additional water, etc.) and the dough is mixed to partial or full development.

The dough is then allowed to ferment for varying time periods. Typically, as  
25 some fermentation has taken place prior to the addition of the remaining ingredients, the time required is minimal (*i.e.*, 10-20 min), however, variations are seen depending upon equipment and product type. Following

the second fermentation step, the dough is then treated as in the Straight Dough Method.

As used herein the term "dough mixture" describes a mixture that minimally comprises a flour or meal and a liquid, such as milk or water.

- 5 As used herein the term "dough" describes an elastic, pliable protein network mixture that minimally comprises a flour, or meal and a liquid, such as milk or water.

- As used herein the term "dough ingredient" may include, but is not exclusive of, any of the following ingredients: flour, water or other liquid, grain,  
10 yeast, sponge, salt, shortening, sugar, yeast nutrients, dough conditioners and preservatives.

- As used herein, the term "baked good" includes but is not exclusive of all bread types, including yeast-leavened and chemically-leavened and white and variety breads and rolls, english muffins, cakes and cookies, confectionery  
15 coatings, crackers, doughnuts and other sweet pastry goods, pie and pizza crusts, pretzels, pita and other flat breads, tortillas, pasta products, and refrigerated and frozen dough products.

- While thioredoxin has been used to reduce albumins in flour, thiol redox proteins have not been used to reduce glutenins and gliadins nor other water  
20 insoluble storage proteins, nor to improve the quality of dough and baked goods. Thiol redox proteins have also not been used to improve the quality of gluten thereby enhancing its value nor to prepare dough from crop cereals such as barley, corn, sorghum, oat, millet and rice or from soybean flour.

Many cereal seeds also contain proteins that have been shown to act as inhibitors of enzymes from foreign sources. It has been suggested that these enzyme inhibitors may afford protection against certain deleterious organisms (Garcia-Olmedo, F. *et al.* (1987), *Oxford Surveys of Plant Molecular and Cell Biology* 4:275-335; Birk, Y. (1976), *Meth. Enzymol.* 45:695-739, and Laskowski, M., Jr. *et al.* (1980), *Ann. Rev. Biochem.* 49:593-626). Two such type enzyme inhibitors are amylase inhibitors and trypsin inhibitors. Furthermore, there is evidence that a barley protein inhibitor (not tested in this study) inhibits an  $\alpha$ -amylase from the same source (Weselake, R.J. *et al.* (1983), *Plant Physiol.* 72:809-812). Unfortunately, the inhibitor protein often causes undesirable effects in certain food products. The trypsin inhibitors in soybeans, notably the Kunitz trypsin inhibitor (KTI) and Bowman-Birk trypsin inhibitor (BBTI) proteins, must first be inactivated before any soybean product can be ingested by humans or domestic animals. It is known that these two inhibitor proteins become ineffective as trypsin inhibitors when reduced chemically by sodium borohydride (Birk, Y. (1985), *Int. J. Peptide Protein Res.* 25:113-131, and Birk, Y. (1976), *Meth. Enzymol.* 45:695-739). These inhibitors like other proteins that inhibit proteases contain intramolecular disulfides and are usually stable to inactivation by heat and proteolysis (Birk (1976), *supra.*; Garcia-Olmedo *et al.* (1987), *supra.*, and Ryan (1980). Currently, to minimize the adverse effects caused by the inhibitors these soybean trypsin inhibitors and other trypsin inhibitors in animal and human food products are being treated by exposing the food to high temperatures. The heat treatment, however, does not fully eliminate inhibitor activity. Further, this process is not only expensive but it also destroys many of the other proteins which have important nutritional value. For example, while 30 min at 120°C leads to complete inactivation of the BBTI of soy flour, about 20% of the original KTI activity remains (Friedman *et al.*, 1991). The prolonged or higher temperature treatments required for full inactivation of inhibitors results in



destruction of amino acids such as cystine, arginine, and lysine (Chae *et al.*, 1984; Skrede and Krogdahl, 1985).

There are also several industrial processes which require  $\alpha$ -amylase activity. One example is the malting of barley which requires active  $\alpha$ -amylase.

- 5 Inactivation of inhibitors such as the barley amylase/subtilisin (asi) inhibitor and its equivalent in other cereals by thiol redox protein reduction would enable  $\alpha$ -amylases to become fully active sooner than with present procedures, thereby shortening time for malting or similar processes.

- Thiol redox proteins have also not previously been used to inactivate trypsin  
10 or amylase inhibitor proteins. The reduction of trypsin inhibitors such as the Kunitz and Bowman-Birk inhibitor proteins decreases their inhibitory effects (Birk, Y. (1985), *Int. J. Peptide Protein Res.* 25:113-131). A thiol redox protein linked reduction of the inhibitors in soybean products designed for consumption by humans and domestic animals would require no heat or lower  
15 heat than is presently required for protein denaturization, thereby cutting the costs of denaturation and improving the quality of the soy protein. Also a physiological reductant, a so-called clean additive (*i.e.*, an additive free from ingredients viewed as "harmful chemicals") is highly desirable since the food industry is searching for alternatives to chemical additives. Further the  
20 ability to selectively reduce the major wheat and seed storage proteins which are important for flour quality (*e.g.*, the gliadins and the glutenins) in a controlled manner by a physiological reductant such as a thiol redox protein would be useful in the baking industry for improving the characteristics of the doughs from wheat and rye and for creating doughs from other grain flours  
25 such as cereal flours or from cassava or soybean flour.

The family of 2S albumin proteins characteristic of oil-storing seeds such as castor bean and Brazil nut (Kreis *et al.* 1989; Youle and Huang, 1981) which

are housed within protein bodies in the seed endosperm or cotyledons (Ashton *et al.* 1976; Weber *et al.* 1980), typically consist of dissimilar subunits connected by two intermolecular disulfide bonds -- one subunit of 7 to 9 kDa and the other of 3 to 4 kDa. The large subunit contains two  
5 intramolecular disulfide groups, the small subunit contains none. The intramolecular disulfides of the 2S large subunit show homology with those of the soybean Bowman-Birk inhibitor (Kreis *et al.* 1989) but nothing is known of the ability of 2S proteins to undergo reduction under physiological conditions.

- 10 These 2S albumin proteins are rich in methionine. Recently transgenic soybeans which produce Brazil nut 2S protein have been generated. Reduction of the 2S protein in such soybeans could enhance the integration of the soy proteins into a dough network resulting in a soybread rich in methionine. In addition, these 2S proteins are often allergens. Reduction of  
15 the 2S protein would result in the cessation of its allergic activity.

Pullulanase ("debranching enzyme") is an enzyme that breaks down the starch of the endosperm of cereal seeds by hydrolytically cleaving  $\alpha$ -1,6 bonds. Pullulanase is an enzyme fundamental to the brewing and baking industries. Pullulanase is required to break down starch in malting and in  
20 certain baking procedures carried out in the absence of added sugars or other carbohydrates. Obtaining adequate pullulanase activity is a problem especially in the malting industry. It has been known for some time that dithiothreitol (DTT, a chemical reductant for thioredoxin) activates pullulanase of cereal preparations (*e.g.*, barley, oat and rice flours). A  
25 method for adequately activating or increasing the activity of pullulanase with a physiologically acceptable system, could lead to more rapid malting methods and, owing to increased sugar availability, to alcoholic beverages such as beers with enhanced alcoholic content.

Death and permanent injury resulting from snake bites are serious problems in many African, Asian and South American countries and also a major concern in several southern and western areas of the United States. Venoms from snakes are characterized by active protein components (generally  
5 several) that contain disulfide (S-S) bridges located in intramolecular (intrachain) cystines and in some cases in intermolecular (interchain) cystines. The position of the cystine within a given toxin group is highly conserved. The importance of intramolecular S-S groups to toxicity is evident from reports showing that reduction of these groups leads to a loss of toxicity in  
10 mice (Yang, C.C. (1967) *Biochim. Biophys. Acta.* 133:346-355; Howard, B.D. *et al.* (1977) *Biochemistry* 16:122-125). The neurotoxins of snake venom are proteins that alter the release of neurotransmitter from motor nerve terminals and can be presynaptic or postsynaptic. Common symptoms observed in individuals suffering from snake venom neurotoxicity include  
15 swelling, edema and pain, fainting or dizziness, tingling or numbing of affected part, convulsions, muscle contractions, renal failure, in addition to long-term necrosis and general weakening of the individual, etc.

The presynaptic neurotoxins are classified into two groups. The first group, the  $\beta$ -neurotoxins, include three different classes of proteins, each having a  
20 phospholipase A<sub>2</sub> component that shows a high degree of conservation. The proteins responsible for the phospholipase A<sub>2</sub> activity have from 6 to 7 disulfide bridges. Members of the  $\beta$ -neurotoxin group are either single chain (*e.g.*, caudotoxin, notexin and agkistrodotoxin) or multichain (*e.g.*, crotoxin, ceruleotoxin and *Vipera* toxin).  $\beta$ -bungarotoxin, which is made up of two  
25 subunits, constitutes a third group. One of these subunits is homologous to the Kunitz-type proteinase inhibitor from mammalian pancreas. The multichain  $\beta$ -neurotoxins have their protein components linked ionically whereas the two subunits of  $\beta$ -bungarotoxin are linked covalently by an intermolecular disulfide. The B chain subunit of  $\beta$ -bungarotoxin, which is

also homologous to the Kunitz-type proteinase inhibitor from mammalian pancreas, has 3 disulfide bonds.

- The second presynaptic toxin group, the facilitatory neurotoxins, is devoid of enzymatic activity and has two subgroups. The first subgroup, the
- 5 dendrotoxins, has a single polypeptide sequence of 57 to 60 amino acids that is homologous with Kunitz-type trypsin inhibitors from mammalian pancreas and blocks voltage sensitive potassium channels. The second subgroup, such as the fasciculins (*e.g.*, fasciculin 1 and fasciculin 2) are cholinesterase inhibitors and have not been otherwise extensively studied.
- 10 The postsynaptic neurotoxins are classified either as long or short neurotoxins. Each type contains S-S groups, but the peptide is unique and does not resemble either phospholipase A<sub>2</sub> or the Kunitz or Kunitz-type inhibitor protein. The short neurotoxins (*e.g.*, erabutoxin a and erabutoxin b) are 60 to 62 amino acid residues long with 4 intramolecular disulfide bonds.
- 15 The long neurotoxins (*e.g.*,  $\alpha$ -bungarotoxin and  $\alpha$ -cobratoxin) contain from 65 to 74 residues and 5 intramolecular disulfide bonds. Another type of toxins, the cytotoxins, acts postsynaptically but its mode of toxicity is ill defined. These cytotoxins show obscure pharmacological effects, *e.g.*, hemolysis, cytolysis and muscle depolarization. They are less toxic than the
- 20 neurotoxins. The cytotoxins usually contain 60 amino acids and have 4 intramolecular disulfide bonds. The snake venom neurotoxins all have multiple intramolecular disulfide bonds.

- The current snake antitoxins used to treat poisonous snake bites following first aid treatment in individuals primarily involve intravenous injection of
- 25 antivenom prepared in horses. Although it is not known how long after envenomation the antivenom can be administered and be effective, its use is recommended up to 24 hours. Antivenom treatment is generally

accompanied by administration of intravenous fluids such as plasma, albumin, platelets or specific clotting factors. In addition, supporting medicines are often given, for example, antibiotics, antihistamines, antitetanus agents, analgesics and sedatives. In some cases, general treatment  
5 measures are taken to minimize shock, renal failure and respiratory failure. Other than administering calcium-EDTA in the vicinity of the bite and excising the wound area, there are no known means of localized treatment that result in toxin neutralization and prevention of toxic uptake into the blood stream. Even these localized treatments are of questionable  
10 significance and are usually reserved for individuals sensitive to horse serum (Russell, F.E. (1983) *Snake Venom Poisoning*, Schollum International, Inc. Great Neck, NY).

The term "individual" as defined herein refers to an animal or a human.

Most of the antivenoms in current use are problematic in that they can  
15 produce harmful side effects in addition to allergic reactions in patients sensitive to horse serum (up to 5% of the patients). Nonallergic reactions include pyrogenic shock, and complement depletion (Chippaur, J.-P. *et al.* (1991) *Reptile Venoms and Toxins*, A.T. Tu, ed., Marcel Dekker, Inc., pp. 529-555).

20 It has been shown that thioredoxin, in the presence of NADPH and thioredoxin reductase reduces the bacterial neurotoxins tetanus and botulinum A *in vitro* (Schiavo, G. *et al.* (1990) *Infection and Immunity* 58:4136-4141; Kistner, A. *et al.* (1992) *Naunyn-Schmiedeberg's Arch Pharmacol* 345:227-234). Thioredoxin was effective in reducing the interchain disulfide link of  
25 tetanus toxin and such reduced tetanus toxin was no longer neurotoxic (Schiavo *et al.*, *supra.*). However, reduction of the interchain disulfide of botulinum A toxin by thioredoxin was reported to be much more sluggish

(Kistner *et al.*, *supra.*). In contrast to the snake neurotoxin studied in the course of this invention, the tetanus research group (Schiavo *et al.*, *supra.*) found no evidence in the work done with the tetanus toxin that reduced thioredoxin reduced toxin intrachain disulfide bonds. There was also no  
5 evidence that thioredoxin reduced intrachain disulfides in the work done with botulinum A. The tetanus and botulinum A toxins are significantly different proteins from the snake neurotoxins in that the latter (1) have a low molecular weight; (2) are rich in intramolecular disulfide bonds; (3) are resistant to trypsin and other animal proteases; (4) are active without  
10 enzymatic modification, *e.g.*, proteolytic cleavage; (5) in many cases show homology to animal proteins, such as phospholipase A<sub>2</sub> and Kunitz-type proteases; (6) in most cases lack intermolecular disulfide bonds, and (7) are stable to agents such as heat and proteases.

Reductive inactivation of snake toxins *in vitro* by incubation with 1 %  $\beta$ -  
15 mercaptoethanol for 6 hours and incubation with 8M urea plus 300mM  $\beta$ -mercaptoethanol has been reported in the literature (Howard, B.D. *et al.* (1977) *Biochemistry* 16:122-125; Yang, C.C. (1967) *Biachim. Biophys. Acta.* 133:346-355). These conditions, however, are far from physiological. As defined herein the term "inactivation" with respect to a toxin protein means  
20 that the toxin is no longer biologically active *in vitro*, in that the toxin is unable to link to a receptor. Also as used herein, "detoxification" is an extension of the term "inactivation" and means that the toxin has been neutralized in an individual as determined by animal toxicity tests.

Bee venom is a complex mixture with at least 40 individual components, that  
25 include major components as melittin and phospholipase A<sub>2</sub>, representing respectively 50% and 12% of the total weight of the venom, and minor components such as small proteins and peptides, enzymes, amines, and amino acids.

Melittin is a polypeptide consisting of 26 amino acids with a molecular weight of 2840. It does not contain a disulfide bridge. Owing to its high affinity for the lipid-water interphase, the protein permeates the phospholipid bilayer of the cell membranes, disturbing its organized structure. Melittin is not by itself a toxin but it alters the structure of membranes and thereby increases the hydrolytic activity of phospholipase A<sub>2</sub>, the other major component and the major allergen present in the venom.

Bee venom phospholipase A<sub>2</sub> is a single polypeptide chain of 128 amino acids, is cross-linked by four disulfide bridges, and contains carbohydrate.

10 The main toxic effect of the bee venom is due to the strong hydrolytic activity of phospholipase A<sub>2</sub> achieved in association with melittin.

The other toxic proteins in bee venom have a low molecular weight and contain at least two disulfide bridges that seem to play an important structural role. Included are a protease inhibitor (63-65 amino acids), MCD or 401-peptide (22 amino acids) and apamin (18 amino acids).

15

Although there are thousands of species of bees, only the honey bee, *Apis mellifera*, is a significant cause of allergic reactions. The response ranges from local discomfort to systemic reactions such as shock, hypotension, dyspnea, loss of consciousness, wheezing and/or chest tightness that can result in death. The only treatment that is used in these cases is the injection of epinephrine.

20

The treatment of bee stings is important not only for individuals with allergic reactions. The "killer" or Africanized bee, a variety of honey bee is much more aggressive than European honey bees and represents a danger in both South and North America. While the lethality of the venom from the Africanized and European bees appears to be the same (Schumacher, M.I.

25

*et al.* (1989) *Nature* 337:413), the behaviour pattern of the hive is completely different. It was reported that Africanized bees respond to colony disturbance more quickly, in greater numbers and with more stinging (Collins, A. M. *et al.* (1982) *Science* 218:72-74). A mass attack by

5 Africanized bees may produce thousands of stings on one individual and cause death. The "killer" bees appeared as a result of the interbreeding between the African bee (*Apis mellifera scutellata*) and the European bee (*Apis mellifera mellifera*). African bees were introduced in 1956 into Brazil with the aim of improving honey production being a more tropically adapted

10 bee. Africanized bees have moved from South America to North America, and they have been reported in Texas and Florida.

In some areas of the world such as Mexico, Brazil, North Africa and the Middle East, scorpions present a life hazard to humans. However, only the scorpions of family Buthidae (genera, *Androctonus*, *Buthus*, *Centruroides*,

15 *Leiurus* and *Tityus*) are toxic for humans. The chemical composition of the scorpion venom is not as complex as snake or bee venom. Scorpion venom contains mucopolysaccharides, small amounts of hyaluronidase and phospholipase, low molecular-weight molecules, protease inhibitors, histamine releasers and neurotoxins, such as serotonin. The neurotoxins

20 affect voltage-sensitive ionic channels in the neuromuscular junction. The neurotoxins are basic polypeptides with three to four disulfide bridges and can be classified in two groups: peptides with from 61 to 70 amino acids, that block sodium channel, and peptides with from 36 to 39 amino acids, that block potassium channel. The reduction of disulfide bridges on the

25 neurotoxins by nonphysiological reductants such as DTT or  $\beta$ -mercaptoethanol (Watt, D.D. *et al.* (1972) *Toxicon* 10:173-181) lead to loss of their toxicity.



Symptoms of animals stung by poisonous scorpions include hyperexcitability, dyspnea, convulsions, paralysis and death. At present, antivenin is the only antidote for scorpion stings. The availability of the venom is a major problem in the production of antivenin. Unlike snake venom, scorpion  
5 venom is very difficult to collect, because the yield of venom per specimen is limited and in some cases the storage of dried venom leads to modification of its toxicity. An additional problem in the production of antivenins is that the neurotoxins are very poor antigens.

The reductive inactivation of snake, bee and scorpion toxins under  
10 physiological conditions has never been reported nor has it been suggested that the thiol redox agents, such as reduced lipoic acid, DTT, or reduced thioredoxin could act as an antidote to these venoms in an individual.

Food allergies also represent a long-standing problem important both nationally and internationally. Up to 5% of children under age 12 and 1% of  
15 adults are clinically affected in the U.S. population (Adverse Reactions to Foods -- AAAI and NIAD Report, 1984, NIH Pub. No. 84-2442, pp. 2, 3). In some countries, the figures are higher, and, throughout the world, the problem is considered to be increasing, especially in infants (T. Matsuda and R. Nakamura 1993 Molecular structure and immunological properties of  
20 Food Allergens, Trends in Food Science & Technology 4, 289-293). The problem extends to a wide range of foods. Food allergies in general have recently achieved an increased profile as a result of the concern about transgenic foods.

Milk represents a significant problem, especially in infants. Wheat and soy  
25 allergies are of growing importance as new populations adopt these foods and are of increased concern in pet (especially dog) foods. Beef, rice and egg

also cause serious allergies in many individuals and again are of significant concern with respect to pet food.

Many of the major allergenic proteins in the above mentioned foods have intramolecular disulfide (S-S) bonds but so far two treatments have been applied commercially to minimize food allergies: (1) heat, and (2) enzymatic proteolysis. In both cases, success has been only partial. While lowering allergenicity, heat treatment has not eliminated the problem, even in the best of cases, because the responsible proteins are typically heat stable. Moreover, heat lowers product quality by destroying nutritionally important amino acids such as lysine, cysteine and arginine. Enzymatic proteolysis is more successful in reducing allergenicity, but desirable food properties such as flavor are usually lost and treatment is costly. Therefore a physiologically safe system that would bring about a decrease in or loss of allergenicity when applied to allergenic foods without a resulting loss in flavor and nutrition would be extremely valuable.

Certain major pollen allergens are known to be disulfide proteins that are highly resistant to temperature. Two pollen proteins are described as major allergens in ragweed pollen. One is a small protein of 5 kDa, Amb a V, containing four disulfide bridges (Goodfriend, L. *et al.* (1985), "Ra5G, a homologue of Ra5 in giant ragweed pollen: isolation, HLA-DR- associated activity and amino acid sequence", *Mol. Immunol.* 22:899-906; Metzler, W.J. *et al.* (1992), "Determination of the three-dimensional solution structure of ragweed allergen Amb t V by nuclear magnetic resonance spectroscopy" *Biochemistry* 31:5117-5127; Mole, L.E., *et al.* (1975), "The amino acid sequence of ragweed pollen allergen Ra5" *Biochemistry* 14:1216-1220; Metzler, W.J., *et al.* (1992), "Proton resonance assignments and three-dimensional solution structure of the ragweed allergen Amb a V by nuclear magnetic resonance spectroscopy" *Biochemistry* 31:8697-8705). This

protein is considered to be homologous in both the short and giant ragweed species. The short ragweed protein which is designated Amb a V and the giant ragweed which is now designated Amb t V, both previously called Ra 5, exhibit a 45 % sequence similarity.

- 5 The other major allergen represents a family of 41 kDa proteins, named Amb a 1.1, Amb a 1.2, Amb a 1.3 and Amb a 1.4. While no disulfide bridges have been described, these proteins contain multiple cysteines (Rafnar, T. *et al.* (1991), "Cloning of Amb a I (antigen E), the major allergen family of short ragweed pollen" *J. Biol. Chem.* 266:1229-1236; Griffith, I.J. *et al.* (1991), "Sequence polymorphism of Amb a I and Amb a II, the major allergens in *Ambrosia artemisiifolia* (short ragweed)" *Int. Arch. Allergy Appl. Immunol.* 96:296-304). Yet other known allergens are disulfide proteins such as the western ragweed, Amb P 5-A and -B, each 8.5 kDa with three disulfide bridges (Ghosh, B. *et al.* (1994), "Immunologic and molecular
- 10 characterization of Amb p V allergens from *Ambrosia psilostachya* (western Ragweed) pollen" *J. Immunol.* 152:2882-2889) and a short ragweed 11.4 kDa plastocyanin like protein, caUed Ra 3, with one disulfide bridge (Klapper, D.G. *et al.* (1980), "Amino acid sequence of ragweed allergen Ra3" *Biochemistry* 19:5729-5734).
- 15
- 20 The 5 kDa Amb V ragweed pollen proteins have a well-defined structure and the positions of the four intrachain disulfide bonds are precisely known (Metzler, W.J. *et al.* (1992) *Biochemistry* 31:5117-5127 and 8697-8705). Previous work has shown that, when reduced under denaturing conditions by chemical agents (urea plus either dithiothreitol or  $\beta$ -mercaptoethanol), the
- 25 immune response shifts from IgE (allergic) to an IgG (defense) because IgG production is enhanced (Zhu, X. *et al.* (1995), "T cell epitope mapping of ragweed pollen allergen *Ambrosia artemisiifolia* (Amb a 5) and *Ambrosia*

trifida (Amb t 5) and the role of free sulfhydryl groups in T cell recognition”  
J. Immunol. 155:5064-73).

Pollen allergies are currently being treated by conventional immunotherapy with undenatured pollen extract. However, such treatment, especially in  
5 children, carries a certain risk of anaphylactic reactions which are potentially lethal. Consequently, there is a need for an attenuated pollen protein or pollen extract for use in immunotherapy that would reduce or eliminate the possibility of anaphylactic reactions. There is also a need for a  
10 physiologically safe system that could determine whether or not an allergen for a particular individual is a disulfide protein. Further, eye drops, nose sprays, aerosols, or dispersants for vaporizers or humidifiers that would alleviate allergy symptoms but also produce less side effects than the currently available products would be extremely valuable.

Two sulfhydryl (SH) groups can often spontaneously react to form an S-S  
15 bond in the presence of oxygen. Therefore, the effects of the NADP-thioredoxin system (NTS) and other reductants to decrease allergenicity and increase digestibility of proteins by pepsin may not be permanent. Consequently, there is also a need for the stabilization of proteins, including allergenic proteins reduced by the NTS and other reductants. Stable forms of  
20 hypoallergenic, hyperdigestible proteins, with a long shelf life that are allergenic in their native state would be extremely valuable in the food industry and in medicine.

### SUMMARY OF THE INVENTION

It is an aspect herein to provide a method for reducing a non thionin cystine  
25 containing protein.

It is a second aspect herein to provide methods utilizing a thiol redox protein alone or in combination with a reductant or reduction system to reduce glutenins or gliadins present in flour or seeds.

- 5 It is also an aspect herein to provide methods using a thiol redox protein alone or in combination with a reductant or reduction system to improve dough strength and baked goods characteristics such as better crumb quality, softness of the baked good and higher loaf volume.

It is a further aspect herein to provide formulations containing a thiol redox protein useful in practicing such methods.

- 10 Still a further aspect herein is to provide a method for producing a dough from rice, corn, soybean, barley, oat, cassava, sorghum or millet flour.

Yet another aspect is to provide a method for producing an improved gluten or for producing a gluten-like product from cereal grains other than wheat and rye.

- 15 It is further an aspect herein to provide a method of reducing an enzyme inhibitor protein having disulfide bonds.

Still another aspect herein is to provide yeast cells genetically engineered to express or overexpress thioredoxin.

- 20 Still yet another aspect herein is to provide yeast cells genetically engineered to express or overexpress NADP-thioredoxin reductase.

1 Still yet a further aspect herein is to provide a method for improving the quality of dough or a baked good using such genetically engineered yeast cells.

Yet still another aspect herein is to provide a method of reducing the  
5 intramolecular disulfide bonds of a non-thionin, non chloroplast protein containing more than one intramolecular cystine comprising adding a thiol redox protein to a liquid or substance containing the cystines containing protein, reducing the thiol redox protein and reducing the cystines containing protein by means of the thiol redox protein.

10 Another aspect herein is to provide an isolated pullulanase inhibitor protein having disulfide bonds and a molecular weight of between 8 to 15 kDa.

Still another aspect herein is to provide a method of increasing the activity of pullulanase derived from barley or wheat endosperm comprising adding  
15 thioredoxin to a liquid or substance containing the pullulanase and reducing the thioredoxin thereby increasing the pullulanase activity.

Still another aspect herein is to provide a method of reducing an animal venom toxic protein having one or more intramolecular cystines comprising contacting the cystine containing protein with an amount of a thiol redox (SH) agent effective for reducing the protein, and maintaining the contact for  
20 a time sufficient to reduce one or more disulfide bridges of the one or more intramolecular cystines thereby reducing the neurotoxin protein. The thiol redox (SH) agent may be a reduced thioredoxin, reduced lipoic acid in the presence of a thioredoxin, DTT or DTT in the presence of a thioredoxin and the snake neurotoxin protein may be a presynaptic or postsynaptic  
25 neurotoxin.

Still a further aspect of the invention is to provide a composition comprising a snake neurotoxin protein and a thiol redox (SH) agent.

- Still yet another aspect of the invention is to provide a method of reducing an animal venom toxic protein having one or more intramolecular cystines
- 5 comprising contacting the protein with amounts of NADP-thioredoxin reductase, NADPH or an NADPH generator system and a thioredoxin effective for reducing the protein, and maintaining the contact for a time sufficient to reduce one or more disulfide bridges of the one or more intramolecular cystines thereby reducing the protein.
- 10 Yet another aspect herein is to provide a method of inactivating, *in vitro*, a snake neurotoxin having one or more intramolecular cystines comprising adding a thiol redox (SH) agent to a liquid containing the toxin wherein the amount of the agent is effective for reducing the toxin.

- Yet a further aspect herein is to provide a method of treating venom toxicity
- 15 in an individual comprising administering, to an individual suffering from venom toxicity, amounts of a thiol redox (SH) agent effective for reducing or alleviating the venom toxicity.

- In accordance with the aspects of the invention, methods are provided for improving dough characteristics comprising the steps of mixing a thiol redox
- 20 protein with dough ingredients to form a dough and baking said dough.

Also, in accordance with the aspects of the invention, a method is provided for inactivating an enzyme inhibitor protein in a grain food product comprising the steps of mixing a thiol redox protein with the seed product, reducing the thiol redox protein by a reductant or reduction system and

reducing the enzyme inhibitor by the reduced thiol redox protein, the reduction of the enzyme inhibitor inactivating the enzyme inhibitor.

- The thiol redox proteins in use herein can include thioredoxin and glutaredoxin. The thioredoxin includes but is not exclusive of *E. coli*
- 5 thioredoxin, thioredoxin h, f and m and animal thioredoxins. A reductant of thioredoxin used herein can include lipoic acid or a reduction system such as NADPH in combination with NADP thioredoxin reductase (NTR). The reductant of glutaredoxin can include reduced glutathione in conjunction with the reduction system NADPH and glutathione reductase. NADPH can be
- 10 replaced with an NADPH generator or generator composition such as one consisting of glucose 6-phosphate, NADP and glucose 6-phosphate dehydrogenase from a source such as yeast. The NADPH generator is added together with thioredoxin and NADP-thioredoxin reductase at the start of the dough making process.
- 15 It should be noted that the invention can also be practiced with cysteine containing proteins. The cysteines can first be oxidized and then reduced via thiol redox protein.

- Further in accordance with the aspects of the invention, a method is provided for decreasing the allergenicity of an allergenic food protein comprising the
- 20 steps of contacting the protein with an amount of thioredoxin, NTR and NADPH or an amount of DTT in the presence of thioredoxin effective for decreasing the allergenicity of the protein and administering the contacted protein in step (a) to an animal, thereby decreasing the allergenic symptoms in said animal that would otherwise occur if the animal received the untreated
- 25 protein.



Another aspect of the invention is to provide a hypoallergenic ingestible food. The food was made hypoallergenic by prior treatment with thioredoxin in the presence of NTR and NADPH. The food can be beef, milk, soy, egg, rice or wheat.

- 5 A further aspect of the invention is to provide a method for improving the proteolysis and therefore the digestibility of food and allergen proteins and consequently to also provide more digestible foods, many of which are allergenic. The foods and allergens are made more susceptible to proteolysis and more digestible by prior treatment with thioredoxin in the presence of
- 10 reductants of thioredoxin such as those described above. Appropriate foods include soy, nuts, milk, whey, beef, egg, bread, other wheat products, and other grain products.

- Still another aspect of the invention is to provide a method for decreasing the allergenicity of an allergenic pollen protein comprising the steps of contacting
- 15 the protein with an amount of reduced thioredoxin effective for decreasing the allergenicity of the protein and administering the thioredoxin reduced protein to an animal in immunotherapeutic doses thereby decreasing the allergenic symptoms of said animal that would otherwise occur if the animal was exposed to the untreated protein.

- 20 Yet another aspect of the invention is to provide a hypoallergenic pollen or pollen protein with reduced disulfide bonds for immunotherapy.

- A further aspect of this invention is to provide a method for determining whether or not an allergen for a particular individual is a disulfide protein comprising administering an allergy test to said individual to identify said
- 25 allergen, treating said identified allergen protein *in vitro* with reduced

thioredoxin and analyzing said treated allergen protein for disulfide bond reduction.

An additional aspect of the invention is to provide a stable hypoallergenic, edible food or food protein with reduced disulfide bonds.

- 5 Another aspect of the invention is to provide a stable hypoallergenic pollen or pollen protein with reduced disulfide bonds for immunotherapy.

Still another aspect of the invention is to provide a stable hyperdigestible edible food or food protein with reduced disulfide bonds.

- 10 A further aspect of the invention is to provide a stable hyperdigestible pollen or pollen protein with reduced disulfide bonds for immunotherapy.

- Still a further aspect of the invention is to provide a method for making a stable hypoallergenic protein by reducing a protein containing disulfide bonds with thioredoxin, nicotinamide adenine dinucleotide phosphate-thioredoxin reductase and NADPH or an amount of dithiothreitol or an amount of lipoic acid effective for reducing the protein and reacting the reduced protein with an amount of a physiological disulfide effective for stabilizing the reduced protein.
- 15

- Still an additional aspect of the invention is to provide a method for making a stable hyperdigestible protein by reducing a protein containing disulfide bonds with thioredoxin, nicotinamide adenine dinucleotide phosphate-thioredoxin reductase and NADPH or an amount of dithiothreitol or an amount of lipoic acid effective for reducing the protein and reacting the reduced protein with an amount of a physiological disulfide effective for stabilizing the reduced protein.
- 20

### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a bar graph showing the mitigation of a skin test response to soy allergen by treating soy allergenic extract with reduced thioredoxin.

Fig. 2 is a bar graph showing the mitigation of a skin test response to milk allergen by treating milk allergenic extract with reduced thioredoxin.

Fig. 3 is a bar graph showing the mitigation of a skin test response to wheat allergen by treating wheat allergenic extract with reduced thioredoxin.

Fig. 4 is a bar graph showing the mitigation of a skin test response to beef allergen by treating beef allergenic extract with reduced thioredoxin at room temperature.

Fig. 5 is a bar graph showing the mitigation of a skin test response to beef allergen by treating beef allergenic extract with reduced thioredoxin at 37°C.

Fig. 6 is a bar graph showing the mitigation of a gastrointestinal allergenic response to soy and wheat by treating diets with reduced thioredoxin.

Fig. 7 is a photograph of an SDS polyacrylamide electrophoretic gel showing the extent of thioredoxin-linked and glutathione-linked reduction by means of fluorescence and protein staining.

Fig. 8 is a drawing of the tertiary structure of oxidized bovine  $\beta$ -lactoglobulin showing the disulfide bridges and free sulfhydryl.

Fig. 9 is a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the effect of thioredoxin-linked reduction on pepsin digestion of bovine  $\beta$ -lactoglobulin.

Fig. 10A represents a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the effect of time on the digestion of untreated milk.

5 Fig. 10B represents a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the effect of time on the digestion of milk following thioredoxin-linked reduction that occurred at 55°C.

Fig. 10C represents a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the effect of time on the digestion of milk following thioredoxin-linked reduction that occurred at 4°C.

10 Fig. 11A is a bar graph showing the thioredoxin-linked mitigation of a skin test response in a dog highly sensitive to milk.

Fig. 11B is a bar graph showing the thioredoxin-linked mitigation of a skin test response in a dog mildly sensitive to milk.

15 Fig. 12A is a bar graph showing the effect of thioredoxin-linked reduction and digestibility on the allergenicity of  $\beta$ -lactoglobulin.

Fig. 12B is a bar graph showing the effect of thioredoxin-linked reduction and digestibility on the allergenicity of milk.

Fig. 13 is a drawing of the tertiary structure of oxidized bovine  $\beta$ -lactoglobulin showing the mouse antibody (MAb) epitopes.

20 Fig. 14 is a graph of a computer generated molecular model of the tertiary structure of bovine  $\beta$ -lactoglobulin showing predicted monoclonal antibody epitope changes after cystine mutagenesis.

Fig. 15 is a photograph of an SDS polyacrylamide electrophoretic gel showing the extent of thioredoxin-linked and glutathione-linked reduction of ragweed pollen allergens by means of fluorescence.

Fig. 16A is a photograph of a protein stained SDS polyacrylamide  
5 electrophoretic gel showing the effect of time on the digestion of untreated ragweed allergen Amb t V.

Fig. 16B is a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the effect of time on the digestion of ragweed allergen Amb t V following thioredoxin-linked reduction.

10 Fig. 17A is a bar graph showing the effect of thioredoxin-linked reduction and pepsin on the allergenicity of giant ragweed pollen extract in dogs that are sensitive to disulfide bond containing proteins.

Fig. 17B is a bar graph showing the effect of thioredoxin-linked reduction and pepsin digestibility on the allergenicity of giant ragweed pollen extract in  
15 dogs less sensitive to disulfide bond containing proteins.

Fig. 18A is a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the effect of time on the digestion of ragweed pollen following thioredoxin-linked reduction that occurred at 4°C.

Fig. 18B is a photograph of a protein stained SDS polyacrylamide  
20 electrophoretic gel showing the effect of time on the digestion of ragweed pollen following thioredoxin-linked reduction at 37°C and 55°C.

Fig. 19A is a photograph of an SDS polyacrylamide electrophoretic gel showing the extent of DTT linked reduction and cystamine and GSSG derivitization and reoxidation of BLG by means of fluorescence.

Fig. 19B is a photograph of a protein stained SDS polyacrylamide  
5 electrophoretic gel showing the extent of DTT linked reduction and cystamine and GSSG derivitization and reoxidation of BLG.

Fig. 20 is a block graph showing the response of a moderately sensitive dog to BLG following reduction with DTT, derivatization with cystamine and GSSG and reoxidation.

10 Fig. 21A is a photograph of an SDS polyacrylamide electrophoretic gel showing the change in the redox state of BLG following reduction by NTS at 4°C and cystamine and GSSG derivitization and reoxidation by means of fluorescence.

Fig. 21B is a photograph of a protein stained SDS polyacrylamide  
15 electrophoretic gel showing the extent of DTT linked reduction and cystamine and GSSG derivitization and reoxidation.

Fig. 22A is a photograph of an SDS polyacrylamide electrophoretic gel showing the change in the redox state of BLG reduced by DTT and thioredoxin (Trx) at 22°C and 55°C by means of fluorescence.

20 Fig. 22B is a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the change in the redox state of BLG reduced by DTT and thioredoxin (Trx) at 22°C and 55°C.

Fig. 23A, 23B and 23C are photographs of protein stained SDS polyacrylamide electrophoretic gels showing the susceptibility of reduced, derivatized and reoxidized BLG to digestion by simulated gastric fluid.

5 Fig. 24 is a line graph showing the time course of digestion of reduced, derivatized and reoxidized BLG by simulated gastric fluid.

Fig. 25A is a photograph of an SDS polyacrylamide electrophoretic gel showing the change in the redox state of milk proteins following reduction by DTT at 55°C and cystamine and GSSG derivatization and reoxidation by means of fluorescence.

10 Fig. 25B is a photograph of a protein stained SDS polyacrylamide electrophoretic gel showing the change in the redox state of milk proteins following reduction by DTT at 55°C and cystamine and GSSG derivatization and reoxidation.

15 Fig. 26 shows panels of photographs A through F of protein stained SDS polyacrylamide electrophoretic gels showing stability of reduced, derivatized and reoxidized milk proteins in simulated gastric fluid.

Fig. 27 is a block graph showing the response of a moderately sensitive dog to milk following reduction with DTT, derivatization with cystamine and GSSG and reoxidation.

## 20 DETAILED DESCRIPTION OF THE INVENTION

In accordance with this detailed description, the following definitions and abbreviations apply:

CM - certain bread wheat  $\alpha$ -amylase inhibitors

- DSG - certain  $\alpha$ -amylase inhibitors isolated from durum wheat  
DTNB - 2'5'-dithiobis (2-nitrobenzoic acid)  
NTR - NADP-thioredoxin reductase  
mBBr - monobromobimane  
5 NADP-MDH - NADP-malate dehydrogenase  
FBPase - fructose-1,6-bisphosphatase  
SDS - sodium dodecyl sulfate  
DTT - dithiothreitol  
Cereal - millet, wheat, oat, barley, rice, sorghum, or rye  
10 BBTI - Bowman-Birk soybean trypsin inhibitor  
KTI - Kunitz soybean trypsin inhibitor  
PAGE - polyacrylamide gel electrophoresis  
TCA - trichloroacetic acid

### ENZYME INHIBITOR PROTEIN EXPERIMENTS

#### 15 Starting Materials

##### Materials

Seeds of bread wheat Triticum aestivum L, cv. Talent) and durum wheat (Triticum durum, Desf., cv. Mondur) were obtained from laboratory stocks.

##### Reagents

- 20 Chemicals and fine chemicals for enzyme assays and sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis were purchased from Sigma Chemical Co. and BioRad Laboratories, respectively. Monobromobimane (mBBr, tradename Thiolite) was purchased from Calbiochem. Other chemicals were obtained from commercial sources and were of the highest  
25 quality available.

##### Enzymes



Thioredoxin and NTR from *E. coli* were purchased from American Diagnostics, Inc. and were also isolated from cells transformed to overexpress each protein. The thioredoxin strain containing the recombinant p]asmid, pFP1, was kindly provided by Dr. J.-P. Jacquot (de la Motte-  
5 Guery, F. *et al.* (1991) *Eur. J. Biochem.* 196:287-294). The NTR strain containing the recombinant plasmid, pPMR21, was kindly provided by Drs. Marjorie Russel and Peter Model (Russel, M. *et al.* (1988) *J. Biol. Chem.* 263:9015-9019). The Isolation procedure used for these proteins was as described in those studies with the following changes: cells were broken in a  
10 Ribi cell fractionator at 25,000 psi and NTR was purified as described by Florencio *et al.* (Florencio, F. J. *et al.* (1988) *Arch. Biochem. Biophys.* 266:496-507) without the red agarose step. Thioredoxin and NTR from *Saccharomyces cerevisiae* (baker's yeast type 1) were isolated by the procedure developed by Florencio *et al.* for spinach leaves with the following  
15 changes: suspended cells [1 part cells:5 parts buffer (w/v)], were broken in a Ribi cell fractionator at 40,000 psi with three passes.

Thioredoxin h and NTR were isolated from wheat germ by the procedure developed for spinach leaves (Florencio, F.J. *et al.* (1988), *Arch. Biochem. Biophys.* 266:496-507). NADP-malate dehydrogenase (NADP-MDH) and  
20 fructose-1,6-bisphosphatase (FBPase) were purified from leaves of corn (Jacquot, J.-P. *et al.* (1981), *Plant Physiol.* 68:300-304) and spinach (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239) respectively. *E. coli* glutaredoxin and calf thymus thioredoxin were obtained from Professor A. Holmgren.

## 25 $\alpha$ -Amylase and trypsin inhibitors

CM-1 protein was isolated from the albumin-globulin fraction of bread wheat flour as described previously (Kobrehel, K. *et al.* (1991), *Cereal Chem.* 68:1-6). A published procedure was also used for the isolation of DSG

proteins (DSG-1 and DSG-2) from the glutenin fraction of durum wheat (Kobrehel, K. *et al.* (1989), *J. Sci. Food Agric.* 48:441-452). The CM-1, DSG-1 and DSG-2 proteins were homogeneous in SDS-polyacrylamide gel electrophoresis. Trypsin inhibitors were purchased from Sigma Chemical Co., except for the one from corn kernel which was from Fluca. In all cases, the commercial preparations showed a single protein component which migrated as expected in SDS-PAGE (Coomassie Blue stain), but in certain preparations, the band was not sharp.

#### **Other proteins**

Purothionin  $\alpha$  from bread wheat and purothionins  $\alpha$ -1 and  $\beta$  from durum wheat were kind gifts from Drs. D.D. Kasarda and B.L. Jones, respectively. The purothionin  $\alpha$  sample contained two members of the purothionin family when examined with SDS-polyacrylamide gel electrophoresis. The purothionin  $\alpha$ -1 and  $\beta$  samples were both homogeneous in SDS-polyacrylamide gel electrophoresis.

#### **Routine Method Steps**

##### **Enzyme activation assays**

The NADP-MDH, FBPase, NTR and Thioredoxin  $h$  assay methods were according to Florencio, F.J. *et al.* (1988), *Arch. Biochem. Biophys.* 266:496-507 with slight modifications as indicated. For enzyme activation assays, the preincubation time was 20 min unless specified otherwise.

##### **mBBR Fluorescent labeling and SDS-polyacrylamide gel electrophoresis analyses**

Direct reduction of the proteins under study was determined by a modification of the method of Crawford *et al.* (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239). The reaction was carried out in 100 mM potassium phosphate buffer, pH 7.1, containing 10 mM EDTA and 16% glycerol in a final volume of 0.1 ml. As indicated, 0.7  $\mu$ g

- (0.1  $\mu$ M) NTR and 1  $\mu$ g (0.8  $\mu$ M) thioredoxin (both routinely from *E. coli* were added to 70  $\mu$ l of the buffer solution containing 1 mM NADPH and 10  $\mu$ g (2 to 17  $\mu$ M) of target protein. When thioredoxin was reduced by dithiothreitol (DTT, 0.5 mM), NADPH and NTR were omitted. Assays with
- 5 reduced glutathione were performed similarly, but at a final concentration of 1 mM. After incubation for 20 min, 100 nmoles of mBBR were added and the reaction was continued for another 15 min. To stop the reaction and derivatize excess mBBR, 10  $\mu$ l of 10% SDS and 10  $\mu$ l of 100 mM  $\beta$ -mercaptoethanol were added and the samples were then applied to the gels.
- 10 In the case of reduction by glutaredoxin, the thioredoxin and NTR were replaced by 1  $\mu$ g (0.8  $\mu$ M) *E. coli* glutaredoxin, 1.4  $\mu$ g (0.14  $\mu$ M) glutathione reductase purified from spinach leaves (Florencio, F.J. *et al.* (1988), *Arch. Biochem. Biophys.* 266:496-507) and 1.5 mM NADPH was used.
- 15 Gels (17.5% w/v, 1.5 mm thickness) were prepared according to Laemmli (Laemmli, U.K. (1970), *Nature* 227:680-685) and developed for 16 hr. at constant current (9 mA). Following electrophoresis, gels were placed in a solution of 40% methanol and 10% acetic acid, and soaked for 4 to 6 hours with several changes of the solution. Gels were then examined for
- 20 fluorescent bands with near ultraviolet light and photographed (exposure time 25 sec) according to Crawford *et al.* (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239). Finally, gels were stained with Coomassie Blue and destained as before (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239).
- 25 **Quantification of labeled proteins**
- To obtain a quantitative indication of the extent of reduction of test proteins by the NADP/thioredoxin system, the intensities of their fluorescent bands seen in SDS-polyacrylamide gel electrophoresis were evaluated, using a

modification of the procedure of Crawford *et al.* (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239). The photographic negatives were scanned using a Pharmacia Ultrascan laser densitometer, and the area underneath the peaks was quantitated by comparison to a standard curve  
 5 determined for each protein. For the latter determination, each protein (at concentrations ranging from 1 to 5  $\mu$ g) was reduced by heating for 3 min at 100°C in the presence of 0.5 mM DTT. Labeling with mBBBr was then carried out as described above except that the standards were heated for 2 min at 100°C after the reaction was stopped with SDS and excess mBBBr  
 10 derivatized with  $\beta$ -mercaptoethanol. Because the intensity of the fluorescent bands was proportional to the amounts of added protein, it was assumed that reduction was complete under the conditions used.

## EXAMPLE 1

### Thioredoxin-linked Reduction of $\alpha$ -Amylase Inhibitors

#### 15 Enzyme Activation Assays

The capability to replace a specific thioredoxin in the activation of chloroplast enzymes is one test for the ability of thiol groups of a given protein to undergo reversible redox change. Even though not physiological in the case of extraplastidic proteins, this test has proved useful in several studies. A  
 20 case in point is purothionin which, when reduced by thioredoxin h activates chloroplast FBPase (Wada, K. *et al.* (1981), *FEBS Lett.* 124:237-240, and Johnson, T.C. *et al.* (1987), *Plant Physiol.* 85:446-451). The FBPase, whose physiological activator is thioredoxin f, is unaffected by thioredoxin h. In this Example, the ability of cystine-rich proteins to activate FBPase as well  
 25 as NADP-MDH was tested as set forth above. The  $\alpha$ -amylase inhibitors from durum wheat (DSG-1 and DSG-2) were found to be effective in enzyme activation; however, they differed from purothionin in showing a specificity for NADP-MDH rather than FBPase (Table I). The  $\alpha$ -amylase inhibitors

were active only in the presence of reduced thioredoxin h, which itself did not significantly activate NADP-MDH under these conditions. DSG-1 and DSG-2 activated NADP-malate dehydrogenase in the presence of DTT-reduced thioredoxin h according to the reaction sequence

5 (DTT → Thioredoxin → DSG → NADP-MDH).

The complete system for activation contained in 200  $\mu$ l of 100 mM Tris-HCl buffer, pH 7.9 was 10 mM DTT, 0.7  $\mu$ g corn leaf NADP-MDH, 0.25  $\mu$ g wheat thioredoxin h and 10  $\mu$ g of DSG-1 or DSG-2. In one study 20 mM  $\beta$ -mercaptoethanol ( $\beta$ -MET) was used instead of DTT. Following

10 preincubation, NADP-MDH was assayed spectrophotometrically.

In the enzyme activation assays, thioredoxin h was reduced by DTT; as expected, monothiols such as  $\beta$ -mercaptoethanol ( $\beta$ -MET), which do not reduce thioredoxin at a significant rate under these conditions (Jacquot, J.-P. *et al.* (1981), *Plant Physiol.* 68:300-304; Nishizawa, A.N. *et al.* (1982),




15 "Methods in Chloroplast Molecular Biology", (M. Edelman, R.B. Hallick and N.-H. Chua, eds.) pp. 707-714, Elsevier Biomedical Press, New York, and Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239), did not replace DTT.

NADP-MDH activity was proportional to the level of added DSG-1 and

20 DSG-2 at a constant thioredoxin h concentration. The same DTT formula was used as set forth above. Except for varying the DSG-1 or DSG-2 concentrations, conditions were identical to those previously described. When tested at a fixed DSG concentration, NADP-MDH showed enhanced activity with increasing thioredoxin h. Except for varying the thioredoxin h

25 concentration, conditions were as described above.

CM-1--the bread wheat protein that is similar to DSG proteins but has a lower molecular weight - also activated NADP-MDH and not FBPase when 20µg of CM-1 were used as shown in Table I. The results indicate that thioredoxin h reduces a variety of α-amylase inhibitors, which, in turn, activate NADP-MDH in accordance with equations 4-6. These proteins were ineffective in enzyme activation when DTT was added in the absence of thioredoxin.



**Table I**

Effectiveness of Thioredoxin-Reduced Trypsin Inhibitors,  
Thionins, and  $\alpha$ -Amylase Inhibitors in Activating Chloroplast  
NADP-Malate Dehydrogenase and Fructose Bisphosphatase  
(DTT-Thioredoxin-Indicated Protein-Target Enzyme)

Activation of NADPH-MDH was carried out as described above in this  
Example except that the quantity of DSG or the other proteins tested was 20  
 $\mu$ g. FBPase activation was tested using the standard DTT assay with 1  $\mu$ g of  
*E. coli* thioredoxin and 20  $\mu$ g of the indicated proteins. The above values are  
corrected for the limited activation seen with *E. coli* thioredoxin under these  
conditions.



	*ACTIVITY, nkat/mg Protein	$M_r$ , kDa	No. of		
			S-S Groups	NADP-MDH	FBPase
15	<u><math>\alpha</math>-Amylase Inhibitors</u>				
	**DSG-2	17	5	2	0
	**DSG-1	14	5	2	0
	‡CM-1	12	5	12	0
	<u>Trypsin Inhibitors</u>				
20	<u>Cystine-rich</u> (plant)				
	Corn kernel	12	5	5	0
	Soybean Bowman-Birk	8	7	3	0
	<u>Other types</u>				
	Ovomucoid	28	9	2	0
25	Soybean Kunitz	20	2	2	0
	Ovoinhibitor	49	14	1	0
	Bovine lung (Aprotinin)	7	3	Trace	2
	<u>Thionins</u>				
	**Purothionin- $\alpha_1$	6	4	1	39
30	**Purothionin- $\beta$	6	4	Trace	5
	‡Purothionin- $\alpha$	6	4	0	14

\* These values compare to the corresponding values of 40 and 550 obtained, respectively,  
with spinach chloroplast thioredoxin  $m_1$  (NADP-MDH) and thioredoxin  $f_2$   
35 \*\* From Durum wheat  
‡ From bread wheat

## EXAMPLE 2

### DTNB Reduction Assays

A second test for thiol redox activity is the ability to catalyze the reduction of the sulfhydryl reagent, 2',5'-dithiobis(2-nitrobenzoic acid) (DTNB),  
 5 measured by an increase in absorbance at 412 nm. Here, the protein assayed was reduced with NADPH via NTR and a thioredoxin. The DTNB assay proved to be effective for the  $\alpha$ -amylase inhibitors from both durum (DSG-1 and 2) and bread wheat (CM-1). When reduced by the NADP/thioredoxin system (in this case using NTR and thioredoxin from *E. coli*), either DSG-1  
 10 or DSG-2 markedly enhanced the reduction of DTNB (NADPH  $\rightarrow$  NTR  $\rightarrow$  Thioredoxin  $\rightarrow$  DSG  $\rightarrow$  DTNB). The DTNB reduction assay was carried out with 10  $\mu$ g thioredoxin and 10  $\mu$ g NTR and 20  $\mu$ g of DSG-1 or DSG-2. CM-1 was also effective in the DTNB reduction assay, and, as with NADP-MDH activation (Table I), was detectably more active  
 15 than the DSG proteins. The conditions for the CM-1 assay were the same as for the DSG/DTNB assay except that the DSG proteins were omitted and purothionin  $\alpha$ , 20  $\mu$ g or CM-1, 20  $\mu$ g was used). The results thus confirmed the enzyme activation experiments in Example 1 and showed that the  $\alpha$ -amylase inhibitors can be reduced physiologically by the  
 20 NADP/thioredoxin system. The role of the  $\alpha$ -amylase inhibitors in promoting the reduction of DTNB under these conditions is summarized in equations 7-9.





### EXAMPLE 3

#### Protein Reduction Measurements

The availability of monobromobimane (mBBr) and its adaptation for use in plant systems has given a new technique for measuring the sulfhydryl groups of plant proteins (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239). When coupled with SDS-polyacrylamide gel electrophoresis, mBBr can be used to quantitate the change in the sulfhydryl status of redox active proteins, even in complex mixtures. This technique was therefore applied to the inhibitor proteins to confirm their capacity for reduction by thioredoxin. Here, the test protein was reduced with thioredoxin which itself had been previously reduced with either DTT or NADPH and NTR. The mBBr derivative of the reduced protein was then prepared, separated from other components by SDS-polyacrylamide gel electrophoresis and its reduction state was examined by fluorescence. In the experiments described below, thioredoxin from *E. coli* was found to be effective in the reduction of each of the targeted proteins. Parallel experiments revealed that thioredoxin h and calf thymus thioredoxins reduced, respectively, the proteins from seed and animal sources.

In confirmation of the enzyme activation and dye reduction experiments, DSG-1 was effectively reduced in the presence of thioredoxin. Following incubation the proteins were derivatized with mBBr and fluorescence visualized after SDS-polyacrylamide gel electrophoresis. Reduction was much less with DTT alone and was insignificant with GSH. A similar requirement for thioredoxin was observed for the reduction of CM-1 and DSG-2 (data not shown). While the thioredoxin used was from *E. coli* similar results were obtained with wheat thioredoxin h. Thioredoxin was also required when DTT was replaced by NADPH and NTR (data not shown).

#### EXAMPLE 4

##### Thioredoxin-linked Reduction of Cystine-Rich Plant Trypsin Inhibitors

Whereas the major soluble cystine-rich proteins of wheat seeds can act as  
5 inhibitors of exogenous  $\alpha$ -amylases, the cystine-rich proteins of most other  
seeds lack this activity, and, in certain cases, act as specific inhibitors of  
trypsin from animal sources. While these proteins can be reduced with  
strong chemical reductants such as sodium borohydride (Birk, Y. (1985), *Int.*  
*J. Peptide Protein Res.* 25:113-131, and Birl, Y. (1976), *Meth. Enzymol.*  
10 45:695-7390), there is little evidence that they can be reduced under  
physiological conditions. It was therefore of interest to test trypsin inhibitors  
for the capacity to be reduced by thioredoxin. The cystine-rich  
representatives tested included the soybean Bowman-Birk and corn kernel  
trypsin inhibitors. The results in both cases were positive: each inhibitor  
15 activated NADP-MDH (but not FBPase) when added in the presence of DTT-  
reduced thioredoxin (Table I) and each reduced DTNB in the presence of  
NADPH, NTR and thioredoxin (data not shown).

As found for the  $\alpha$ -amylase inhibitors, the thioredoxin-dependent reduction of  
the cystine-rich trypsin inhibitors could be directly monitored by the  
20 mBBBr/SDS-polyacrylamide gel electrophoresis technique. Thus, significant  
reduction by DTT was observed only in the presence of reduced thioredoxin  
with both the Bowman-Birk (BBTI) inhibitor which showed a highly  
fluorescent fast moving band and corn kernel (CKTI) trypsin inhibitor which  
showed a highly fluorescent band migrating behind thioredoxin.

## EXAMPLE 5

### Thioredoxin-linked Reduction of Other Trypsin Inhibitors and Purothionins

In view of the finding that cystine-rich trypsin inhibitors from seeds can  
5 undergo specific reduction by thioredoxin, the question arose as to whether  
other types of trypsin inhibitor proteins share this property. In the course of  
this study, several such inhibitors -- soybean Kunitz, bovine lung aprotinin,  
egg white ovoinhibitor and ovomucoid trypsin inhibitors -- were tested.  
While the parameters tested were not as extensive as with the cystine-rich  
10 proteins described above, it was found that the other trypsin inhibitors also  
showed a capacity to be reduced specifically by thioredoxin as measured by  
both the enzyme activation and mBBR/SDS-polyacrylamide gel  
electrophoresis methods. As was the case for the cystine-rich proteins  
described above, the trypsin inhibitors tested in this phase of the study  
15 (soybean Kunitz and animal trypsin inhibitors) activated NADP-MDH but not  
FBPase (Table I). Bovine lung aprotinin was an exception in that it activated  
FBPase more effectively than NADP-MDH. It might also be noted that  
aprotinin resembles certain of the seed proteins studied here in that it shows a  
high content of cystine (ca. 10%) (Kassel, B. *et al.* (1965), *Biochem.*  
20 *Biophys. Res. Commun.* 20:463-468).

The fluorescence evidence for the thioredoxin-linked reduction of one of  
these proteins, the Kunitz inhibitor, was shown by a highly fluorescent slow  
moving band in an mBBR/SDS-polyacrylamide electrophoretic gel. In its  
reduced form, the Kunitz inhibitor also yielded a fluorescent fast moving  
25 band. The nature of this lower molecular mass species is not known. Its  
position on the gel suggests that it could represent Bowman-Birk inhibitor  
present as a contaminant in the Kunitz preparation; however, such a  
component was not evident in Coomassie Blue stained SDS gels. The animal  
inhibitors which yielded a single fluorescent band of the expected molecular

weight, also showed a thioredoxin requirement for reduction (data not shown).

In confirmation of earlier results, thioredoxin-reduced purothionin consistently activated FBPase and the type tested earlier, purothionin- $\alpha$ ,  
5 failed to activate NADP-MDH (Table I) (Wada, K. *et al.* (1981), *FEBS Lett.* 124:237-240). However, in contrast to purothionin- $\alpha$  from bread wheat, two purothionins previously not examined (purothionins  $\alpha$ -1 and  $\beta$  from durum wheat) detectably activated NADP-MDH (Table I). The two durum wheat purothionins also differed in their ability to activate FBPase. The activity  
10 differences between these purothionins were unexpected in view of the strong similarity in their amino acid sequences (Jones, B.L. *et al.* (1977), *Cereal Chem.* 54:511-523) and in their ability to undergo reduction by thioredoxin. A requirement for thioredoxin was observed for the reduction of purothionin (here the  $\alpha$ -type) by the SDS-PAGE fluorescence procedure.

15

## EXAMPLE 6

### Quantitation of Reduction

The above Examples demonstrate that thioredoxin reduces a variety of proteins, including  $\alpha$ -amylase, such as the CM and DSG inhibitors, and trypsin inhibitors from seed as well as animal sources. While clear in the  
20 qualitative sense, the above results give no quantitative indication of the extent of reduction. Therefore, an experiment was conducted following the protocol of Crawford *et al.* (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239).

As shown in Table II, the extent of reduction of the seed inhibitor proteins by  
25 the *E. coli* NADP/thioredoxin system was time-dependent and reached, depending on the protein, 15 to 48% reduction after two hours. The results, based on fluorescence emitted by the major protein component, indicate that

thioredoxin acts catalytically in the reduction of the  $\alpha$ -amylase and trypsin inhibitors. The ratio of protein reduced after two hours to thioredoxin added was greater than one for both the most highly reduced protein (soybean Bowman-Birk trypsin inhibitor) and the least reduced protein (corn kernel trypsin inhibitor) -- *i.e.*, respective ratios of 7 and 2 after a two-hour reduction period. It should be noted that the values in Table II were obtained under standard assay conditions and no attempt was made to optimize reduction by modifying those conditions.

**Table II**

10      Extent of Reduction of Seed Proteins by the NADP/Thioredoxin System  
           Using the mBBr/SDS-Polyacrylamide Gel Electrophoresis Procedure

The following concentrations of proteins were used (nmoles): thioredoxin, 0.08; NTR, 0.01; purothionin- $\beta$ , 1.7; DSG-1, 0.7; corn kernel trypsin inhibitor, 1.0; Bowman-Birk trypsin inhibitor, 1.3; and Kunitz trypsin  
 15      inhibitor, 0.5. Except for the indicated time difference, other conditions were as in Examples 1-4.

---

	<u>Protein</u>	<u>20 min</u>	<u>% Reduction After</u> <u>120 min</u>
20	Purothionin- $\beta$	15	32
	DSG-1	22	38
	Corn kernel trypsin inhibitor	3	15
	Bowman-Birk trypsin inhibitor	25	48
	Kunitz trypsin inhibitor	14	22



---

25

## EXAMPLE 7

### *E. coli* Glutaredoxin as Reductant

Bacteria and animals are known to contain a thiol redox protein, glutaredoxin, that can replace thioredoxin in reactions such as ribonucleotide  
5 reduction (Holmgren, A. (1985), *Annu. Rev. Biochem.* 54:237-271).  
Glutaredoxin is reduced as shown in equations 10 and 11.



So far there is no evidence that glutaredoxin interacts with proteins from  
10 higher plants. This ability was tested, using glutaredoxin from *E. coli* and  
the seed proteins currently under study. Reduction activity was monitored by  
the mBBR/SDS polyacrylamide gel electrophoresis procedure coupled with  
densitometric scanning. It was observed that, under the conditions previously  
described, glutaredoxin could effectively replace thioredoxin in some, but not  
15 all cases. Thus, glutaredoxin was found to be active in the reduction of the  
following (the numbers indicate the percentage reduction relative to *E. coli*  
thioredoxin): DSG-1 and CM-1  $\alpha$ -amylase inhibitors (147 and 210%,  
respectively); corn kernel trypsin inhibitor (424%); and purothionin (82, 133,  
and 120% for the  $\alpha$ ,  $\alpha 1$  and  $\beta$  forms, respectively). Glutaredoxin was  
20 ineffective in the reduction of the DSG-2  $\alpha$ -amylase inhibitor and the soybean  
Bowman-Birk and Kunitz trypsin inhibitors. The trypsin inhibitors from  
animal sources also showed a mixed response to glutaredoxin. Egg white  
ovoinhibitor was effectively reduced (55% reduction relative to *E. coli*  
thioredoxin) whereas egg white ovomucoid inhibitor and bovine lung  
25 aprotinin were not affected. Significantly, as previously reported (Wolosiuk,  
R.A. *et al.* (1977), *Nature* 266:565-567), glutaredoxin failed to replace

thioredoxin as the immediate reductant in the activation of thioredoxin-linked enzymes of chloroplasts, FBPase and NADP-MDH (data not shown).

The above Examples demonstrate that some of the enzyme inhibitor proteins tested can be reduced by glutaredoxin as well as thioredoxin. Those specific  
5 for thioredoxin include an  $\alpha$ -amylase inhibitor (DSG-2), and several trypsin inhibitors (Kunitz, Bowman-Birk, aprotinin, and ovomucoid inhibitor). Those proteins that were reduced by either thioredoxin or glutaredoxin include the purothionins, two  $\alpha$ -amylase inhibitors (DSG-1, CM-1), a  
10 cystine-rich trypsin inhibitor from plants (corn kernel inhibitor), and a trypsin inhibitor from animals (egg white ovoidinhibitor). These results raise the question of whether glutaredoxin occurs in plants. Glutaredoxin was reported to be present in a green alga (Tsang, M.L.-S. (1981), *Plant Physiol.* 68:1098-1104) but not in higher plants.

Although the activities of the NADP-MDH and FBPase target enzymes  
15 shown in Table I are low relative to those seen following activation by the physiological chloroplast proteins (thioredoxin m or f), the values shown were found repeatedly and therefore are considered to be real. It seems possible that the enzyme specificity shown by the inhibitor proteins, although not relevant physiologically, reflects a particular structure achieved on  
20 reduction. It remains to be seen whether such a reduced structure is related to function within the seed or animal cell.

The physiological consequence of the thioredoxin (or glutaredoxin) linked reduction event is of considerable interest as the function of the targeted proteins is unclear. The present results offer a new possibility. The finding  
25 that thioredoxin reduces a wide variety of inhibitor proteins under physiological conditions suggests that, in the absence of compartmental barriers, reduction can take place within the cell.

### EXAMPLE 8

#### Inactivation of Soybean Trypsin Inhibitor in Soybean Meal

The goal of this Example is to inactivate the Bowman-Birk and Kunitz trypsin inhibitors of soybeans. The following protocol applies to animal feed preparations. To 10 g of soybean meal are added 0.2  $\mu$ g thioredoxin, 0.1  $\mu$ g NADP-thioredoxin reductase and 500 nanamoles NADPH together with 1 M Tris-HCl buffer, pH 7.9, to give 5.25 ml of 30 mM Tris-HCl. The above mixture is allowed to sit for about 30 min at room temperature. Direct reduction of the soybean trypsin inhibitor is determined using the mBBr fluorescent labeling/SDS-polyacrylamide gel electrophoresis method previously described (Kobrehel, K. *et al.* (1991), *J. Biol. Chem.* 266:16135-16140). An analysis of the ability of the treated flour for trypsin activity is made using modifications of the insulin and BAEE (Na-benzoyl-L-arginine ethyl ester) assays (Schoellmann, G. *et al.* (1963), *Biochemistry* 252:1963; Gonias, S.L. *et al.* (1983), *J. Biol. Chem.* 258:14682). From this analysis it is determined that soybean meal so treated with the NADP/thioredoxin system does not inhibit trypsin.

### EXAMPLE 9

#### Inactivation of $\alpha$ -Amylase Inhibitors in Cereals

To 10 g of barley malt are added 0.2  $\mu$ g thioredoxin, 0.1  $\mu$ g NADP-thioredoxin reductase and 500 nanamoles NADPH together with 1 M Tris-HCl buffer, pH 7.9, to give 5.25 ml of 30 mM Tris-HCl. The above mixture is allowed to sit for about 30 min at room temperature. Direct reduction of the  $\alpha$ -amylase inhibitors is determined using the mBBr fluorescent labeling/SDS-polyacrylamide gel electrophoresis method previously described (Kobrehel, K. *et al.* (1991), *J. Biol. Chem.* 266:16135-16140).  $\alpha$ -Amylase activity is monitored by following the release of maltose



from starch (Bernfeld, P. (1955), *Methods in Enzymol.* 1:149). From this analysis it is determined that barley so treated with the NADP/thioredoxin system does not inhibit  $\alpha$ -amylase.

## REDUCTION OF CEREAL PROTEINS

5

### Materials and Methods

#### Plant Material

Seeds and semolina of durum wheat (*Triticum durum*, Desf. cv. Monroe) were kind gifts of Dr. K. Kahn.

#### Germination of wheat seeds

- 10 Twenty to 30 seeds were placed in a plastic Petri dish on three layers of Whatman #1 filter paper moistened with 5 ml of deionized water. Germination was carried out for up to 4 days at room temperature in a dark chamber.

#### Reagents/Fine Chemicals

- 15 Biochemicals and lyophilized coupling enzymes were obtained from Sigma Chemical Co. (St. Louis, MO). *E. coli* thioredoxin and NTR were purchased from American Diagnostica, Inc. (Greenwich, CT). Wheat thioredoxin h and NTR were isolated from germ, following the procedures developed for spinach leaves (Florencio, F.J. *et al.* (1988), *Arch. Biochem.*
- 20 *Biophys.* 266:496-507). *E. coli* glutaredoxin was a kind gift of Professor A. Holmgren. Reagents for SDS-polyacrylamide gel electrophoresis were purchased from Bio-Rad Laboratories (Richmond, CA). Monobromobimane (mBBBr) or Thiolite was obtained from Calbiochem Co. (San Diego, CA). Aluminum lactate and methyl green were products of Fluka Chemicals Co.
- 25 (Buchs, Switzerland).

### **Gliadins and Glutenins**

For isolation of insoluble storage proteins, semolina (0.2 g) was extracted sequentially with 1 ml of the following solutions for the indicated times at 25°C: (1) 50 mM Tris-HCl, pH 7.5 (20 min); (2) 70% ethanol (2 hr); and  
5 (3) 0.1 M acetic acid (2 hr). During extraction, samples were placed on an electrical rotator and, in addition, occasionally agitated with a vortex mixer. After extraction with each solvent, samples were centrifuged (12,000 rpm for 5 min) in an Eppendorf microfuge and, supernatant fractions were saved for analysis. In between each extraction, pellets were washed with 1 ml of  
10 water, collected by centrifugation as before and the supernatant wash fractions were discarded. By convention, the fractions are designated: (1) albumin/globulin; (2) gliadin; and (3) glutenin.

### ***In vitro* mBBr labelling of proteins**

Reactions were carried out in 100 mM Tris-HCl buffer, pH 7.9. As  
15 indicated, 0.7 µg NTR and 1 µg thioredoxin (both from *E. coli* unless specified otherwise) were added to 70 µl of this buffer containing 1 mM NADPH and 10 µg of target protein. When thioredoxin was reduced by dithiothreitol (DTT), NADPH and NTR were omitted and DTT was added to 0.5 mM. Assays with reduced glutathione were performed similarly, but at a  
20 final concentration of 1 mM. After incubation for 20 min, 100 nmoles of mBBr were added and the reaction was continued for another 15 min. To stop the reaction and derivatize excess mBBr, 10 µl of 10% SDS and 10 µl of 100 mM β-mercaptoethanol were added and the samples were then applied to the gels. For reduction by glutaredoxin, the thioredoxin and NTR were  
25 replaced by 1 µg *E. coli* glutaredoxin, 1.4 µg glutathione reductase (purified from spinach leaves) and 1.5 mM NADPH.

### ***In vivo* mBBR labelling of proteins**

- At the indicated times, the dry seeds or germinating seedlings (selected on the basis of similar radical length) were removed from the Petri dish and their embryos or germinated axes were removed. Five endosperms from each lot
- 5 were weighed and then ground in liquid N<sub>2</sub> with a mortar and pestle. One ml of 2.0 mM mBBR in 100 mM Tris-HCl, pH 7.9, buffer was added just as the last trace of liquid N<sub>2</sub> disappeared. The thawed mixture was then ground for another minute and transferred to a microfuge tube. The volume of the suspension was adjusted to 1 ml with the appropriate mBBR or buffer
- 10 solution. Protein fractions of albumin/globulin, gliadin and glutenin were extracted from endosperm of germinated seedlings as described above. The extracted protein fractions were stored at -20°C until use. A buffer control was included for each time point.

### **SDS-Polyacrylamide Gel Electrophoresis**

- 15 SDS-polyacrylamide electrophoresis of the mBBR-derivatized samples was performed in 15 % gels at pH 8.5 as described by Laemmli, U.K. (1970), *Nature* 227:680-685. Gels of 1.5 mm thickness were developed for 16 hr. at a constant current of 9 mA.

### **Native Gel Electrophoresis**

- 20 To resolve the different types of gliadins, native polyacrylamide gel electrophoresis was performed in 6% gels (a procedure designed to separate gliadins into the four types) as described by Bushuk and Zillman (Bushuk, W. *et al.* (1978), *Can. J. Plant Sci.* 58:505-515) and modified for vertical slab gels by Sapirstein and Bushuk (Sapirstein, H.D. *et al.* (1985), *Cereal*
- 25 *Chem.* 62:372-377). A gel solution in 100 ml final volume contained 6.0 g acrylamide, 0.3 g bisacrylamide, 0.024 g ascorbic acid, 0.2 mg ferrous sulfate heptahydrate and 0.25 g aluminum lactate. The pH was adjusted to 3.1 with lactic acid. The gel solution was degassed for 2 hr. on ice and then

0.5 ml of 3 % hydrogen peroxide was added as a polymerization catalyst. The running buffer, also adjusted to pH 3.1 with lactic acid, contained 0.5 g aluminum lactate per liter. Duration of electrophoresis was approximately 4 hr., with a constant current of 50 mA. Electrophoresis was terminated when  
5 the solvent front, marked with methyl green tracking dye, migrated to about 1 cm from the end of the gel.

#### **mBBR Removal/Fluorescence Photography**

Following electrophoresis, gels were placed in 12 % (w/v) trichloroacetic acid and soaked for 4 to 6 hr. with one change of solution to fix the proteins; gels  
10 were then transferred to a solution of 40 % methanol/10 % acetic acid for 8 to 10 hr. to remove excess mBBR. The fluorescence of mBBR, both free and protein bound, was visualized by placing gels on a light box fitted with an ultraviolet light source (365 nm). Following removal of the excess (free)  
15 mBBR, gels were photographed with Polaroid Positive/Negative Landfilm, type 55, through a yellow Wratten gelatin filter No. 8 (cutoff=460 nm) (exposure time ranged from 25 to 60 sec at f4.5) (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239).

#### **Protein Staining/Destaining/Photography**

SDS-gels were stained with Coomassie Brilliant Blue R-250 in 40 %  
20 methanol/10 % acetic acid for 1 to 2 hr. and destained overnight as described before (Crawford, N.A. *et al.* (1989), *Arch. Biochem. Biophys.* 271:223-239). Aluminum lactate native gels were stained overnight in a filtered solution containing 0.1 g Coomassie Brilliant Blue R-250 (dissolved in 10 ml 95 % ethanol) in 240 ml 12 % trichloroacetic acid. Gels were destained  
25 overnight in 12 % trichloroacetic acid (Bushuk, W. *et al.* (1978), *Can. J. Plant Sci.* 58:505-515, and Sapirstein, H.D. *et al.* (1985), *Cereal Chem.* 62:372-377).

Protein stained gels were photographed with Polaroid type 55 film to produce prints and negatives. Prints were used to determine band migration distances and loading efficiency.

5 The Polaroid negatives of fluorescent gels and prints of wet protein stained gels were scanned with a laser densitometer (Pharmacia-LKB UltroScan XL). Fluorescence was quantified by evaluating peak areas after integration with GelScan XL software.

### Enzyme Assays

10 The following activities were determined in crude extracts with previously described methods: hexokinase (Baldus, B. *et al.* (1981), *Phytochem.* 20:1811-1814), glucose-6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase (Schnarrenberger, C. *et al.* (1973), *Arch. Biochem. Biophys.* 154:438-448), glutathione reductase, NTR and thioredoxin h (Florencio, F.J. *et al.* (1988), *Arch. Biochem. Biophys.* 266:496-507).

### 15 Protein Determination

Protein concentrations were determined by the Bradford method (Bradford, M. (1976) *Anal. Biochem.* 72:248-256), with Bio-Rad reagent and bovine serum albumin as a standard.

### Subunit Molecular Weight Determination

20 The subunit molecular weight of gliadins and glutenins was estimated on SDS-PAGE gels by using two sets of molecular weight standards (kDa). The first set consisted of BSA (66), ovalbumin (45), soybean trypsin inhibitor (20.1), myoglobin (17), cytochrome c (12.4) and aprotinin (6.5). The other set was the BioRad Prestained Low SDS-PAGE standards: phosphorylase b  
25 (110), BSA (84), ovalbumin (47), carbonic anhydrase (33), soybean trypsin inhibitor (24) and lysozyme (16).

## EXAMPLE 10

### Reduction of Gliadins

As a result of the pioneering contributions of Osborne and coworkers a century ago, seed proteins can be fractionated on the basis of their solubility  
5 in aqueous and organic solvents (20). In the case of wheat, preparations of endosperm (flour or semolina) are historically sequentially extracted with four solutions to yield the indicated protein fraction: (i) water, albumins; (ii) salt water, globulins; (iii) ethanol/water, gliadins; and (iv) acetic acid/water, glutenins. A wide body of evidence has shown that different proteins are  
10 enriched in each fraction. For example, the albumin and globulin fractions contain numerous enzymes, and the gliadin and glutenin fractions are in the storage proteins required for germination.

Examples 1, 4 and 5 above describe a number of water soluble seed proteins (albumins/globulins, *e.g.*,  $\alpha$ -amylase inhibitors, cystine-rich trypsin  
15 inhibitors, other trypsin inhibitors and thionines) that are reduced by the NADP/thioredoxin system, derived either from the seed itself or *E. coli*. The ability of the system to reduce insoluble storage proteins from wheat seeds, viz., representatives of the gliadin and glutenin fractions, is described below. Following incubation with the indicated additions, the gliadin  
20 proteins were derivatized with mBBBr and fluorescence was visualized after SDS-polyacrylamide gel electrophoresis. The lanes in this first gliadin gel were as follows: 1. Control: no addition. 2. GSH/GR/NADPH: reduced glutathione, glutathione reductase (from spinach leaves) and NADPH. 3. NGS: NADPH, reduced glutathione, glutathione reductase (from spinach  
25 leaves) and glutaredoxin (from *E. coli*). 4. NTS: NADPH, NTR, and thioredoxin (both proteins from *E. coli*). 5. MET/T(Ec):  $\beta$ -mercaptoethanol and thioredoxin (*E. coli*). 6. DTT. 7. DTT/T(Ec): DTT and thioredoxin (*E. coli*). 8. DTT/T(W): Same as 7 except with wheat thioredoxin h. 9. NGS,-

Gliadin: same as 3 except without the gliadin protein fraction. 10. NTS,-  
Gliadin: same as 4 except without the gliadin protein fraction. Based on its  
reactivity with mBBR, the gliadin fraction was extensively reduced by  
thioredoxin. The major members undergoing reduction showed a Mr ranging  
5 from 25 to 45 kDa. As seen in Examples 1, 4 and 5 with the seed  $\alpha$ -amylase  
and trypsin inhibitor proteins, the gliadins were reduced by either native  $h$  or  
*E. coli* type thioredoxin (both homogeneous); NADPH (and NTR) or DTT  
could serve as the reductant for thioredoxin. Much less extensive reduction  
was observed with glutathione and glutaredoxin--a protein able to replace  
10 thioredoxin in certain *E. coli* and mammalian enzyme systems, but not known  
to occur in higher plants.

The gliadin fraction is made up of four different protein types, designated  $\alpha$ ,  
 $\beta$ ,  $\gamma$  and  $\omega$ , which can be separated by native polyacrylamide gel  
electrophoresis under acidic conditions (Bushuk, W. *et al.* (1978), *Can. J.*  
15 *Plant Sci.* 58:505-515; Kasarda, D.D. *et al.* (1976), *Adv. Cer. Sci. Tech.*  
*1*:158-236; Sapirstein, H.D. *et al.* (1985), *Cereal Chem.* 62:372-377; and  
Tatham, A.S. *et al.* (1990), *Adv. Cer. Sci. Tech.* 10:1-78). Except for the  $\omega$   
gliadins, each species contains cystine (S-S) groups and thus has the potential  
for reduction by thioredoxin. In this study, following incubation with the  
20 indicated additions, proteins were derivatized with mBBR, and fluorescence  
was visualized after acidic-polyacrylamide gel electrophoresis. The lanes in  
the second gliadin gel in this study were as follows: 1. Control: no addition.  
2. GSH: reduced glutathione. 3. GSH/GR/NADPH: reduced glutathione,  
glutathione reductase (from spinach leaves) and NADPH. 4. NGS: NADPH,  
25 reduced glutathione, glutathione reductase (from spinach leaves) and  
glutaredoxin (from *E. coli*). 5. NGS+NTS: combination of 4 and 6. 6.  
NTS: NADPH, NTR, and thioredoxin (both proteins from *E. coli*).  
7. MET/T(Ec):  $\beta$ -mercaptoethanol and thioredoxin (*E. coli*). 8. DTT/T(Ec):  
DTT and thioredoxin (*E. coli*). 9. NTS(-T): same as 6 except without

thioredoxin. 10. NGS+NTS,-Gliadin: same as 5 except without the gliadin fraction.

- When the thioredoxin-reduced gliadin fraction was subjected to native gel electrophoresis, the proteins found to be most specifically reduced by thioredoxin were recovered in the  $\alpha$  fraction. There was active reduction of the  $\beta$  and  $\gamma$  gliadins, but as evident from the densitometer results summarized in Table III, the reduction within these groups was nonspecific, *i.e.*, relatively high levels of reduction were also achieved with glutathione and glutaredoxin. There was especially strong reduction of the  $\gamma$  gliadins by DTT-reduced thioredoxin. As anticipated, there was no reduction of the  $\omega$  gliadins. The evidence indicates that thioredoxin (either native h or *E. coli*) specifically reduces certain of the gliadins, especially the  $\alpha$  type.

**Table III**

**Reductant Specificity of the Different Types of Gliadins**

- 15 The area under  $\alpha$ ,  $\beta$ ,  $\gamma$  and aggregate peaks following reduction by the NADP/thioredoxin system were: 4.33, 8.60, 5.67 and 0.74 Absorbance units times millimeters, respectively. These combined areas were about 65 % of those observed when thioredoxin was reduced by DTT with the second gel, with the reaction conditions as in Example 10.

20	<u>Reductant</u>	<u>Gliadin, % Relative Reduction</u>			
		$\alpha$	$\beta$	$\gamma$	Aggregate*
	None	22.4	30.4	24.3	29.2
	Glutathione	36.4	68.1	60.6	60.1
	Glutaredoxin	43.5	83.3	79.7	61.5
25	Thioredoxin	100.0	100.0	100.0	100.0

---

\*Proteins not entering the gel



## EXAMPLE 11

### Reduction of Glutenins

The remaining group of seed proteins to be tested for a response to thioredoxin -- the glutenins -- while the least water soluble, are perhaps of greatest interest. The glutenins have attracted attention over the years because of their importance for the cooking quality of flour and semolina (MacRitchie, F. *et al.* (1990), *Adv. Cer. Sci. Tech.* 10:79-145). Testing the capability of thioredoxin to reduce the proteins of this group was, therefore, a primary goal of the current investigation.

Several glutenins were reduced specifically by thioredoxin when the mBBBr/SDS-page technique was applied and the conditions were as in Example 10 with the first gel. The most extensive reduction was observed in the low molecular mass range (30 - 55 kDa). The reduction observed in the higher molecular mass range was less pronounced, but still obvious-- especially in the 100 kDa region and above. Though not shown reduction may also occur in the 130 kDa range. Like the gliadins, certain of the glutenins were appreciably reduced by glutathione and glutaredoxin. However, in all cases, reduction was greater with thioredoxin and, in some cases, specific to thioredoxin (Table IV, note proteins in the 30 - 40 and 60 - 110 kDa range). As observed with the other wheat proteins tested, both the native h anal *E. coli* thioredoxins were active and could be reduced with either NADPH and the corresponding NTR or with DTT. Thus as found for the gliadins, certain glutenins were reduced *in vitro* specifically by thioredoxin, whereas others were also reduced, albeit less effectively, by glutathione and glutaredoxin.

**Table IV**  
**Reductant Specificity of Glutenins**

Reaction conditions as in the Example 1 study of the effect of thioredoxin h concentration on the activation of NADP-MDH by DSG-1 or -2  $\alpha$ -amylase inhibitors.

<u>Reductant</u>	<u>Glutenin, % Relative Reduction*</u>		
	60-110 kDa	40-60 kDa	30-40 kDa
None	8	23	16
Glutathione	31	51	29
10 Glutaredoxin	50	72	40
Thioredoxin*	100	100	100

---

\*Area under the three molecular weight classes (from high to low) following reduction by the NADP/thioredoxin system were: 1.5, 5.67 and 5.04  
15 Absorbance units times millimeters, respectively.

## EXAMPLE 12

### *In vivo* Reduction Experiments

The above Example demonstrates that thioredoxin specifically reduces components of the wheat gliadin and glutenin fractions when tested *in vitro*.

20 The results, however, provide no indication as to whether these proteins are reduced *in vivo* during germination -- a question that, to our knowledge, had not been previously addressed (Shutov, A.D. *et al.* (1987), *Phytochem.* 26:1557-1566).

To answer this question, the mBBR/SDS-PAGE technique was applied to  
25 monitor the reduction status of proteins in the germinating seed. We observed that reduction of components in the Osborne fractions increased progressively with time and reached a peak after 2 to 3 days germination. The observed increase in reduction ranged from 2-fold with the gliadins, to

3-fold with the albumin/globulins and 5-fold with the glutenins. The results suggest that, while representatives of the major wheat protein groups were reduced during germination, the net redox change was greatest with the glutenins.

- 5 Although providing new evidence that the seed storage proteins undergo reduction during germination, the results give no indication as to how reduction is accomplished, *i.e.*, by glutathione or thioredoxin. To gain information on this point, the *in vivo* reduction levels of the principal thioredoxin-linked gliadins (30 - 50 kDa) and glutenins (30 - 40, 40 - 60
- 10 kDa) was compared with the reduction determined from *in vitro* measurements (cf. Table IV). For this purpose, the ratio of fluorescence to Coomassie stained protein observed *in vivo* during germination and *in vitro* with the appropriate enzyme reduction system was calculated. The results (principal thioredoxin linked gliadins were those in the Mr range from 25 to
- 15 45 kDa, and glutenins were those in the Mr range from 30 to 60 kDa) suggest that, while glutathione could account for a significant part of the *in vivo* reduction of the gliadin fraction (up to 90%), this was not the case with the glutenins whose reduction seemed to require thioredoxin. The level of reduction that could be ascribed to glutathione (or glutaredoxin) was
- 20 insufficient to account for the levels of reduced glutenin measured in the germinating seed.

### EXAMPLE 13

#### Enzyme Measurements

- The source of NADPH needed for the NTR linked reduction of thioredoxin h
- 25 was also investigated. Semolina was analyzed for enzymes that function in the generation of NADPH in other systems, notably dehydrogenases of the oxidative phosphate pathway. The results summarized in Table V confirm earlier evidence that endosperm extracts contain the enzymes needed to

- generate NADPH from glucose via this pathway: hexokinase, glucose 6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase (Tatham, A.S. *et al.* (1990), *Adv. Cer. Sci. Tech.* 10:1-78). It is noteworthy that the glucose 6-phosphate dehydrogenase activity seen in Table V was insensitive to reduced thioredoxin (data not shown). In this respect the endosperm enzyme resembles its cytosolic rather than its chloroplast counterpart from leaves (Fickenscher, K. *et al.* (1986), *Arch. Biochem. Biophys.* 247:393-402; Buchanan, B.B. (1991), *Arch. Biochem. Biophys.* 288:1-9; Scheibe, R. *et al.* (1990), *Arch. Biochem. Biophys.* 274:290-297).
- 10 As anticipated from earlier results with flour (Johnson, T.C. *et al.* (1987), *Planta* 171:321-331; Suske, G. *et al.* (1979), *Z. Naturforsch. C* 34:214-221), semolina also contained thioredoxin h and NTR (Table V). Interestingly, based on activity measurements, NTR appeared to be a rate-limiting component in preparations from the cultivar examined.

15

**Table V**

Activities of Enzymes Effecting  
the Reduction of Thioredoxin h in Semolina  
(Glucose-Glu-6-P-6-P-Gluconate-NADP-Thioredoxin h)

20	<u>Protein</u>	Activity (nkat/mg protein)
	Hexokinase	0.28
	Glucose-6-P dehydrogenase	0.45
	6-P-Gluconate dehydrogenase	0.39
	NTR	0.06
25	Thioredoxin <u>h</u>	0.35

The present results suggest that thioredoxin h functions as a signal to enhance metabolic processes associated with the germination of wheat seeds.

Following its reduction by NTR and NADPH (generated via the oxidative

pentose phosphate pathway), thioredoxin h appears to function not only in the activation of enzymes, but also in the mobilization of storage proteins.

#### EXAMPLE 14

##### Improvement of Dough Quality

5 Dough quality was improved by reducing the flour proteins using the NADP/thioredoxin system. Reduced thioredoxin specifically breaks sulfur-sulfur bonds that cross-link different parts of a protein and stabilize its folded shape. When these cross-links are cut the protein can unfold and link up with other proteins in bread, creating an interlocking lattice that forms the elastic  
10 network of dough. The dough rises because the network traps carbon dioxide produced by yeast in the fermenting process. It is proposed that the reduced thioredoxin activated the gliadins and glutenins in flour letting them recombine in a way that strengthened the dough. Reduced thioredoxin strengthened the protein network formed during dough making. For these  
15 tests (using 10 gm of either intermediate quality wheat flour obtained from a local miller in Montpellier, France, or poor quality wheat also obtained from a local miller in Montpellier, France, this poor quality wheat being mainly of the Apollo cultivar), 0.2  $\mu\text{g}$  *E. coli* thioredoxin, 0.1  $\mu\text{g}$  *E. coli* NADP-thioredoxin reductase and 500 nanomoles NADPH were added together with  
20 1 M Tris-HCl, pH 7.9 buffer to give 5.25 ml of a 30 mM Tris-HCl enzyme system mixture. The reaction was carried out by mixing the enzyme system mixture with the 10 gm of the flour in a micro-farinograph at 30°C. The resulting farinograph measurements showed a strengthening of the dough by the added NADP/thioredoxin system. With a flour of poor quality, the  
25 farinograph reading was stable for at least 4 min after the dough was formed in the presence of the reduction system, whereas the reading dropped immediately after dough formation in the control without this addition. The improving effect was persistent and was maintained throughout the run. Expressed another way, the micro-farinograph reading is 375 Brabender

units, 7 min after dough formation with the poor quality wheat control (no added enzyme system) versus 450 Brabender units for the same poor quality wheat treated with components of the NADP/thioredoxin system (NADPH, thioredoxin and NADP-thioredoxin reductase).

- 5 Another farinograph study was carried out as above with 10 gm of Apollo flour only the concentration of NADPH was 500  $\mu$ moles instead of nanomoles. The farinograph measurements showed that this amount of NADPH also resulted in a definite improvement in the quality of the dough.

- Higher farinograph measurements of dough correspond to improved dough  
10 strength and improved baked good characteristics such as better crumb quality, improved texture and higher loaf volume. Also, based on *in vivo* analyses with the isolated proteins, the native wheat seed NADP/thioredoxin system will also be effective in strengthening the dough.

- For purposes of baking and other aspects of this invention, ranges of about  
15 0.1 to 3.0  $\mu$ g of a thioredoxin (preferably *E. coli* or thioredoxin h) and from about 0.1 to 2.0  $\mu$ g reductase and about 30 to 500 nanomoles of NADPH are added for about every 10 gm of flour. The optimal levels of thioredoxin and reductase depend on flour quality. In general, the higher the flour quality, the higher the level of thioredoxin and reductase required. Thioredoxin can  
20 also be reduced by lipoic acid instead of by the NADPH/NADP-thioredoxin reductase reduction system. The other dough ingredients such as milk or water are then added. However, the liquid may first be added to the NTR/thioredoxin system and then added to the flour. It is preferred that yeast for purposes of leavening be added after the reduced thioredoxin has  
25 had a chance to reduce the storage proteins. The dough is then treated as a regular dough proofed, shaped, etc. and baked.

NADPH can be replaced in this Example as well as in the following Examples with an NADPH generator such as one consisting of 100  $\mu$ M glucose 6-phosphate, 100  $\mu$ M NADP and 0.05 units (0.2  $\mu$ gram) glucose 6-phosphate dehydrogenase from a source such as yeast. The NADPH generator is added together with thioredoxin and NADP-thioredoxin reductase at the start of the dough making process.

A higher farinograph measurement was obtained when 10 gm of Apollo cultivar (CV) wheat were reacted with 20  $\mu$ l NADP (25 mM), 20  $\mu$ l G6P (25 mM), 0.25  $\mu$ g G6PDase, 0.1  $\mu$ g NTR and 0.2  $\mu$ g thioredoxin h contained in 4.25 ml H<sub>2</sub>O and 0.90 ml Tris-HCl (30 mM, pH 7.9). A higher farinograph measurement was also obtained when 10 gm of Apollo wheat were reacted with this same reaction mixture but without any NTR or thioredoxin.

## EXAMPLE 15

### Wheat Bread Baking Studies

The baking tests were carried out by using a computer monitored PANASONIC baking apparatus.

Composition of bread:

20	<u>Control:</u>	Flour*:	200 gm (dry)
		Water:	70% hydratation
		Salt (NaCl):	5.3 g
		Yeast:	4.8 g ( <i>Saccharomyces cerevisiae</i> , SafInstant) (dry yeast powder)

\* Flour samples were obtained from pure bread wheat cultivars having contrasting baking quality (including animal feed grade and other grades having from poor to good baking quality).

### Assays

The dough for the assays contained all the components of the control plus as indicated varying amounts of the NADP Thioredoxin System (NTS) and/or the NADP generating System.

### Experimental conditions

- Flour and salt are weighed and mixed
- The volume of water needed to reach a hydration of 70% was put into the baking pan.
- 5 -- The mixture of flour and salt was added to the water and the baking program monitored by the computer was started. The complete program lasted 3 hrs 9 min and 7 secs.
- In the case of the assays, enzyme system components are added to the water before the addition of the flour-salt mixture.
- 10 -- Yeast was added automatically after mixing for 20 min and 3 secs.

The program monitoring the Panasonic apparatus was:

	<u>Segments</u>	<u>Duration</u>	<u>Mixing Conditions</u>	<u>Heating</u>
	Mixing	00:00:03	T1	off
15	Mixing	00:05:00	T2	off
	Mixing	00:05:00	T1	off
	Rest	00:10:00	T0	off
	Mixing	00:17:00	T2	off
	Mixing	00:07:00	T1	off
20	Rest	00:30:00	T0	to reach 32°C
	Mixing	00:00:04	T1	32°C
	Rest	01:15:00	T0	32°C
	Baking	00:14:00	T0	to reach 180°C
	Baking	00:26:00	T0	180°C
25	Mixing Conditions: T0 = no mixing (motor at rest) T1 = normal mixing T2 = alternately 3 second mixing, 3 second rest			

Bread loaf volume was determined at the end of the baking, when bread loaves reached room temperature.



### Cultivar THESEE Assay

The french wheat cultivar Thesee is classified as having good breadmaking quality. Table VI below sets forth the results of the assay.

Table VI

5		NADPH (moles)	NTR ( $\mu$ g)	Th ( $\mu$ g)	Loaf Volume	
					Relative (cm <sup>3</sup> )	Units
	Control	0	0	0	1690	100
	Samples	6.0	30	60	1810	107
10		6.0	30	0	1725	102
		6.0	0	60	1720	102
		6.0	0	0	1550	92
		0	30	60	1800	107
15	*NADPH Generating syst.		30	60	1620	96
	*NADPH Generating syst. plus ATP, glucose		30	60	1630	96
20	NTR and Th from yeast	6.0	9.4	20	1750	104
*Composition of the NADPH generating system, ATP and glucose.						
25					Volume Added	
	NADP, 25 mMolar				700 $\mu$ l (17.5 $\mu$ moles)	
	Glucose-6-phosphate, 25 mMolar				700 $\mu$ l (17.5 $\mu$ moles)	
	Glucose-6-phosphate dehydrogenase (50 $\mu$ g/ml)				175 $\mu$ l (8.75 $\mu$ g)	
	ATP, 25 mMolar				700 $\mu$ l (17.5 $\mu$ moles)	
30	Glucose, 25 mMolar				700 $\mu$ l (17.5 $\mu$ moles)	

As shown in Table VI, an increased loaf volume was obtained when the complete NTS at concentrations of 6.0  $\mu$ moles NADPH, 30  $\mu$ g NTR and 60  $\mu$ g Th was used to bake loaves from 200 g of Thesee flour with the amounts and conditions described above in this Example. Unless otherwise stated, the

5 NTR and thioredoxin (th) were from *E. coli*. No similar increase occurred when the generating system was used or when either NTR or Th were omitted. Also no significant effect on loaf volume occurred when amounts of the components in the system were about half or less than half of the amounts of above.

#### 10 **Cultivar APOLLO Assay**

This French wheat cultivar is classified as having poor breadmaking quality. The NTR and thioredoxin used in this assay were from *E. coli*. Table VII below sets forth the results of this assay using 200 gm of Apollo flour. Again unless otherwise stated the amounts and conditions are those described

15 above at the beginning of the Example.

Table VII

		<u>Loaf Volume</u>				
		<u>NADPH</u>	<u>NTR</u>	<u>Th</u>	<u>Relative</u>	
		<u>(μmoles)</u>	<u>(μg)</u>	<u>(μg)</u>	<u>(cm3)</u>	<u>Units</u>
5	Control	0	0	0	1400	100
	Samples	6.0	30	60	1475	105
	*NADPH Generating syst. plus ATP, glucose		30	60	1530	109
10	*NADPH Generating syst. plus ATP, glucose		0	0	1430	102
	*NADPH Generating syst.		6	0	1430	102
15	*NADPH Generating syst.		6	7	1440	103

\*The composition of the generating system, ATP and glucose is as in Table VI.

## 20 Cultivar ARBON Assay

The French wheat cultivar Arbon is used for feed and is classified as non suitable for breadmaking. Tables VIII and IX below show that an improved bread loaf volume can be obtained from Arbon using the NTS or NADPH and NTR with the dough components and conditions described at the beginning of the Example. The amounts of NTR, thioredoxin, NADPH and the NADPH generating system components used in the assay are set forth in Tables VIII and IX. The improvement in Arbon bread quality using the complete NTS as set forth in Table IX was clearly visible when compared to the control.

Table VIII

		NADPH ( $\mu$ moles)	NTR ( $\mu$ g)	Th ( $\mu$ g)	<u>Loaf Volume</u> (cm <sup>3</sup> )
5	Control	0	0	0	1350
	Samples	0.1-0.6	3-4	3-4	up to 20% higher than the control
10		> 2.0	> 20	> 20	less than the control

Table IX

		<u>Loaf Volume</u> (cm <sup>3</sup> )	<u>Relative</u> <u>Units</u>
15	<u>Treatment</u>		
	Complete NTS	1650	122
	minus Thioredoxin	1690	125
	minus NTR	1520	113
	minus Thioredoxin, NTR	1540	114
20	minus NADPH	1440	107
	minus NADPH, plus *NADPH generating system	1560	116
	minus NTS (control)	1350	100
<hr/>			
25	NADPH, 0.6 $\mu$ moles Thioredoxin, 3.5 $\mu$ g NTR, 3 $\mu$ g		
	*Generating System:		
	3.5 $\mu$ moles NADP		
30	3.5 $\mu$ moles glucose-6-phosphate 1.75 $\mu$ g glucose-6-phosphate dehydrogenase		

## EXAMPLE 16

### Triticale Bread Baking Study

Triticale is a wheat/rye hybrid and is generally used for chicken feed. It is more nutritious than wheat but is not generally considered appropriate for breadmaking, especially in the more developed nations. The effect of the NTS system and variations thereof on loaves baked from Triticale flour was consequently studied. Unless otherwise stated, the baking conditions and dough ingredient were as described for wheat flour in Example 15. As shown in Table X there is an improvement in loaf volume when the triticale dough contained thioredoxin, NTR and the NADPH generating system in the amounts set forth in that Table. However, no corresponding improvement was seen when the NTS (*i.e.*, thioredoxin, NTR and NADPH) was used. An improvement in the texture of the bread, making it more cohesive and stable, also occurred when NTR, Th and the NADPH generating system as set forth in Table X were used.

**TABLE X**

Effect of the NADP/Thioredoxin System (NTS)  
on Loaves Baked from Triticale Flour (cv. Juan)

5	Treatment	Loaf Volume	
		(cm <sup>3</sup> )	Relative Units
	Complete NTS	1230	94
	minus NTS (control)	1310	100
10	minus NADPH, plus *NADPH	1390	106
<hr/>			
	NADPH, 0.6 $\mu$ moles		
	Thioredoxin, 3.5 $\mu$ g		
	NTR, 3.0 $\mu$ g		
15	Generating System:		
	4.5 $\mu$ moles NADP		
	4.5 $\mu$ moles glucose-6-phosphate		
	4.5 $\mu$ g glucose-6-phosphate dehydrogenase		

**EXAMPLE 17**

- 20 The effect of the NADPH/thioredoxin system on flour from sorghum, corn and rice was also determined. The baking conditions were as described for wheat flour in Example 15. The amounts of the components of the NTS as used in this assay were as follows: 8  $\mu$ moles NADPH, 40.5  $\mu$ g NTR and 54  $\mu$ g thioredoxin. Both the thioredoxin and NTR were from *E. coli*. The
- 25 breads in this study containing the NTS, especially corn and sorghum exhibited improved texture and stability.

### EXAMPLE 18

#### **Reduction of Ethanol-Soluble and Myristate-Soluble Storage Proteins from Triticale, Rye, Barley, Oat, Rice, Sorghum, Corn and Teff**

Unless otherwise stated, the materials and methods used in this Example are  
5 according to those set forth above in the section titled "Reduction of Cereal Proteins, Materials and Methods."

#### **Triticale, Rye, Barley, Oat and Teff**

The reactions were carried out in 30 mM Tris-HCl buffer, pH 7.9. As indicated, 0.7  $\mu$ g of NTR and 1  $\mu$ g of thioredoxin from *E. coli* or 2  $\mu$ g of  
10 thioredoxin from yeast, as identified, were added to 70  $\mu$ L of this buffer containing 1 mM NADPH and 25 to 30  $\mu$ g of extracted storage protein. The ethanol extracted storage proteins were obtained by using 50 ml of 70% ethanol for every 10 gm of flour and extracting for 2 hr. In the case of teff, 200 mg of ground seeds were extracted. The myristate extracted proteins  
15 were obtained by extracting 1 gm of flour with 8 mg sodium myristate in 5 ml of distilled H<sub>2</sub>O for 2 hrs. The combination of NADPH, NTR and thioredoxin is known as the NADP/thioredoxin system (NTS). As indicated, glutathione (GSH), 2.5 mM, was added as reductant in either the absence (GSH) or presence of 1.5 mM NADPH and 1.4  $\mu$ g of spinach leaf  
20 glutathione reductase (GR/GSH/NADPH). After incubation for 20 min, 100 nmol of mBBBr was added and the reaction was continued for another 15 min. To stop the reaction and derivatize excess mBBBr, 10  $\mu$ L of 10% SDS and 10  $\mu$ L of 100 mM 2-mercaptoethanol were added, and the samples were then applied to the gels. The procedure for SDS-polyacrylamide gel  
25 electrophoresis was as described by N. A. Crawford *et al.* (1989 *Arch. Biochem. Biophys.* 271:223-239).

### Rice, Sorghum and Corn

- The reactions were carried out in 30 mM Tris-HCl buffer, pH 7.9. When proteins were reduced by thioredoxin, the following were added to 70  $\mu$ L of buffer: 1.2 mM NADPH, 10 to 30  $\mu$ g of seed protein fraction, 0.5  $\mu$ g *E. coli* NTR and 1  $\mu$ g *E. coli* thioredoxin. For reduction with glutathione, thioredoxin and NTR were replaced with 2.5 mM reduced glutathione and 1  $\mu$ g glutathione reductase (baker's yeast, Sigma Chemical Co.). For reduction with dithiothreitol, NADPH, thioredoxin, and NTR were omitted and 0.5 mM dithiothreitol was added. In all cases, incubation time was 20 min.
- Then 10  $\mu$ l of a 10 mM mBBR solution was added and the reaction continued for an additional 15 min. To stop the reaction and derivatize excess mBBR, 10  $\mu$ l of 10% SDS and 10  $\mu$ l of 100 mM 2-mercaptoethanol were added and the samples applied to the gels. In each case, to obtain the extracted protein, 1 g ground seeds was extracted with 8 mg of sodium myristate in 5 ml distilled water. With the exception of the initial redox state determination of the proteins, samples were extracted for 2 hr at 22°C and then centrifuged 20 min at 16,000 rpm prior to the addition of the mBBR. With the initial redox state determination, the mBBR was added under a nitrogen atmosphere along with the myristate followed by extraction.
- Separate SDSS-polyacrylamide electrophoretic gels of the reduction studies of myristate-extracted proteins from flour of oat, triticale, rye, barley and teff were prepared. A gel showing the extent of thioredoxin linked buffer and ethanol-extracted proteins for teff was also prepared. In all of the oat, triticale, rye, barley, teff/myristate extractions studies, the flour was first extracted with buffer, 50 mM Tris-HCl, pH 7.5 for 20 min and then with 70% ethanol for 2 hr. In addition, gels were prepared for the myristate-extracted proteins from corn, sorghum and rice. With corn, sorghum and rice, the ground seeds were extracted only with myristate. Therefore, with corn, sorghum and rice, the myristate extract represents total protein,



whereas with oat, triticale, rye, barley and teff, the myristate extract represents only the glutenin-equivalent fractions since these flours had been previously extracted with buffer and ethanol. The results, depicted in the gels, show that the NTS is most effective, as compared to GSH or

5 GSH/GR/NADPH, with myristate-extracted (glutenin-equivalent) proteins from oat, triticale, rye, barley and teff. The NTS is also most effective with the total proteins from rice. Reduced glutathione is more effective with the total proteins from corn and sorghum.

### **Conclusions from the corn, sorghum and rice**

10 In the first gel relating to the effect of NTS vs. glutathione reductase on the reduction status of the myristate-extracted proteins, in treatment (1), extraction with myristate in the presence of mBBr was carried out under a nitrogen atmosphere; in treatment (2), to the myristate extracted proteins mBBr was added without prior reduction of the proteins; in treatment (3), the

15 myristate extracted proteins were reduced by the NADP/thioredoxin system (NTS); in treatment (4) the myristate extracted proteins were reduced by NADPH, glutathione and glutathione reductase. In the second gel relating to the *in vivo* reduction status and thioredoxin linked *in vitro* reduction of the myristate-extracted proteins, treatment (1) is like treatment (2) in the first

20 gel; in treatment (2) the seeds were extracted with myristate in the presence of mBBr under nitrogen; in treatment (3), seeds were extracted with myristate and reduced by the NTS and then mBBr was added; and in treatment (4) conditions as in (3) except that proteins were reduced by DTT. Treatment (1) in the first gel and treatment (2) in the second gel showed the initial redox

25 state of the proteins in the grains. For all three cereals, the proteins in the seed were highly reduced. If extracted in air, the proteins became oxidized especially the sorghum and rice. The oxidized proteins can be re-reduced, maximally with NTS in all cases. With rice, the reduction was relatively specific for thioredoxin; with corn, glutathione is as effective as thioredoxin

and with sorghum glutathione is slightly more effective than thioredoxin. Dithiothretol showed varying effectiveness as a reductant. These experiments demonstrated that the storage proteins of these cereals are less specific than in the case of wheat and suggest that thioredoxin should be tested both in the  
5 presence and absence of glutathione when attempting to construct a dough network.

Gels were also prepared resulting from the reduction studies of wheat glutenins and gliadins, respectively, by a yeast NADP/thioredoxin system. The glutenins were obtained by using 50 ml of 0.1 M acetic acid for every 10  
10 gm of flour and extracting for 2 hr. The gliadins were obtained by using 50 ml of 70% ethanol for every 10 gm of flour and extracting for 2 hr. The gels showed that the yeast system is highly active in reducing the two major groups of wheat storage proteins.

gels for the reduction of ethanol-extracted proteins from flour of triticale,  
15 rye, oat and barley, respectively, were also prepared. The results showed that the NTS is most effective with the ethanol-extracted proteins from triticale, rye and oat. The ethanol-extracted barley proteins were reduced in the control and thioredoxin or glutathione had little effect.

## EXAMPLE 19

### Effect of Thioredoxin-linked Reduction on the Activity and Stability of the Kunitz and Bowman-Birk Soybean Trypsin Inhibitor Proteins

#### Materials and Methods

##### 5 Plant materials

Durum wheat (*Triticum durum*, Desf. cv. Monroe) was a kind gift of Dr. K. Kahn. Wheat germ was obtained from Sigma Chemical Co. (St. Louis, MO).

##### Chemicals and Enzymes

Reagents for sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) were obtained from Bio-Rad Laboratories (Richmond, CA), and DTT was from Boehringer Mannheim Biochemicals (Indianapolis, IN). L-1-Tosylamide-2-phenylethyl chloromethyl ketone (TPCK)-treated trypsin (type XIII, T8640), subtilisin (type VIII: bacterial subtilisin Carlsberg, P538O), KTI (T9003), BBTI (T9777), azocasein, and other chemicals were  
15 purchased from Sigma Chemical Co. (St. Louis, MO). *E. coli* thioredoxin and NTR were isolated from cells transformed to overexpress each protein. The thioredoxin strain containing the recombinant plasmid, pFPI, was kindly provided by Dr. J.-P. Jacquot (de La Motte-Guery *et al.*, 1991). The NTR strain containing the recombinant plasmid, pPMR21, was kindly provided by  
20 Drs. Marjorie Russel and Peter Model (Russel and Model, 1988). The isolation procedures used for these proteins were as described in those studies with the following changes: cells were broken in a Ribi cell fractionator at 25,000 psi and NTR was purified as described by Florencio *et al.* (1988) without the red agarose step. The *E. coli* thioredoxin and NTR were,  
25 respectively, 100% and 90% pure as determined by SDS-polyacrylamide gel electrophoresis. Wheat thioredoxin *h* was purified as previously described (Johnson *et al.*, 1987).

### **Germination of Wheat Seeds**

Wheat seeds were sterilized by steeping in 50% (v/v) of Generic Bleach for 1 h at room temperature, followed by a thorough wash with distilled water.

- The sterilized seeds were placed in a plastic Petri dish on two layers of
- 5 Whatman filter paper moistened with distilled water containing 100 µg/ml of chloramphenicol. Germination was continued at room temperature in a dark chamber for up to 5 days.

### **Preparation of Wheat Proteases**

- The endosperm (10-15 g fresh weight) isolated from 5-day germinated wheat
- 10 seeds by excising the roots and shoots was extracted for 30 minutes at 4°C with 5 volumes of 200 mM sodium acetate, pH 4.6, containing 10 mM β-mercaptoethanol. The homogenate was centrifuged for 20 minutes at 48,000 g, 4°C. The pellet was discarded and the supernatant fluid was fractionated with 30-70% ammonium sulfate. This fraction, which
- 15 represented the protease preparation, was resuspended in a minimum volume of 20 mM sodium acetate, pH 4.6, containing 10 mM β-mercaptoethanol, and dialyzed against this buffer overnight at 4°C. When assayed with azocasein as substrate, the protease preparation had an optimal pH of about 4.6 and was stable for at least one week at 4°C.

### **20 Reduction and Proteolytic Susceptibility of Trypsin Inhibitors**

- Unless indicated, the reduction of the trypsin inhibitors (0.4 mg/ml) was carried out in 0.1 ml of 20 mM sodium phosphate buffer, pH 7.9 containing 10 mM EDTA at 30°C for 2 hours. The concentrations of thioredoxin, NTR, and NADPH were 0.024 mg/ml, 0.02 mg/ml, and 0.25 mM, respectively.
- 25 With DTT as reductant, EDTA and components of the NADP/thioredoxin system were omitted. Following reduction, aliquots of the inhibitor mixture were withdrawn either for determination of trypsin inhibitory activity or proteolytic susceptibility. In the subtilisin tests, the inhibitor mixture (50 µl)

was directly mixed with subtilisin and incubated at room temperature for 1 hour. With the wheat protease preparation, the pH of the inhibitor mixture (50 µl) was first adjusted to 4.7 by mixing with 35 µl of 200 mM sodium acetate, pH 4.6; 10 µl of the wheat protease preparation was then added and  
5 incubation was continued for 2 hours at 37°C. To stop digestion with subtilisin, 2 µl of 100 mM phenylmethylsulfonyl fluoride (PMSF) and 10 µl of 10% SDS were added to the digestion mixture. With the plant protease preparation, digestion was stopped by adding an equal volume of SDS sample buffer [0.125 M Tris-HCl, pH 6.8, 4% (w/v) SDS, 20% (v/v) glycerol, 10%  
10 (v/v) β-mercaptoethanol, and 0.02% (w/v) bromophenol blue]. Proteolytic products were analyzed by electrophoresis with 12% or 16% SDS polyacrylamide slab gels (Laemmli, 1970). The dried slab gels were scanned with a laser densitometer (Pharmacia-LKB UltraScan XL) and the peak area of the KTI or BBTI protein band was obtained by integration with a  
15 Pharmacia GelScan XL software program.

### Assays

Thioredoxin and NTR were assayed as previously described by Florencio *et al.* (1988). Trypsin activity was measured in 50 mM Tris-HCl, pH 7.9, by following the increase in absorbance at 253 nm with N-benzoyl-L-arginine  
20 ethyl ester as substrate (Mundy *et al.*, 1984) or by the release of azo dye into the trichloroacetic acid (TCA)-soluble fraction from azocasein substrate (see below). For trypsin inhibition assays, trypsin (5 to 10 µg) was preincubated with appropriate amounts of KTI or BBTI for 5 minutes at room temperature in 50 mM Tris-HCl, pH 7.9 and proteolytic activity was then  
25 determined. While the two substrates yielded similar data, results are presented with only one substrate.

Wheat protease activity was measured by following the release of azo dye into TCA solution from azocasein substrate at pH 4.7. Fifty µl of wheat

protease in a solution of 20 mM sodium acetate, pH 4.6, and 10 mM  $\beta$ -mercaptoethanol were added to 50  $\mu$ l of 200 mM sodium acetate, pH 4.6, and 100  $\mu$ l of 2% azocasein (in 20 mM sodium phosphate, pH 7.0). Following 1-hour incubation at 37°C, 1 ml of 10% TCA was added and the mixture was allowed to stand for 10 minutes at room temperature. After centrifugation for 5 minutes in a microfuge (8000 g), 1 ml of the supernatant solution was withdrawn and mixed with 1 ml of 1 N NaOH. The absorbance was read at 440 nm. Protein concentration was determined with Bio-Rad reagent using bovine serum albumin as a standard (Bradford, 1976).

10

## Results

### Trypsin Inhibitory Activity

The 20 kDa Kunitz and 8 kDa Bowman-Birk trypsin inhibitors of soybean contain 2 and 7 disulfide groups, respectively (Birk, 1976; Wilson, 1988). Although their physiological functions have not been established, the two types of inhibitors have been extensively investigated owing to their wide distribution in legume seeds and their potential to cause nutritional disorders, *e.g.*, hypertrophy and associated malfunctions of the pancreas. As shown in Tables I and II and described in previous Examples, KTI and BBTI are reduced specifically by the NADP/thioredoxin system from either *E. coli* or plants. The reduced forms of glutathione and glutaredoxin (a thiol protein capable of replacing thioredoxin in certain animal and bacterial systems, but not known to occur in plants (Holmgren, 1985)) were without effect.

To determine the consequence of reduction by thioredoxin, the trypsin inhibitory activity of the oxidized and reduced forms of KTI and BBTI was compared. As shown in Table XI, preincubation with the NADP/thioredoxin system (NTS) for 2 hours at 30°C resulted in a substantial loss of trypsin inhibitory activity (*i.e.*, there was an increase in trypsin activity relative to the uninhibited control). More specifically, the NADP/thioredoxin system

effected a 3- and 6-fold increase in trypsin activity for KTI and BBTI, respectively. Similar results were obtained with DTT, a nonphysiological substitute for thioredoxin, and with thioredoxin reduced by lipoic acid, a naturally occurring dithiol. Extended incubation with DTT alone (overnight at room temperature) led to complete or almost complete inactivation of both inhibitors (data not shown). Unlike DTT, lipoic acid did not reduce (inactivate) KTI and BBTI significantly in the absence of thioredoxin.

**Table XI**

Changes in the Ability of Soybean Trypsin Inhibitors to Inhibit Trypsin Following Reduction by the NADP/Thioredoxin System, DTT or Reduced Lipoic Acid

	<u>Treatment</u>	Relative Trypsin Activity*	
		<u>KTI</u>	<u>BBTI</u>
	No inhibitor	100	100
15	Inhibitor		
	Oxidized	17.0	11.5
	Reduced by NTS <sup>1</sup>	55.6	70.6
	Reduced by DTT <sup>2</sup>	68.6	88.9
	Reduced by LA/Trx <i>h</i> <sup>3</sup>	40.5	87.8

\*The specific activity of the uninhibited control trypsin was 0.018  $\Delta A_{253nm}/\mu g/min$  using N-benzoyl-L-arginine ethyl ester as substrate.

<sup>1</sup> Reduction by *E. coli* NTS (NADP/thioredoxin system) was conducted at 30°C for 2 hours.

<sup>2</sup> Reduction by DTT (1 mM) was conducted at 30°C for 1 hour.

<sup>3</sup> Reduction by lipoic acid (LA, 0.4 mM) and wheat thioredoxin *h* (Trx *h*) was conducted at 30°C for 1 hour. In the presence of lipoic acid alone (0.4 mM), trypsin activity was 20.0% for KTI and 12.5% for BBTI.

Friedman and colleagues observed that heating soybean flour in the presence of sulfur reductants (sodium sulfite, N-acetyl-L-cysteine, reduced glutathione, or L-cysteine) inactivated trypsin inhibitors, presumably as a result of the reduction or interchange of disulfide groups with other proteins in soy flour  
5 (Friedman and Gumbmann, 1986; Friedman *et al.*, 1982, 1984). Inactivation of the trypsin inhibitors by these reductants improved the digestibility and nutritive value of flours in tested rats (Friedman and Gumbman, 1986). Taken together with these earlier observations, the present findings demonstrate that disulfide bonds of both KTI and BBTI targeted by  
10 thioredoxin are important to maintenance of trypsin inhibitory activity.

### Heat Stability

Protease inhibitor proteins are typically stable to inactivation treatments such as heat. This stability is attributed, at least in part, to the cross-linking of disulfide bonds (Birk, 1976; Ryan, 1981). It is known that breaking the  
15 disulfide bonds by reduction decreases heat stability (Friedman *et al.*, 1982). The question arises as to whether reduction by thioredoxin yields similar results.

The results as shown in TABLE XII provide a positive answer to this question. When heated at 80°C for 15 minutes, the thioredoxin-reduced form  
20 of KTI completely lost its ability to inhibit trypsin, whereas its oxidized counterpart retained about half of the original activity (Table XII). Oxidized BBTI was even more stable, retaining the bulk of its trypsin inhibitory activity after heating at 100°C for 25 minutes. Nonetheless, as with KTI, the reduced form of BBTI was fully inactivated by heat (Table XII). These  
25 results are consistent with prior observations (i) that KTI and BBTI show increased sensitivity to heat on reduction; and (ii) that pure BBTI in solution is more heat-stable than pure KTI in solution. The reverse is true for flour



(i.e., KTI is more heat-stable than BBTI (Friedman *et al.*, 1982 and 1991; and DiPietro and Liener, 1989)).

**Table XII**

Heat Stability of the Kunitz and Bowman-Birk  
Trypsin Inhibitors: Oxidized and Following Reduction  
by the *E. coli* NADP/thioredoxin System

		Relative Trypsin Activity*	
<u>Treatment</u>		<u>KTI</u>	<u>BBTI</u>
	No inhibitor	100	100
10	Inhibitor, unheated		
	Oxidized	26.6	9.4
	Reduced	76.4	82.4
	Inhibitor, heated 15 min at 80°C		
	Oxidized	52.3	nd <sup>1</sup>
15	Reduced	98.7	nd
	Inhibitor, heated 25 min at 100°C		
	Oxidized	nd	17.2
	Reduced	nd	98.4
20	*The specific activity of trypsin was 0.319 $\Delta A_{440\text{nm}}$ /mg/min using azocasein as substrate. The temperatures used for inactivation were determined in initial experiments designed to show the heat stability of the trypsin inhibitors under our conditions.		

<sup>1</sup> nd: not determined.

## 25 Protease Susceptibility

To test whether the reduced forms of KTI and BBTI show decreased stability to proteases other than trypsin, both the reduced and oxidized forms of KTI and BBTI were incubated with a wheat protease preparation or with subtilisin and the proteolytic products were analyzed by SDS-PAGE. The extent of  
30 proteolysis was determined by measuring the abundance of intact protein on

SDS gels by laser densitometer. When tested with a protease preparation from 5-day germinated wheat seeds, the oxidized form of the Kunitz inhibitor was almost completely resistant to digestion whereas the thioredoxin-reduced form was susceptible to protease. As shown in Table XIII, about 80% of

5 KTI was degraded in a reaction that depended on all components of the NADP/thioredoxin system (NTS). BBTI showed the same pattern except that the oxidized protein showed greater proteolytic susceptibility relative to KTI. Similar effects were observed with both inhibitors when the plant protease preparation was replaced by subtilisin (data not shown). The nature of the

10 proteolytic reaction was not investigated, but it is noted that peptide products were not detected on SDS gels.

**Table XIII**

Effect of Thioredoxin-linked Reduction on the Susceptibility  
of Kunitz and Bowman-Birk Trypsin Inhibitors to  
Proteolysis by a Plant Protease Preparation<sup>1</sup>

5	<u>Treatment</u>		Relative Abundance <sup>2</sup>	
		<u>KTI</u>	<u>BBTI</u>	
	No protease	100	100	
	Protease			
	No reduction system	97.9	67.2	
10	<i>E. coli</i> NTS <sup>3</sup>	22.1	16.0	
	NTS minus thioredoxin	90.2	nd <sup>4</sup>	
	NTS minus NADPH	97.7	nd	
	NTS minus NTR	97.9	nd	

15 <sup>1</sup> Following reduction by *E. coli* thioredoxin system at 30°C for 2 hours, pH was adjusted to 4.7 by addition of 200 mM sodium acetate, pH 4.6. Wheat protease preparation was then added and incubated at 37°C for 2 hours, followed by SDS-PAGE analyses.

<sup>2</sup> Determined by laser densitometer.

20 <sup>3</sup> NTS: NADP/thioredoxin system.

<sup>4</sup> nd: not determined.

This Example shows that reduction by thioredoxin, or dithiothreitol (DTT), leads to inactivation of both proteins and to an increase in their heat and protease susceptibility. The results indicate that thioredoxin-linked reduction of the inhibitor proteins is relevant both to their industrial processing and to seed germination.

These results confirm the conclusion that disulfide bonds are essential for the trypsin inhibitory activity of KTI and BBTI (Birk, 1985; Friedman and Gumbmann, 1986; Friedman *et al.*, 1982,1984). These studies also show that reduction (inactivation) can take place under physiological conditions (*i.e.*, at low temperature with NADPH-reduced thioredoxin). The

ability to inactivate the trypsin inhibitors at lower temperatures provides a potential method for full inactivation of both trypsin inhibitors, thereby improving the quality of soybean products and saving energy. The need for a method for the complete inactivation of KTI is significant since 20% of its activity is consistently retained in soy flour under conditions in which BBTI is fully inactivated (Friedman *et al.*, 1991).

The present results also add new information on the protease susceptibility of KTI and BBTI. Their increase in protease susceptibility following reduction suggests that, if exposed to the protease inhibitors during seed germination, the NADP/thioredoxin system could serve as a mechanism by which the inhibitor proteins are modified (inactivated) and eventually degraded (Baumgartner and Chrispeels, 1976; Chrispeels and Baumgartner, 1978; Orf *et al.*, 1977; Wilson, 1988; Yoshikawa *et al.*, 1979). As stated previously, there is evidence that the NADP-thioredoxin system plays a similar role in mobilizing proteins during the germination of wheat seeds.

## EXAMPLE 20

### Reduction of Castor Seed 2S Albumin Protein by Thioredoxin

The results of the following study of sulfhydryl agents to reduce the 2S protein from castor seed (Sharief and Li, 1982; Youle and Huang, 1978) shows that thioredoxin actively reduces intramolecular disulfides of the 2S large subunit but not the intermolecular disulfides joining the two subunits.

## Materials and Methods

### Materials

Seeds of castor (*Ricinus communis* L. var Hale) were obtained from Bothwell Enterprises, Plainview, TX). Biochemicals were obtained from Sigma

Chemical Co. (St. Louis, MO). *E. coli* thioredoxin and NTR were isolated from cells transformed to overexpress each protein. The thioredoxin strain containing the recombinant plasmid pFPI, was kindly provided by Dr. J.-P. Jacquot (de La Mott-Guery *et al.* 1991). The strain containing the  
5 recombinant plasmid, pPMR21, was kindly provided by Drs. Marjorie Russel and Peter Model (Russel and Model, 1988). Thioredoxin and NTR were purified by the respective procedures of de La Mott-Guery *et al.* (1991) and Florencio *et al.* (1988). Reagents for SDS-polyacrylamide gel electrophoresis were purchased from Bio-Rad Laboratories (Richmond, CA).  
10 Monobromobimane (mBBr) or Thiolite was obtained from Calbiochem (San Diego, CA). Other chemicals were obtained from commercial sources and were of the highest quality available. NADP-malate dehydrogenase and fructose-1,6-bisphosphatase were purified from leaves of corn (Jacquot *et al.* 1981) and spinach (Nishizawa *et al.* 1982), respectively. Thioredoxin *h* was  
15 isolated from wheat seeds by following the procedure devised for the spinach protein (Florencio *et al.* 1988). Glutathione reductase was prepared from spinach leaves (Florencio *et al.* 1988).

### Isolation of Protein Bodies

Protein bodies were isolated by a nonaqueous method (Yatsu and Jacks,  
20 1968). Shelled dry castor seeds, 15 g, were blended with 40 ml of glycerol for 30 sec in a Waring blender. The mixture was filtered through four layers of nylon cloth. The crude extract was centrifuged at 272 x g for 5 min in a Beckman J2-21M centrifuge using a JS-20 rotor. After centrifugation, the supernatant fraction was collected and centrifuged 20 min at 41,400 x g. The  
25 pellet, containing the protein bodies, was resuspended in 10 ml glycerol and centrifuged as before (41,400 x g for 20 min) collecting the pellet. This washing step was repeated twice. The soluble ("matrix") fraction was obtained by extracting the pellet with 3 ml of 100 mM Tris-HCl buffer (pH 8.5). The remaining insoluble ("crystalloid") fraction, collected by

centrifugation as before, was extracted with 3 ml of 6M urea in 100 mM Tris-HCl buffer (pH 8.5).

### 2S Protein Purification Procedure

- The 2S protein was prepared by a modification of the method of Tully and Beevers (1976). The matrix protein fraction was applied to a DEAE-cellulose (DE-52) column equilibrated with 5 mM Tris-HCl buffer, pH 8.5 (Buffer A) and eluted with a 0 to 300 mM NaCl gradient in buffer A. Fractions containing the 2S protein were pooled and concentrated by freeze drying. The concentrated fraction was applied to a Pharmacia FPLC Superose-12 (HR 10/30) column equilibrated with buffer A containing 150 mM NaCl. The fraction containing 2S protein from the Superose-12 column was applied to an FPLC Mono Q HR 5/5 column equilibrated with buffer A. The column was eluted sequentially with 3 ml of buffer A, 20 ml of a linear gradient of 0 to 300 mM NaCl in buffer A and finally with buffer A containing 1 M NaCl. The 2S protein purified by this method was free of contaminants in SDS polyacrylamide gels stained with Coomassie blue (Kobrehel *et al.*, 1991).

### Analytical Methods

- Reduction of proteins was monitored by the monobromobimane (mBBr)/SDS polyacrylamide gel electrophoresis procedure of Crawford *et al.* (1989). Labeled proteins were quantified as described previously in the "Reduction of Cereal Proteins, Materials and Methods" section. Protein was determined by the method of Bradford (1976).

### Enzyme Assays/Reduction Experiments

- The Wada *et al.*, 1981 protocol was used for assaying NADP-malate dehydrogenase and fructose 1,6 biphosphatase in the presence of thioredoxin and 2S protein. Assays were conducted under conditions in which the

amount of added thioredoxin was sufficient to reduce the castor 2S protein but insufficient to activate the target enzyme appreciably. All assays were at 25°C. Unless otherwise indicated, the thioredoxin and NTR used were from *E. coli*. The 2S protein was monitored during purification by mBBR/SDS-  
5 polyacrylamide gel electrophoresis following its reduction by dithiothreitol and *E. coli* thioredoxin (Crawford *et al.*, 1989; Kobrehel *et al.*, 1991).

The reduction of the matrix and crystalloid proteins from castor seed were determined by the mBBR/SDS-polyacrylamide gel electrophoresis procedure. The lanes for the gels (not shown) were as follows, 1 and 7, Control: no  
10 addition; 2 and 8, GSH/GR/NADPH: reduced glutathione, glutathione reductase (from spinach leaves) and NADPH; 3 and 9, NGS: NADPH, reduced glutathione, glutathione reductase (from spinach leaves) and glutaredoxin from *E. coli*; 4 and 10, NTS: NADPH, NTR, and thioredoxin (both proteins from *E. coli*); 5 and 11, NADPH; 6 and 12, NADPH and *E.*  
15 *coli* NTR. Reactions were carried out in 100 mM Tris-HCl buffer, pH 7.8. As indicated, 0.7 µg NTR and 1 µg thioredoxin were added to 70 µl of this buffer containing 1 mM NADPH and target protein: 8 µg matrix protein for treatments 1-6 and 10 µg crystalloid protein for treatments 7-12. Assays with  
20 glutathione were performed similarly, but at a final concentration of 2 mM, 1.4 µg glutathione reductase, 1 µg glutaredoxin, and 1.5 mM NADPH. Reaction time was 20 min.

The mBBR/SDS-Page technique was also used to determine the specificity of thioredoxin for reducing the disulfide bonds of castor seed 2S protein. The lanes for the gel (not shown) were as follows, (1) Control (no addition);  
25 (2) Control + NTS (same conditions as with the matrix and crystalloid proteins); (3) Control (heated 3 min at 100°C); (4) Control + 2 mM DTT (heated 3 min at 100°C). The samples containing 5 µg 2S protein and the indicated additions were incubated for 20 min in 30 mM Tris-HCl (pH 7.8).

mBBR, 80 nmol, was then added and the reaction continued for another 15 min prior to analysis by the mBBR/SDS polyacrylamide gel electrophoresis procedure.

## Results

- 5 The castor storage proteins, which are retained within a protein body during seed maturation, can be separated into two fractions on the basis of their solubility. The more soluble proteins are housed in the protein body outer section ("matrix") and the less soluble in the inner ("crystalloid"). In the current study, the matrix and crystalloid components were isolated to
- 10 determine their ability to undergo reduction by cellular thiols, viz., glutathione, glutaredoxin and thioredoxin. Glutaredoxin, a 12 kDa protein with a catalytically active thiol group, can replace thioredoxin in certain enzymic reactions of bacteria and animals (Holmgren *et al.* 1985) but is not known to occur in plants.
- 15 The results showed that, while a number of storage proteins of castor seed were reduced by the thiols tested, only a low molecular weight protein, corresponding to the large subunit of the 2S protein of the matrix, showed strict specificity for thioredoxin. Certain higher molecular weight proteins of the crystalloid fraction underwent reduction, but in those cases there was little
- 20 difference between glutaredoxin and thioredoxin. The castor seed 2S large subunit thus appeared to resemble cystine-containing proteins previously discussed in undergoing thioredoxin-specific reduction. These experiments were designed to confirm this specificity and to elucidate certain properties of the reduced protein. As expected, owing to lack of disulfide groups, the 2S
- 25 small subunit showed essentially no reaction with mBBR with any of the reductants tested.



When its fluorescent band was monitored by laser densitometry, the reduction of the castor seed 2S large subunit was found to depend on all components of the NADP/thioredoxin system (NADPH, NTR and thioredoxin) (Table XIV). As for other thioredoxin-linked proteins (including chloroplast enzymes), the thioredoxin active in reduction of the 2S large subunit could be reduced either chemically with dithiothreitol (DTT) or enzymatically with NADPH and NTR. The extent of reduction by the NADP thioredoxin system, DTT alone, and DTT + thioredoxin was 84%, 67% and 90%, respectively, after 20 min at 25°C. Similar, though generally extensive reduction was observed with the disulfide proteins discussed above (Johnson et al. 1987). As with the other seed proteins, native wheat thioredoxin *h* and *E. coli* thioredoxins could be used interchangeably in the reduction of the 2S protein by DTT (data not shown).

**Table XIV**

Extent of reduction of the castor seed 2S protein by different sulfhydryl reductants. Reaction conditions as with the matrix and crytalloid protein determination. A reduction of 100% corresponds to that obtained when the 2S protein was heated for 3 min in the presence of 2% SDS and 2.5%  $\beta$ -mercaptoethanol. NTS: NADPH, NTR, and thioredoxin (both proteins from *E. coli*); GSH/GR/NADPH: reduced glutathione, glutathione reductase (from spinach leaves) and NADPH; NGS: NADPH, reduced glutathione, glutathione reductase (from spinach leaves) and glutaredoxin (from *E. coli*).

	<u>Treatment</u>	<u>Relative Reduction (%)</u>
25	Control	0
	NADP/thioredoxin system, complete	84
	NADP minus thioredoxin	0
	NADP minus NADPH	0
	NADP minus NTR	0

Reduced glutathione	0
NADP/glutaredoxin system, complete	0
DTT	67
DTT + thioredoxin	90

- 5 The capability of thioredoxin to reduce the castor seed 2S protein was also evident in enzyme activation assays. Here, the protein targeted by thioredoxin (in this case 2S) is used to activate a thioredoxin-linked enzyme of chloroplasts, NADP-malate dehydrogenase or fructose 1,6-bisphosphatase. As with most of the proteins examined so far, the 2S protein more effectively
- 10 activated NADP-malate dehydrogenase and showed little activity with the fructose bisphosphatase (2.6 vs. 0.0 nmoles/min/mg protein).

The castor seed 2S protein contains inter-as well as intramolecular disulfides. The 2S protein thus provides an opportunity to determine the specificity of thioredoxin for these two types of bonds. To this end, the castor seed 2S

15 protein was reduced (i) enzymically with the NADP/thioredoxin system at room temperature, and (ii) chemically with DTT at 100°C. Following reaction with mBBBr the reduced proteins were analyzed by SDS-polyacrylamide gel electrophoresis carried out without additional sulfhydryl agent. The results indicate that while thioredoxin actively reduced

20 intramolecular disulfides, it was much less effective with intermolecular disulfides.

The present results extend the role of thioredoxin to the reduction of the 2S protein of castor seed, an oil producing plant. Thioredoxin specifically reduced the intramolecular disulfides of the large subunit of the 2S protein

25 and showed little activity for the intermolecular disulfides joining the large and small subunits. Based on the results with the trypsin inhibitors of soybean, it is clear that reduction of intramolecular disulfides by thioredoxin

markedly increases the susceptibility of disulfide proteins to proteolysis (Jiao *et al.* 1992a). It, however, remains to be seen whether reduction of the 2S protein takes place prior to its proteolytic degradation (Youle and Huang, 1978) as appears to be the case for the major storage proteins of wheat. A  
5 related question raised by this work is whether the 2S protein of castor, as well as other oil producing plants such as brazil nut (Altenbach *et al.*, 1987; Ampe *et al.*, 1986), has a function in addition to that of a storage protein.

## EXAMPLE 21

### 10 Thioredoxin-Dependent Deinhhibition of Pullulanase of Cereals by Inactivation of a Specific Inhibitor Protein

#### Assay of Pullulanase

##### 1. Standard curve of maltotriose:

A series of concentrations of maltotriose (0 to 2 mg) in 0.1 to 0.2 ml water or buffer were made in microfuge tubes. To this was added 0.2 ml of  
15 dinitrosalicylic acid (DA) reagent (mix 1 g of DA, 30 g of sodium potassium tartrate, and 20 ml of 2N NaOH with water to final volume of 100 ml). The reagents were dissolved in a warm water bath. The mixture was heated at 100°C for 5 min and cooled down in a water bath (room temperature). Each sample was transferred to a glass tube that contained 2 ml of water. Read  
20  $A_{493}$  vs water.  $\Delta A_{493}$  [ $A_{493}$  of sample containing maltotriose was subtracted from  $A_{493}$  of the blank (no maltotriose)] was plotted against maltotriose concentrations.

##### 2. Pullulanase activity assay:

Pullulanase activity is measured as the release of reducing sugar from the  
25 substrate pullulan. Typically 10-100  $\mu$ l of pullulanase sample (in 20 mM Tris-HCl, pH 7.5, or in 5-20 acetate-NA, pH 4.6) was mixed with 25-100  $\mu$ l of 200 mM Acetate-NA, pH 5.5 (this buffer serves to bring final pH of the

assay to 5.5) and 10-20  $\mu$ l of 2 % (w/v) pullulan. The mixture was incubated at 37°C for 30 to 120 min, depending on the activity of pullulanase. The reaction was stopped by adding 200  $\mu$ l of DA reagent. Reducing sugar was then determined as above.

5 Note:

1. When a crude extract of pullulanase obtained by the dialysis of crude extracts or pullulanase obtained from a dialyzed 30-60% ammonium sulfate fraction is used as a pullulanase source, it must be thoroughly dialysed before assay because there are reduced sugars in the crude extract. In other words the background of crude pullulanase samples from dialysed crude  
10 extracts or a dialysed 30-60% ammonium sulfate fraction is very high. In this case, the blank is made as follows: 200  $\mu$ l of DA reagent are added first, followed by the addition of enzyme sample, pullulan and buffer.

2. When final concentrations of DTT (or  $\beta$ -mercaptoethanol (MET) or GSH) are higher than 2 mM in the assay mixture, the OD<sub>493</sub> values will be  
15 greater than those of the minus-DTT (MET, GSH) samples. DTT (MET, GSH) should be added to the blank, samples without DTT during assay at the end of the reaction. Care should be taken to make sure the final concentration of DTT in the assay mixture is below 2 mM.

20 **Purification of Pullulanase Inhibitor**  
**Extraction and Ammonium Sulfate Fractionation**

200 g of barley malt was ground to fine powder with an electric coffee grinder and extracted with 600 ml of 5 % (w/v) NaCl for 3 h at 30°C. Following centrifugation at 30,000 g and at 4°C for 25 min, the supernatant  
25 was fractionated by precipitation with solid ammonium sulfate. Proteins precipitated between 30% and 60% saturated ammonium sulfate were dissolved in a minimum volume of 20 mM Tris HCl, pH 7.5, and dialyzed against this buffer at 4°C overnight.

### **DE52 Chromatography**

The dialyzed sample was centrifuged to remove insoluble materials and the supernatant adjusted to pH 4.6 with 2N formic acid. After pelleting the acid-denatured protein, the supernatant was readjusted to pH 7.5 with  $\text{NH}_4\text{OH}$  and  
5 loaded onto a DE52 column (2.5 x 26 cm) equilibrated with 20 mM Tris-HCl, pH 7.5. Following wash with 80 ml of the above buffer, the column was eluted with a linear 0-500 mM Tris-HCl, pH 7.5. Fractions of 6.7 ml were collected. Pullulanase was eluted at about 325 mM NaCl and its inhibitor came off at about 100 mM NaCl. Pullulanase was further purified  
10 through CM32 (20 mM sodium acetate, pH 4.6) and Sephacryl-200 HR (30 mM Tris-HCl, pH 7.5, containing 200 mM NaCl and 1 mM EDTA) chromatography. Pullulanase inhibitor protein was purified as described below.

### **CM32 Chromatography**

15 The pullulanase inhibitor sample (about 90 ml) from the DE52 step was placed in a 150-ml flask and incubated at 70°C water-bath for 20 min. Following centrifugation, the clarified sample was then adjusted to pH 4.6 with 2N formic acid and dialyzed against 20 mM sodium acetate, pH 4.6. The precipitate formed during dialysis was removed by centrifugation and the  
20 supernatant was chromatographed on a CM32 column (2.5 X 6 cm) equilibrated with 20 mM sodium acetate, pH 4.6. Proteins were eluted with a linear 0-0.4 M NaCl in 200 ml of 20 mM sodium acetate, pH 4.6. Fractions (5.0 ml/fraction) containing pullulanase inhibitory activity were pooled, dialyzed, and rechromatographed on a CM32 column (2.5 x 6 cm)  
25 with a linear 0.2-1 M NaCl gradient in 200 ml of 20 mM sodium acetate, pH 4.0.

### **Sephadex G-75 Filtration**

Pullulanase inhibitor fractions from the second CM32 chromatography step were concentrated in a dialysis bag against solid sucrose and then separated by a Sephadex G-75 column (2.5 x 85 cm) equilibrated with 30 mM Tris-HCl, pH 7.5, containing 200 mM Na Cl and 1 mM EDTA. Fractions (3.6 ml/fraction) showing pullulanase inhibitory activity were pooled, concentrated by dialysis against solid sucrose, and then dialysed against 10 mM Tris-HCl, pH 7.5.

### **Identification and Purification of Pullulanase Inhibitor**

During germination, starch is converted to glucose by  $\alpha$ -,  $\beta$ -amylases, and pullulanase (also called debranching enzyme, R-enzyme). While extensive studies have been conducted for the regulation of amylases, little is known about the regulation of pullulanase in seeds. Yamada (Yamada, J. (1981) *Carbohydrate Research* 90:153-157) reported that incubation of cereal flours with reductants (*e.g.*, DTT) or proteases (*e.g.*, trypsin) led to an activation of pullulanase activity, suggesting that reduction or proteolysis might be a mechanism by which pullulanase is activated during germination. Like in rice flour, pullulanase extracts from germinated wheat seeds or from barley malt showed lower activity, and were activated 3 to 5-fold by preincubation with DTT for 20 to 30 min. However, following purification of the crude extract (a dialysate of 30-60% ammonium sulfate fraction) by anion or cation exchange chromatography, the total pullulanase activity increased 2 to 3-fold over that of the sample applied to the column when assayed without preincubation with DTT, and DTT has no or little effect on pullulanase. One possibility was that pullulanase might be activated by proteolysis during the process of purification, since germinated wheat seeds or barley malt show high protease activity. If this was the case, addition of protease inhibitor cocktail would prevent pullulanase activation during purification. In contrast to this point, many experiments with protease inhibitors failed to prove this.

Another possibility was that there is an inhibitor that is precipitated by ammonium sulfate and inhibits pullulanase. The role of DTT is to reduce and thus inactivate this protein inhibitor, leading to activation of pullulanase. Along this line, the 30-60% ammonium sulfate fraction from barley malt was applied to a DE52 column (2.5 x 26 cm) equilibrated with 20 mM Tris-CH<sub>3</sub>COOH, pH 7.5. Following elution with a linear salt gradient, "deinhibited" ("activated") pullulanase was identified as a protein peak coming off at about 325 mM NaCl (from fraction numbers 44 to 60). Assay of pullulanase activity in the preincubation mixture consisting of 50 µl of the peak pullulanase activity fraction (fraction number 45) with 50 µl of other protein fractions indicated that a protein peak that showed pullulanase inhibitory activity was eluted from the DE52 column by about 100 mM NaCl between fraction numbers 8 to 25.

The pullulanase inhibitor sample was further purified by two consecutive cation exchange chromatography steps with CM32 at pH 4.6 and 4.0, and filtration with Sephadex G-75.

#### **Properties of Pullulanase Inhibitor**


Preliminary experiments showed that pullulanase inhibitor protein is resistant to treatment of 70°C for 10 min and pH 4.0. Based on the profile of Sephadex G-75 gel filtration and SDS-PAGE, pullulanase inhibitor has a molecular weight between 8 to 15 kDa  $\pm$  2 kDa. The study further showed that the protein contains thioredoxin-reducible (S-S) bonds.

These studies, as shown in Table XV, found that the ubiquitous dithiol protein, thioredoxin, serves as a specific reductant for a previously unknown disulfide-containing protein that inhibits pullulanase of barley and wheat endosperm.


**Table XV**  
**Activity Change in Pullulanase Inhibitor Protein**  
**Following Reduction by NADP/Thioredoxin System**

5	<u>Treatment</u>	<u>Relative Pullulanase Activity</u>
	No inhibitor	100
	Inhibitor	
10	Oxidized	30.1
	Reduced by DTT	46.1
	Reduced by <i>E. coli</i> Trx/DTT	95.1
	Reduced by <i>E. coli</i> NTS	40.4
	Reduced by GSH/NADPH/GR	33.6

Reduction of the inhibitor protein eliminated its ability to inhibit pullulanase,  
 15 thereby rendering the pullulanase enzyme active. These studies as shown in  
 Table XV illustrate that it is possible to render the pullulanase enzyme active  
 with a physiological system consisting of NADPH, NADP-thioredoxin  
 reductase (NTR) and thioredoxin (the NADP/thioredoxin system) as well as  
 with thioredoxin (Trx) and dithiothreitol. These findings also elucidate how  
 20 reductive activation of pullulanase takes place (*i.e.*, that a specific (previously  
 unknown) inhibitor is reduced and thereby inactivated, so that the enzyme  
 becomes active). The thioredoxin active in this reaction can be obtained from  
 several sources such as *E. coli* or seed endosperm (thioredoxin *h*). The role  
 of thioredoxin in reductively inactivating the inhibitor protein (I) and  
 25 deinhibiting the pullulanase enzyme (E) is given in Equations 1 and 2.







In summary, the crude endosperm extracts were fractionated by column chromatography procedures. These steps served to separate the protein inhibitor from the pullulanase enzyme. The inhibitor protein was then highly purified by several steps. By use of the mBBR/SDS-PAGE procedure, it was determined that disulfide group(s) of the new protein are specifically reduced by thioredoxin and that the reduced protein loses its ability to inhibit pullulanase. Like certain other disulfide proteins of seeds (*e.g.*, the Kunitz and Bowman-Birk trypsin inhibitors of soybean), the pullulanase inhibitor protein showed the capability to activate chloroplast NADP-malate dehydrogenase. In these experiments, dithiothreitol was used to reduce thioredoxin, which in turn reduced inhibitor and thereby activated the dehydrogenase enzyme.

15

## EXAMPLE 22

### Engineering of Yeast Cells to Overexpress Thioredoxin and NADP-Thioredoxin Reductase

The two *Saccharomyces cerevisiae* thioredoxin genes (Muller, E.G.D. (1991), *J. Biol. Chem.* 266:9194-9202), TRX1 and TRX2, are cloned in high copy number episomal vectors, an example of which is YEp24, under the control of strong constitutive promoter elements, examples of which are the glycolytic promoters for the glyceraldehyde-3-P dehydrogenase, enolase, or phosphoglycerate kinase genes. Recombinant constructs are assessed for the overexpression of thioredoxin by quantitative Western blotting methods using an antithioredoxin rabbit antiserum (Muller, E.G.D. *et al.* (1989), *J. Biol. Chem.* 264:4008-4014), to select the optimal combination of thioredoxin genes and promoter elements. The cells with the optimal thioredoxin

overexpression system are used as a source of thioredoxin for dough improvement.

The NADP-thioredoxin reductase gene is cloned by preparing an oligonucleotide probe deduced from its amino terminal sequence. The  
5 enzyme is prepared from yeast cells by following a modification of the procedure devised for spinach leaves (Florencio, F.J. *et al.* (1988), *Arch. Biochem. Biophys.* 266:496-507). The amino terminus of the pure reductase enzyme is determined by microsequencing by automated Exman degradation with an Applied Biosystems gas-phase protein sequencer. On the basis of this  
10 sequence, and relying on codon usage in yeast, a 20-base 24-fold degenerate DNA probe is prepared. The probe is hybridized to isolated yeast DNA cleaved with *EcoRI* and *PstI* by Southern blot analysis. The most actively region is extracted from the agarose gels and introduced into a pUC19 plasmid vector (Szekeres, M. *et al.* (1991), *J. Bacteriol.* 173:1821-1823).  
15 Following transformation, plasmid-containing *E. coli* colonies are screened by colony hybridization using the labeled oligonucleotide probe (Vogeli, G. *et al.* (1987), *Methods Enzymol.* 152:407-415). The clone is identified as carrying the gene for NADP-thioredoxin reductase by sequencing the DNA as given in Szekeres *et al.* above. Once identified, the NADP-thioredoxin  
20 reductase gene is overexpressed in yeast as described above for the TRX1 and TRX2 yeast thioredoxin genes. The yeast cells which overexpress NADP-thioredoxin reductase are used as a source of reductase to improve dough quality.

### EXAMPLE 23

#### Improvement in Dough Quality Using Genetically Engineered Yeast Cells

*Saccharomyces cerevisiae* cells engineered to overexpress the two yeast  
5 thioredoxins and the yeast NADP-thioredoxin reductase as set forth in  
Example 23 are lysed by an established procedure such as sonication and then  
freeze dried. The dried cells from the cultures overexpressing thioredoxin  
and NADP-thioredoxin reductase are combined and then used to supplement  
10 flour to improve its dough quality. Two-tenths gram of the combined lysed  
dried cells are added together with about 300 to about 500 nanomoles  
NADPH to 1 M Tris-HCl buffer, pH 7.9, to give 5.25 ml of 30 mM Tris-  
HCl. The reaction is carried out in a microfarinograph at 30°C as described  
in Example 14. An improvement in dough quality is observed which is  
similar to the improvement shown in Example 14.

### EXAMPLE 24

#### Improvement of Gluten

The positive effects of the NADP/thioredoxin system on dough quality  
presents the option of applying this system to flour in the preparation of  
gluten. The purpose is to alter the yield and the properties of gluten, thereby  
20 enhancing its technological value: (1) by obtaining stronger glutes (increased  
elasticity, improved extensibility); (2) by increasing gluten yield by capturing  
soluble proteins, reduced by the NADP-thioredoxin system, in the protein  
network, thereby preventing them from being washed out during the  
production of gluten. In this procedure (using 10 g flour), 0.2 µg *E. coli*  
25 thioredoxin, 0.1 µg *E. coli* NADP-thioredoxin reductase and 300 to 500  
nanomoles NADPH are added together with 1 M Tris-HCl, pH 7.9, buffer to  
give 5.25 ml of 30 mM Tris-HCl. The gluten is made at room temperature  
according to the common lixiviation method. The yield of the gluten is  
determined by weight and the strength of the gluten is determined by the

classical manual stretch method. The gluten product which are obtained by this treatment with the NADP/thioredoxin system is used as an additive in flour or other grain.

### EXAMPLE 25

#### 5           **Method of Producing Dough from a Non-wheat or Rye Flour**

For this test (using 10 gm of milled flour from corn, rice or sorghum), 0.2  $\mu$ g *E. coli* thioredoxin, 0.1  $\mu$ g *E. coli* NADP-thioredoxin reductase and 500 nanomoles NADPH are added together with 1 M Tris-HCl, pH 7.9, buffer to give 5.25 ml of 30 mM Tris-HCl. The reaction is carried out by mixing the  
10 10 gm of milled flour with the enzyme system in a micro-farinograph at 30°C. The farinograph measurements show wheat-like dough characteristics by the added NADP-thioredoxin system. In the controls without the enzyme system, no microfarinograph reading is possible because the mixture fails to form a dough. The dough which is formed is persistent and its consistency is  
15 maintained throughout the run. The end product is similar to the network formed in dough derived from wheat.

### **Reduction of Animal Toxins**

The invention provides a method for chemically reducing toxicity causing proteins contained in bee, scorpion and snake venome and thereby altering  
20 the biological activity of the venoms well as reducing the toxicity of animal toxins specifically snake neurotoxins by means of thiol redox (SH) agents namely a reduced thioredoxin, reduced lipoic acid in the presence of a thioredoxin or DTT. The reduction of the thioredoxin occurs preferably via the NADP-thioredoxin system (NTS). As stated previously, the NTS  
25 comprises NADPH, NADP-thioredoxin reductase (NTR) and a thioredoxin.

The term "thiol Redox agent" has been used sometimes in the literature to denote both an agent in the nonreduced state and also in the reduced or sulfhydryl (SH) state. As defined herein the term "thiol redox (SH) agent" means a reduced thiol redox protein or synthetically prepared agent such as DTT.

The reduction of the neurotoxin may take place in a medium that is liquid such as blood, lymph or a buffer, etc. or in a medium that is solid such as cells or other living tissue. As used herein the term "liquid" by itself does not refer to a biological fluid present in an individual.

10 Presumably the proficiency of the thiol redox (SH) agents to inactivate the venom *in vitro* and to detoxify the venom in individuals depends upon the ability of the agents of the invention to reduce the intramolecular disulfide bonds in these toxicity causing venom components.

15 All snake neurotoxins, both presynaptic and postsynaptic can be reduced and at least partially inactivated *in vitro* by the thiol redox (SH) agents of the invention. Snake neurotoxins inactivated *in vitro* according to the invention are useful as antigens in the preparation of antivenoms. The neurotoxins are inactivated preferably by incubation with a thiol redox (SH) agent in an appropriate buffer. The preferred buffer is Tris-HCl buffer but other buffers  
20 such as phosphate buffer may be used. The preferred thiol redox (SH) agent is a reduced thioredoxin.

Effective amounts for inactivating snake neurotoxins range from about 0.1  $\mu\text{g}$  to 5.0  $\mu\text{g}$ , preferably about 0.5  $\mu\text{g}$  to 1.0  $\mu\text{g}$ , of a reduced thioredoxin; from about 1 nanomole to 20 nanomoles, preferably from 5 nanomoles to 15  
25 nanomoles, of reduced lipoic acid in the presence of about 1.0  $\mu\text{g}$  of a thioredoxin and from about 10 nanomoles to 200 nanomoles, preferably 50

nanomoles to 100 nanomoles, of reduced DTT (preferably in the presence of about 1.0  $\mu\text{g}$  of a thioredoxin) for every 10  $\mu\text{g}$  of snake neurotoxin in a volume of 100  $\mu\text{l}$ .

- The effective amounts for inactivating a snake neurotoxin using the
- 5 components in the NADP-thioredoxin system range from about 0.1  $\mu\text{g}$  to 5.0  $\mu\text{g}$ , preferably about 0.5  $\mu\text{g}$  to 1.0  $\mu\text{g}$ , of thioredoxin; from about 0.1  $\mu\text{g}$  to 2.0  $\mu\text{g}$ , preferably from 0.2  $\mu\text{g}$  to 1.0  $\mu\text{g}$ , of NTR and from about 0.05 micromoles to 0.5 micromoles, preferably about 0.1 micromoles to 0.25 micromoles, of NADPH for every 10  $\mu\text{g}$  of snake neurotoxin in a
- 10 volume of 100  $\mu\text{l}$ .

Upon inactivation the buffer containing the inactivated neurotoxin and thiol redox (SH) agent, etc. may be injected into an animal such as a horse to produce an antivenom or prior to injection it may be further treated with heat or formaldehyde.

- 15 The thiol redox (SH) agents of the invention may also be used to treat individuals who are suffering the effects of neurotoxicity caused by a venomous snake bite. The preferred method of administering the reduced thiol redox (SH) agent to the individual is by multiple subcutaneous injections around the snake bite.
- 20 Of course the correct amount of a thiol redox (SH) agent used to detoxify a neurotoxin in an individual will depend upon the amount of toxin the individual actually received from the bite. However, effective amounts for detoxifying or reducing the toxicity of snake neurotoxins in mice usually range from about 0.01  $\mu\text{g}$  to 0.3  $\mu\text{g}$ , preferably about 0.02  $\mu\text{g}$  to 0.05  $\mu\text{g}$ ,
- 25 of a reduced thioredoxin; from about 0.1 nanomole to 3.0 nanomoles, preferably from 0.2 nanomole to 1.0 nanomole, of reduced lipoic acid in the

presence of about 0.05  $\mu\text{g}$  of a thioredoxin; from about 1.0 nanomole to 30 nanomoles, preferably from 2.0 nanomoles to 5.0 nanomoles, of DTT, preferably in the presence of 0.05  $\mu\text{g}$  of a thioredoxin, for every gm of mouse body weight.

- 5 The effective amounts for detoxifying a snake neurotoxin in a mouse using the components of the NADP-thioredoxin system range from about 0.01  $\mu\text{g}$  to 0.3  $\mu\text{g}$ , preferably about 0.02  $\mu\text{g}$  to 0.05  $\mu\text{g}$  of a thioredoxin; from about 0.005  $\mu\text{g}$  to 0.12  $\mu\text{g}$ , preferably from 0.01  $\mu\text{g}$  to 0.025  $\mu\text{g}$ , of NTR and from about 5 nanomoles to 30 nanomoles, preferably 10 nanomoles to
- 10 15 nanomoles, NADPH for every gm of mouse body weight.

The preferred method of administering the NTS to an individual is also by multiple subcutaneous injections. The preferred thiol redox agent for human use is human thioredoxin administered via the NADP-thioredoxin system or with lipoic acid or DTT.

15

A partial list of the venomous snakes which produce the neurotoxins which can be inactivated or detoxified by the methods of this invention appears on pages 529-555 of Chippaur, J.-P. *et al.* (1991) *Reptile Venoms and Toxins*, A.T. Tu, ed., Marcel Dekker, Inc., which is herein incorporated by

20 reference.

Other features and advantages of the invention with respect to inactivating and detoxifying venome can be ascertained from the following examples.

## EXAMPLE 26

### Reduction Studies of Bee, Scorpion and Snake Venoms and Labeling with mBBr

- Reactions were carried out with 50  $\mu$ g venom (final volume of 100  $\mu$ l) in 30 mM Tris-CHl buffer pH 7.9 containing the following protease inhibitors: phenylmethylsulfonyl fluoride (PMSF), leupeptin and pepstatin (final concentrations used in the assay respectively: 100  $\mu$ M, 1  $\mu$ M and 1  $\mu$ M). With NADPH as a reductant, the mixture also contained 4  $\mu$ g thioredoxin, 3.5  $\mu$ g NTR (both from *E. coli*) and 12.5 mM NADPH. When thioredoxin (4  $\mu$ g, *E. coli* or human) was reduced by DTT, NADPH and NTR were omitted and DTT was added to 0.5 mM. Assays with GSH were performed similarly but at a final concentration of 5 mM and in the presence of 1.5  $\mu$ g glutathione reductase and 12.5 mM NADPH. The mixture was incubated for 20 min at room temperature, mBBr was then added to 1.5 mM and the reaction was continued for 15 min at room temperature. The reaction was stopped and excess mBBr derivitized by adding 10  $\mu$ l of 100 mM  $\beta$ -mercaptoethanol, 5  $\mu$ l of 20% SDS and 10  $\mu$ l of 50% glycerol. Samples were then analyzed by SDS-polyacrylamide gel electrophoresis as previously described.
- The same experiment with the NADP-thioredoxin system was performed without adding protease inhibitors.

- After 20 min incubation at room temperature with different reductants and in the presence of protease inhibitors, the samples were derivitized with mBBr and separated by electrophoresis and fluorescence was determined. It was shown that in all cases thioredoxin (*E. coli* or human) specifically reduced components of the venoms. The gel also showed that thioredoxin reduces venom components in a similar way when the reaction was performed in the absence of protease inhibitors.



The reduction of bee, scorpion and snake venoms by the NADP-Thioredoxin system with and without protease inhibitors was also shown using the SDS-Polyacrylamide gel mBBr procedure. After 20 min incubation at room temperature with NTS in the presence or absence of any protease inhibitors, 5 the samples were derivatized with mBBr, separated by electrophoresis, and fluorescence was determined as previously described.

### Materials

Venoms: Bee venom from *Apis mellifera*, scorpion venom from *Androctonus australis*, and snake venom from *Bungarus multicinctus* were 10 purchased from Sigma chemical Co. (St. Louis, MO).

Protease Inhibitors: Phenylmethylsulfonyl fluoride (PMSF), Leupeptin and Pepstatin were purchased from Sigma Chemical Co. (St. Louis, MO).

### Venom Detoxification

Detoxification of bee, scorpion and snake venoms is determined by 15 subcutaneous injection into mice. Assays are done in triplicate. Prior to injection, the venom is diluted in phosphate-saline buffer (0.15 M NaCl in 10 mM  $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$  pH 7.2) at concentrations ranging up to twice the  $\text{LD}_{50}$  (per g mouse): bee venom from *Apis mellifera*, 2.8  $\mu\text{g}$ ; scorpion venom from *Androctonus australis*, 0.32  $\mu\text{g}$ ; and snake venom from *Bungarus* 20 *multicinctus*, 0.16  $\mu\text{g}$ . At 5, 10, 30, 60 minutes and 4, 12 and 24 hr after injection, separate groups of challenged mice are injected (1) intravenously and (2) subcutaneously (multiple local injections around the initial injection site). The thioredoxin is reduced with : (1) *E. coli* NADP-thioredoxin system, using 0.08  $\mu\text{g}$  thioredoxin, 0.07  $\mu\text{g}$  NTR and 25 nmoles NADPH; 25 (2) Thioredoxin reduced by DTT or reduced lipoic acid (0.08  $\mu\text{g}$  *E. coli* or human thioredoxin added to 1 nmole dithiothreitol or 1 nmole of reduced

lipoic acid). Concentrations are per  $\mu\text{g}$  venom injected into the animal; all solutions are prepared in phosphate-saline buffer.

The effect of thioredoxin on detoxification is determined by (1) comparing the  $\text{LD}_{50}$  with the control group without thioredoxin and (2) following the  
5 extent of the local reaction, as evidenced by necrosis, swelling and general discomfort to the animal.

### **Reduction Studies for Reducing Snake Neurotoxins - Materials and Methods**

#### **Toxins**

10 Porcine pancreas phospholipase  $\text{A}_2$ , erabutoxin b and  $\beta$ -bungarotoxin were purchased from Sigma Chemical Co. (St. Louis, Mo). As the phospholipase  $\text{A}_2$  was provided in 3.2 M  $(\text{NH}_4)_2\text{SO}_4$  solution pH 5.5, the protein was dialysed in 30 mM Tris-HCl buffer, pH 7.9, using a centricon 3 KDa cutoff membrane.  $\alpha$ -Bungarotoxin and  $\alpha$ -bungarotoxin $^{125}\text{I}$  were a kind gift from  
15 Dr. Shalla Verrall.

#### **Reagents and fine chemicals**

DL- $\alpha$ -Lipoic acid, L- $\alpha$ -phosphatidylcholine from soybean, NADPH and  $\beta$ -mercaptoethanol were purchased from Sigma Chemical Co. (St Louis, MO) and monobromobimane (mBBBr, trade name thiolite) from Calbiochem (San  
20 Diego, CA). Reagents for sodium dodecylsulfate (SDS)-polyacrylamide gel electrophoresis were purchased from Bio-Rad Laboratories (Richmond, CA).

#### **Proteins and Enzymes**

Thioredoxin and NTR were purified from *E. coli* as is described by Jiao *et al.*, (1992) *Ag. Food Chem. (in press)*. Thioredoxin *h* was purified from  
25 wheat germ (Florencio, F.J. *et al.* (1988) *Arch Biochem. Biophys.* 266:496-507) and thioredoxins *f* and *m* from spinach leaves (Florencio, F.J. *et al.*, *supra.*). Human thioredoxin was a kind gift of Dr. Emanuelle Wollman.

NADP-malate dehydrogenase was purified from corn leaves (Jacquot, J.-P. *et al.* (1981) *Plant Physiol.* 68:300-304) and glutathione reductase from spinach leaves (Florencio, F.J. *et al.*, *supra.*). *E. coli* glutaredoxin was a kind gift of Professor A. Holmgren.

## 5 SDS-Polyacrylamide Gel Electrophoresis

SDS-polyacrylamide gel electrophoresis was performed in 10-20% gradient gels of 1.5 mm thickness that were developed for 3 hr at a constant current of 40 mA. Following electrophoresis, gels were soaked for 2 hr in 12% (w/v) trichloroacetic acid and then transferred to a solution containing 40% methanol and 10% acetic acid for 12 hr to remove excess mBBr. The fluorescence of protein-bound mBBr was determined by placing gels on a light box fitted with an ultraviolet light source (365 nm). Gels were photographed with Polaroid positive/ negative Landfilm, type 55, through a yellow Wratten gelatin filter No. 8 (cutoff=460 nm) (exposure time 40 sec. at  $f4.5$ ). Gels were stained for protein for 1 hr in solution of 0.125% (w/v) Coomassie blue R-250 in 10% acetic acid and 40% methanol. Gels were destained in this same solution from which Coomassie blue was omitted.

Polaroid negatives of fluorescent gels and dry stained gels were scanned with a laser densitometer (Pharmacia-LKB Ultrosan XL). The bands were quantified by evaluating areas or height of the peaks with Gelscan XL software.

## EXAMPLE 27

### Reduction of Toxins and Labeling with mBBr

Reactions were carried out with 10  $\mu$ g of target toxin in a final volume of 100  $\mu$ l in 30 mM Tris-HCl buffer, pH 7.9, with 0.8  $\mu$ g thioredoxin, 0.7  $\mu$ g NTR (both from *E. coli*) and 2.5 mM NADPH. When thioredoxin was reduced by DTT, NADPH and NTR were omitted and DTT was added to 0.5

mM. Assays with GSH were performed similarly, but at a final concentration of 1 mM. For reduction by glutaredoxin, the thioredoxin and NTR were replaced by 1  $\mu$ g *E. coli* glutaredoxin, 0.38  $\mu$ g glutathione reductase (partially purified from spinach leaves), 1 mM GSH and 2.5 mM  
5 NADPH (the combination of these four components is called NADP/glutaredoxin system). Reduction by the reduced form of lipoic acid, was carried out in a volume of 100  $\mu$ l at two concentrations, 100  $\mu$ M and 200  $\mu$ M, both alone and with 0.8  $\mu$ g of thioredoxin. The mixture was incubated for 2 hr at 37°C in the case of erabutoxin b and  $\alpha$ -bungarotoxin, 1 hr at room  
10 temperature for  $\beta$ -bungarotoxin and 20 min at room temperature for phospholipase A<sub>2</sub>. After incubation, mBBR was added to 1.5 mM and the reaction continued for 15 min at room temperature. The reaction was stopped and excess mBBR derivatized by adding 10  $\mu$ l of 100 mM  $\beta$ -mercaptoethanol, 5  $\mu$ l of 20 % SDS and 10  $\mu$ l 50 % glycerol. Samples were then analyzed by  
15 SDS-polyacrylamide gel electrophoresis.

Total toxin reduction was accomplished by boiling samples for 3 min in 2 mM DTT. After cooling, the samples were labeled with mBBR and treated as before, except that all samples were again boiled for 2 min prior to loading in the gel. Dithiothreitol (DTT) and the reduced forms of thioredoxin and  
20 lipoic acid are dithiol reductants as opposed to monothiol reductants like 2-mercaptoethanol and glutathione. DTT is a synthetically prepared chemical agent, whereas thioredoxin and lipoic acid occur within the cell. Erabutoxin b was significantly reduced by the NTS, DTT and thioredoxin and reduced lipoic acid and thioredoxin. With erabutoxin b lipoic acid was shown to be  
25 more specific reductant than dithiothreitol. Dithiothreitol reduced the toxin partly without thioredoxin whereas reduced lipoic acid did not (lane 8). The results also showed that the NTS or DTT plus thioredoxin are specific reductants for  $\alpha$ -bungarotoxin and  $\beta$ -bungarotoxin.

## EXAMPLE 28

### NADP-Malate Dehydrogenase Activation

The ability of snake toxins to activate chloroplast NADP-malate dehydrogenase was carried out by preincubating 5 µg toxin with a limiting  
5   thioredoxin concentration (to restrict activation of the enzyme by the  
thioredoxin): *E. coli* thioredoxin, 0.25 µg; human, 0.9 µg; wheat, 1.15 µg;  
spinach *f* and *m*, 0.375 and 0.125 µg, respectively. Purified corn NADP-  
malate dehydrogenase, 1.4 µg, was added to a solution containing 100 mM  
Tris-HCl, pH 7.9, thioredoxin as indicated, and 10 mM DTT (final volume  
10   0.2 ml). After 25 min, 160 µl of the preincubation mixture was injected into  
a 1 cm cuvette of 1 ml capacity containing (in 0.79 ml) 100 mM Tris HCl,  
pH 7.9, and 0.25 mM NADPH. The reaction was started by the addition of  
50 µl of 50 mM oxalacetic acid. NADPH oxidation was followed by  
monitoring the change in absorbance at 340 nm with a Beckman  
15   spectrophotometer fitted with a four-way channel changer. The results of this  
experiment showed that the reduction by different reduced thioredoxins of  
erabutoxin b significantly alters the toxin's biological ability to activate  
NADP-malate dehydrogenase. The results demonstrate that, although there  
are differences in effectiveness, all thioredoxins tested function to some  
20   extent in limiting the effect of the toxin.

## EXAMPLE 29

### Proteolysis assay of Erabutoxin b

Erabutoxin b, 10 µg was incubated for 2 hr at 37°C with 30 mM Tris-HCl  
buffer pH 7.9 (total volume, 100 µl). As indicated, the buffer was  
25   supplemented with 0.8 µg thioredoxin, 0.7 µg NTR and 2.5 mM NADPH.  
When thioredoxin was reduced by DTT the NTR and NADPH were omitted  
and DTT was added to 0.5 mM. Following incubation, samples were  
digested with 0.4 and 2 µg of trypsin for 10 min at 37°C. DTT, 4.8 µl of 50  
mM solution, 5 µl of 20% SDS and 10 µl of 50% glycerol were added,

samples were boiled for 3 min, and then subjected to SDS-polyacrylamide gel electrophoresis. Gels were stained with Coomassie blue and the protein bands quantified by densitometric scanning as described above. The results of the assay are shown in Table XVI below. These results show that reduction of a snake neurotoxin (erabutoxin b) renders the toxin more susceptible to proteolysis. An extension of this conclusion would indicate that administration of reduced thioredoxin as a toxin antidote should help to destroy the toxin owing to the increase in proteolytic inactivation by proteases of the venom.

10

**Table XVI**  
**Susceptibility of the Oxidized and**  
**Reduced Forms of Erabutoxin b to Trypsin**

	<u>Treatment</u>	<u>% Erabutoxin b digested</u>	
		<u>0.4 µg trypsin</u>	<u>2 µg trypsin</u>
15	Control	0.0	34.1
	Reduced, NTS	21.1	57.8
	Reduced, DTT	3.1	40.6
	Reduced, DTT + Trx	28.0	71.8

20 Erabutoxin b, 10 µg was preincubated for 2 hours at 37°C in 30 mM Tris-HCl buffer, pH 7.9, as follows: control, no addition; reduced by *E. coli* NADP/thioredoxin system (NTS), thioredoxin, NTR and NADPH; reduced by DTT, DTT; and reduced by DTT plus thioredoxin, DTT supplemented with *E. coli* thioredoxin. After preincubation 0.4 µg and 2 µg of trypsin were added to the indicated which then were analyzed by SDS-  
25 polyacrylamide gel electrophoresis.

### EXAMPLE 30

#### Phospholipase A<sub>2</sub> Assay

Activity of the oxidized and reduced forms of the phospholipase A<sub>2</sub> component of  $\beta$ -bungarotoxin was determined spectrophotometrically following change in acidity as described by Lobo de Araujo *et al.* (1987) *Toxicon* 25:1181-1188. For reduction experiments, 10  $\mu$ g toxin was incubated in 30 mM Tris-HCl buffer, pH 7.9, containing 0.8  $\mu$ g thioredoxin, 0.7  $\mu$ g NTR and 7 mM NADPH (final volume, 35  $\mu$ l). After 1 hr incubation at room temperature, 20  $\mu$ l of the reaction mixture was added to a 1 cm  
10 cuvette containing 1 ml of assay solution (adjusted to pH 7.6) that contained 10 mM CaCl<sub>2</sub>, 100 mM NaCl, 4 mM sodium cholate, 175  $\mu$ M soybean phosphatidylcholine and 55  $\mu$ M phenol red. The reaction was followed by measuring the change in the absorbance at 558 nm in a Beckman Du model 2400 spectrophotometer. The results of this experiment demonstrated that  $\beta$ -  
15 bungarotoxin loses most of its phospholipase activity when reduced by thioredoxin. The results are consistent with the conclusion that the administration of reduced thioredoxin following a snake bite would help detoxify the toxin by eliminating phospholipase A<sub>2</sub> activity.

### EXAMPLE 31

#### 20 $\alpha$ -Bungarotoxin binding to Acetylcholine Receptor

$\alpha$ -Bungarotoxin binding was assayed with cultured mouse cells by using radiolabeled toxin (Gu, Y. *et al.* (1985) *J. Neurosci.* 5:1909-1916). Mouse cells, line C<sub>2</sub>, were grown as described by Gu *et al* (Gu, Y. *et al.*, *supra.*) and plated in 24-well plastic tissue culture plates (Falcon) at a density of  
25 about 3000 cells per well. Growth medium was replaced by fusion medium after 48 hr and again after 96 hr. Cultures were used for assay after an additional 2 days growth.

$\alpha$ -Bungarotoxin binding was determined with cells subjected to three different treatments: [A] 10 nM  $\alpha$ -bungarotoxin<sup>125</sup>I (262 Ci/mmol) was preincubated 2 hr at 37°C in 200  $\mu$ l of phosphate-saline buffer (0.15M NaCl in 10 mM Na<sub>2</sub>HPO<sub>4</sub>/NaH<sub>2</sub>PO<sub>4</sub> pH 7.2) with 4  $\mu$ g thioredoxin, 3.5  $\mu$ g NTR (both from *E. coli*) and 6.25 mM NADPH. In certain cases, the NTR and NADPH were replaced by 1.25 mM DTT. After 2 hr incubation, the mixture was transferred to a well containing mouse cells, washed two times with phosphate-saline, and incubated for 2 hours at 37°C. [B] After washing the cells two times with phosphate-saline buffer, 10 nM  $\alpha$ -bungarotoxin<sup>125</sup>I (in 200  $\mu$ l of phosphate-saline) was added per well. Following a 2 hr incubation at 37°C, cells were washed again with phosphate-saline buffer to remove unbound toxin. As indicated, 200  $\mu$ l saline, supplemented with 0.68 mM CaCl<sub>2</sub>, 0.49 mM MgCl<sub>2</sub>, 4  $\mu$ g thioredoxin, 3.5  $\mu$ g NTR and 6.25 mM NADPH were added to the well. The plate was incubated 2 hr at 37°C. NTR and NADPH were omitted from treatment with DTT which was added at 1.25 mM. [C] After washing cells twice with phosphate-saline buffer, 200  $\mu$ l of a solution containing 4  $\mu$ g thioredoxin, 3.5  $\mu$ g NTR and 6.25 mM NADPH, were added to each well. In some cases, NTR and NADPH were replaced with 1.25 mM DTT. The plate was incubated for 2 hr at 37°C. Cells were then washed twice with phosphate-saline buffer to remove the added reductant. Phosphate-saline buffer, 200  $\mu$ l, containing 0.68 mM CaCl<sub>2</sub> and 0.49 mM MgCl<sub>2</sub> and 10 nM  $\alpha$ -bungarotoxin<sup>125</sup>I was added to each well. Incubation was continued for 2 hr at 37°C. The results of this assay are shown in Table XVII. This experiment shows that when reduced by thioredoxin,  $\beta$ -bungarotoxin can no longer bind to the acetylcholine receptor. When extended to the whole animal, the thioredoxin-linked reduction mechanism would result in detoxification by eliminating binding of the toxin to its target receptor.



Each  $\alpha$ -bungarotoxin binding assay was done in triplicate. Nonspecific binding was measured by adding 100-fold excess unlabeled  $\alpha$ -bungarotoxin to the incubation mixture. After the incubation period, the cells in all cases were washed with phosphate-saline to remove unbound toxin. The amount of  
5 toxin bound was determined by solubilizing the cells in 0.1 M NaOH and measuring radioactivity in a gamma counter.

**Table XVII**  
**Binding of  $\alpha$ -Bungarotoxin to the**  
**Acetylcholine Receptor of Mouse Cells**

		<u>%Binding</u>
5	<u>Treatment A</u>	
	Toxin + Reductant $\xrightarrow[2 \text{ hr}, 37^\circ\text{C}]{} \xrightarrow[2 \text{ hr}, 37^\circ\text{C}]{} \text{cells} \xrightarrow{\text{STOP}}$	
	No reductant	100.0
	NTS	0.0
10	DTT plus Thioredoxin	0.0
	NTS minus NTR	63.0
	NTS minus Thioredoxin	78.0
	NTS minus NADPH	101.0
	<u>Treatment B</u>	
15	Toxin + Cells $\xrightarrow[2 \text{ hr}, 37^\circ\text{C}]{} \xrightarrow[2 \text{ hr}, 37^\circ\text{C}]{} \text{wash cells} \xrightarrow[+ \text{ reductant}]{\text{STOP}}$	
	No reductant	100.0
	NTS	78.0
20	DTT plus Thioredoxin	76.0
	<u>Treatment C</u>	
	Cells + Reductant $\xrightarrow[2 \text{ hr}, 37^\circ\text{C}]{} \xrightarrow[2 \text{ hr}, 37^\circ\text{C}]{} \text{wash cells} \xrightarrow[+ \text{ toxin}]{\text{STOP}}$	
25	No reductant	100.0
	NTS	68.7
	DTT	85.0
	DTT plus Thioredoxin	68.8

*E. coli* NTS: thioredoxin, NTR and NADPH

### EXAMPLE 32

#### Example for Detoxification in an Animal

- Detoxification of snake neurotoxins is determined by subcutaneous injection into mice. Assays are done in triplicate. Prior to injection, the toxin is
- 5 diluted in phosphate-saline buffer (0.15 M NaCl in 10 mM  $\text{Na}_2\text{HPO}_4/\text{NaH}_2\text{PO}_4$  pH 7.2) at concentrations ranging up to twice the  $\text{LD}_{50}$  dose. ( $\text{LD}_{50}$  is defined as that dose of toxin that kills 50% of a given group of animals.) For toxicity tests, the following neurotoxin concentrations correspond to the  $\text{LD}_{50}$  (per g mouse): erabutoxin b, 0.05  $\mu\text{g}$  - 0.15  $\mu\text{g}$ ;  $\alpha$ -
- 10 bungarotoxin, 0.3  $\mu\text{g}$ ; and  $\beta$ -bungarotoxin, 0.089  $\mu\text{g}$ . At 5, 10, 30, 60 minutes and 4, 12 and 24 hr after injection, separate groups of the challenged mice are injected (1) intravenously, and (2) subcutaneously (multiple local injections around the initial injection site). The thioredoxin is reduced with:
- (1) the *E. coli* NADP-thioredoxin system, using 0.08  $\mu\text{g}$  thioredoxin, 0.07
- 15  $\mu\text{g}$  NTR and 25 nanomoles NADPH; (2) Thioredoxin plus 1-2 nanomoles of reduced lipoic acid, using 0.08  $\mu\text{g}$  *E. coli* or 0.20  $\mu\text{g}$  human thioredoxin, and (3) using 0.08  $\mu\text{g}$  *E. coli* or 0.20  $\mu\text{g}$  human thioredoxin with 5 nanomoles dithiothreitol (concentrations are per  $\mu\text{g}$  toxin injected into the animal; all solutions are prepared in phosphate-saline buffer).
- 20 The effect of thioredoxin on detoxification is determined by (1) comparing the  $\text{LD}_{50}$  with the control group without thioredoxin; (2) following the extent of the local reaction, as evidenced by necrosis, swelling and general discomfort to the animal; (3) following the serum levels of creatin kinase, an indicator of tissue damage. Creatin kinase, which is released into the blood
- 25 as a result of breakage of muscle cells, is monitored using the standard assay kit obtained from Sigma Chemical Co. (St. Louis, MO).

The symptoms of snake bite are multiple and depend on a variety of factors. As a consequence, they vary from patient to patient. There are, nonetheless,

common symptoms that thioredoxin treatment should alleviate in humans. Specifically, the thioredoxin treatment should alleviate symptoms associated with neurotoxic and related effects resulting from snake bite. Included are a decrease in swelling and edema, pain and blistering surrounding the bite;  
5 restoration of normal pulse rate; restriction of necrosis in the bite area; minimization of the affected part. A minimization of these symptoms should in turn result in improvement in the general health and state of the patient.

### **Reduction of Food and Pollen Proteins and Allergens**

The invention provides a method for chemically reducing the disulfide bonds  
10 in major allergen proteins particularly food and pollen allergen proteins and for decreasing or eliminating the allergenicity that occurs when foods containing those proteins are ingested or pollens containing those proteins are inhaled, ingested or come in contact with mucus membranes.

One method of decreasing or eliminating an allergic reaction involves  
15 subjecting an animal over a period of time to immunotherapy with varying doses of an allergen that has been reduced by reduced thioredoxin. The allergen may be a pollen protein or an allergen protein found in plants such as poison oak or in animals such as dust mites.

The invention also provides a method for increasing the digestibility of  
20 allergen proteins and therefore food but also other proteins such as pollens that may be swallowed as well as inhaled. The disulfide bonds of the proteins were reduced to the sulfhydryl (SH) group by thioredoxin. The other major cellular thiol reductant, glutathione, was ineffective in this capacity. The proteins are allergenically active in the oxidized (S-S) state;  
25 when treated with reduced thioredoxin they are reduced (SH state) and lose allergenicity. Thioredoxin achieves this reduction when activated (reduced) either by NADPH via the enzyme NADP-thioredoxin reductase

(physiological reducing system), or by dithiothreitol (DTT), a synthetic chemical reductant or by lipoic acid a physiological reductant. While use of NADPH as a reductant is often preferred in almost all instances, physiologically acceptable lipoic acid may be used and DTT may also be  
5 acceptable for the treatment of some proteins. If the allergen contains thioredoxin then NADPH/NTR or lipoic acid may be used in some cases without added thioredoxin.

Presumably the proficiency of reduced thioredoxin to decrease or eliminate the allergenicity caused by the food depends upon the ability of the reduced  
10 thioredoxin to reduce the intramolecular disulfide bonds in the allergenic proteins in the food. Also, an allergen that has had its disulfide bonds reduced by thioredoxin will cause less allergenicity (*i.e.*, it will be hypoallergenic) but will still retain its ability to be an effective immunotherapeutic agent.

15 Food proteins that have intramolecular disulfide bonds can be reduced and the allergenicity of foods containing these proteins can be at least decreased and the digestibility increased by reduced thioredoxin when the food is treated with the thioredoxin for an effective period of time at an effective temperature. The effective temperature for incubating the food is from about  
20 4°C to 55°C but any temperature which will allow the reduction of the disulfide bonds is acceptable. The effective time for incubation is from about 20 min to 2 hrs but again any time that allows for reduction of the disulfide bonds without significantly degrading food quality is acceptable. For example, the proteins may also be reduced and remain reduced by incubation  
25 at about 4°C for about 48 hrs.

Examples of the foods that will exhibit reduced allergenicity and increased digestibility upon treatment with reduced thioredoxin are beef, milk, soy, egg, rice, wheat and nuts.

A preferred method for decreasing the allergenicity and increasing the digestibility of food ingested by a mammal is to pretreat or incubate the food using the components in the NADP-thioredoxin system (NTS). In general, the effective amounts of the components range from about 0.01 mg to 4.0 mg, preferably about 0.10 mg to 2 mg of thioredoxin; from about 0.01 mg to 4 mg, preferably from about 0.20 mg to 2 mg of NTR and from about 1 micromole to 250 micromoles, preferably about 25 micromoles to 100 micromoles of NADPH for every 1.0 gm of protein in the food. Of course, the effective amounts of each component will vary depending upon the amount of the other components, the time and temperature of incubation, as well as the particular food and the amount of innate thioredoxin and its reductants in that food. For example, the above amounts of thioredoxin were determined on the basis of using 1 mg NTR and 100 micromoles NADPH per gram of food protein. The range of NADPH was determined on the basis of using 1 mg NTR and 1 mg thioredoxin. The appropriate component amounts may also be affected by prior treatment of the food and the type of animal and its condition.

The NADP/thioredoxin system is also able to reduce the disulfide allergens of airborne allergens such as ragweed pollen and grasses and dermal contact allergens such as dust mites and thereby alleviate the IgE response of an animal to those allergens when the animal receives these reduced allergens systemically.

The effective temperature for treating an allergen such as a pollen allergen with the NADP/thioredoxin system is from about 4°C to 55°C but again any

temperature which will allow the reduction of the disulfide bonds is acceptable. The effective time for incubation is similar to the time for treating food (from about 20 min to about 1 hr) but again any time that allows for reduction of the disulfide bonds without significantly degrading the protein is acceptable. For example, the proteins may also be reduced and remain reduced by incubation at about 4°C for about 24 hrs.

Examples of allergens that will exhibit reduced allergenicity and increased digestibility upon treatment with reduced thioredoxin include disulfide containing allergenic proteins such as ragweed pollens, grass pollens, poison oak and ivy allergens and dust mite allergen protein.

A preferred method for decreasing the allergenicity and alleviating the IgE response of a mammal to an allergen is to pretreat or incubate the allergen using the components in the NADP-thioredoxin system (NTS). In general, the effective amounts of the components for airborne and contact allergens like food range from about 0.1 mg to 6.0 mg, preferably about 0.1 mg to 4.0 mg of thioredoxin; from about 0.1 mg to 8.0 mg, preferably from about 0.2 mg to 7.0 mg of NTR and from about 1 micromole to 500 micromoles, preferably about 25 micromoles to 400 micromoles of NADPH for every 1 gm of protein of the allergen. Again, the effective amounts of each component will vary depending upon the amount of the other components, the time and temperature of incubation, as well as the particular allergen protein or allergen extract and the amount of innate thioredoxin and its reductants in that allergen extract. The above amounts of thioredoxin were determined on the basis of using 1 mg NTR and 100 micromoles NADPH per gram of allergen protein. The range of NADPH was determined on the basis of using 1.0 µg NTR and 1.0 µg thioredoxin per 1.0 gm pollen protein.

Varying doses of the thioredoxin reduced allergen extract are then injected into the animal over a prescribed period of time.

The appropriate component amounts may also be affected by prior treatment of the allergen, the particular allergen and the type of animal and its  
5 condition.

Other features and advantages of the invention with respect to decreasing the allergenicity and increasing the digestibility of a particular allergen can be ascertained from the following examples.

### EXAMPLE 33

#### 10           **Treatment of Milk, Soy, Wheat and Beef with Thioredoxin**

For this study a 1 to 5 physiological saline (PBS) dilution of a stock, 1:20 weight/volume (w/v), cow's milk allergenic extract (Catalogue No. 3390JG, Miles, Inc., Elkhart, Indiana), a 1 to 10 PBS dilution of a stock, 1:10 w/v, soy allergenic extract (Catalogue No. 3597ED, Miles, Inc., Elkhart, Indiana)  
15 and a 1 to 5 PBS dilution of a stock, 1:10 w/v, wheat allergenic extract (Catalogue No. 3708ED, Miles, Inc., Elkhart, Indiana) were prepared. A 1 to 5 PBS dilution of a stock, 1:10 w/v, commercial beef allergenic extract (Catalogue No. 3078JF, Miles, Inc., Elkhart, Indiana) was also prepared.

In the case of cow's milk, 0.1 ml of the dilution was treated with the  
20 NADP/thioredoxin system (NTS) which comprised incubating the allergen in this instance with 4.8 micrograms thioredoxin, 4.2 micrograms NTR and 1 mM NADPH (final volume, 0.2 ml). A second sample was also prepared using 0.1 ml of the diluted milk allergen incubated with 1.5 mM DTT and 4.8 micrograms thioredoxin (final volume, 0.2). In all the allergen studies,  
25 including the studies in this Example and the ones following, the thioredoxin



and NTR used were purified as previously described from *E. coli* that had been transformed to overproduce those proteins (de la Motte-Guery, F. *et al.* (1991) *Eur. J. Biochem.* 196:287-294, and Russel, M. *et al.* (1988) *J. Biol. Chem.* 263:9015-9019). For soy and wheat, 0.05 ml of the dilutions were  
5 incubated with 2.4 micrograms thioredoxin, 2.1 micrograms NTR and 1 mM NADPH. In addition, an identical treated control sample of PBS was prepared. DTT treated soy and wheat samples were also prepared using 0.05 ml of the separate allergen dilutions incubated with 1.5 mM DTT and 2.4 micrograms thioredoxin. The final volume for the control and all the soy and  
10 wheat samples was 0.1 ml. The milk and wheat preparations were incubated at room temperature for 25 min while the soy preparation was incubated at 37°C for 1 hr and 25 min. With the beef, 0.05 ml of the dilution was incubated with 2.4 micrograms thioredoxin, 2.1 micrograms NTR and 1 mM NADPH (final volume, 0.1 ml) for 25 min at 37°C. Another 0.05 ml  
15 sample was treated the same way but at room temperature. Following incubation 1 ml dilutions ranging from  $1 \times 10^3$  to  $1 \times 10^6$  or from  $1 \times 10^3$  to  $1 \times 10^7$  were prepared for each treated food extract preparation. The diluted samples were used for testing within 30 min.

#### EXAMPLE 34

20        **Determination of the Reduction of Food Allergens by the  
         NADP/Thioredoxin System and Increase in Proteolysis Using the  
         mBBr Fluorescent Labeling/SDS-Polyacrylamide Gel Electrophoresis**

##### **Method**

For this study a 1 to 2.5, 1 to 5 and 1 to 1.5 dilution with PBS of the stock  
25 cow's milk, beef and wheat extracts described in Example 33 were respectively prepared. To 50 microliters of these dilutions were added 2.4 micrograms thioredoxin, 2.1 micrograms NTR and 1 mM NADPH (final volume, 0.1 ml). The controls consisted of 50 microliters of the particular

diluted extract and 50 microliters of PBS. The preparations were incubated for 25 min at room temperature and also at 37°C. After incubation, 8 µl of 80 mM mBBr was added and the reaction continued for 15 min at room temperature. The reaction was stopped and excess mBBr derivatized by adding 10 µl of 100 mM β-mercaptoethanol, 10 µl of 20% SDS and 10 µl of 50% glycerol. The samples were analyzed by the mBBr/SDS-polyacrylamide gel electrophoresis technique previously described. The results showed that the NTS effectively reduced the proteins in the allergenic extracts at both room temperature and 37°C. In an additional study, where a PBS dilution of the soy stock extract described in Example 33 was similarly treated with the NTS, an analysis using the mBBr labeling/SDS-PAGE method showed that thioredoxin also reduced the soy proteins. However, when soy, cow's milk, wheat, egg and beef allergenic proteins were incubated with glutathione, glutathione reductase and NADPH, there was minimal or no reduction of those treated allergenic proteins.

A PBS dilution of a commercial rice allergenic extract (Catalogue No. 3549ED, Miles, Inc., Elkhart, Indiana) is also similarly incubated with the NTS and analyzed using the mBBr/SDS-PAGE technique to show that reduced thioredoxin reduces rice allergen proteins.

In a separate study, it was also observed that food allergen proteins from the commercial extracts described in Example 33 that had been reduced by the NTS and were further incubated with trypsin had an increased susceptibility to proteolysis over controls that had not been treated with the NTS. The analysis of the reduction was done using the mBBr/SDS-PAGE techniques. Further in this study, when 10 µg of an NTS reduced purified milk allergen protein, β-lactoglobulin (Sigma Chemical Co.), was treated with 2 µg of trypsin, proteolysis was 100% as compared with only 50% for the identically trypsin treated oxidized β-lactoglobulin. When 10 µg of another purified

milk allergen, oxidized  $\alpha$ -lactalbumin (Sigma Chemical Co.), was similarly treated with 2  $\mu$ g trypsin, there was no noticeable proteolysis. However, 10  $\mu$ g of purified  $\alpha$ -lactalbumin reduced by the NTS was proteolyzed 80% by trypsin. Also 10  $\mu$ g of  $\alpha$ -lactalbumin reduced by 0.8  $\mu$ g of thioredoxin and 5 0.5 mM DTT was 100% proteolyzed by 2  $\mu$ g of trypsin.

### EXAMPLE 35

#### Reduction of Egg White Proteins

Dried chicken egg white was purchased from Sigma Chemical Co. About 80% of the total proteins in egg white are allergens. A solution of 20 mg/ml  
10 egg white was resuspended in PBS. Since not all the material was dissolved, it was centrifuged at 14,000 RPM for 2 min. The soluble egg white proteins were used for the reduction study using mBBBr fluorescent labelling and SDS-polyacrylamide gel electrophoresis analysis. The treatments used were the control (no reductant), NTS, DTT plus thioredoxin, reduced glutathione  
15 (GSH) and reduced glutathione/glutathione reductase/NADPH. Reactions were carried out in PBS with 23 microliters of the soluble proteins from the 20 mg/ml egg white suspension in a final volume of 100 microliters. In the NTS, 7.5 mM NADPH, 2.4 micrograms thioredoxin and 2.1 micrograms NTR were used. When thioredoxin was reduced by DTT, NADPH and NTR  
20 were omitted and DTT was added to 1.5 mM. The GSH concentration used was 3 mM. For reduction by the GSH/GR/NADPH system, 3 mM GSH, 4 micrograms GR and 7.5 mM NADPH were used. The mixture was incubated for 1 hr at room temperature. After incubation, 7 microliters of 80 mM mBBBr was added and the reaction continued for 15 min at room  
25 temperature. The reaction was stopped and excess mBBBr was derivatized by adding 10 microliters of 100 mM mercaptoethanol, 10 microliters of 20% SDS and 10 microliters of 50% glycerol. Samples were then analyzed by SDS-polyacrylamide gel electrophoresis. The results of this experiment

showed that the NTS and DTT plus thioredoxin are very effective in reducing egg white proteins which are about 80% allergens. GSH or GSH/GR/NADPH showed the same level of reduction as the control and therefore is an ineffective reductant of egg white proteins.

5

### EXAMPLE 36

#### Sensitization of Animals for Allergenicity Studies

The animals used in this study were atopic dogs born to different pairs of littermates from an in-bred colony of high IgE-producing spaniels.

10 A litter of 9 pups (4 males, 5 females) was born to an in-bred IgE-responder bitch sired by her brother. On newborn day 1, for the cow's milk, soy and rice studies, a nine-pup litter was divided into two groups: Group I of 5 pups was injected subcutaneously (SQ) in the right axilla with 1  $\mu$ g the commercial soybean extract described in Example 33 in 0.2 ml alum; Group II of 4 pups was injected SQ in the right axilla with 1  $\mu$ g of commercial dried cow's milk  
15 extract (described in Example 33) solubilized in 0.2 ml saline and 0.2 ml alum. All 9 pups were also given 1  $\mu$ g of a stock 1:10 w/v rice allergenic extract (Catalogue No. 3549ED, Miles, Inc. Elkhart, Indiana) in 0.2 ml alum SQ in the left axilla.

20 At ages 3, 7 and 11 weeks, all pups were vaccinated with 0.5 ml of live attenuated distemper-hepatitis vaccine (Pitman-Moore, Washington's Crossing, PA) in the shoulder SQ. Two and 9 days after each vaccination, they were given the same food antigens that they received in the neonatal period, e.g., 5 dogs in Group I received 1  $\mu$ g soybean extract in 0.2 ml alum, 4 dogs in Group II received 1  $\mu$ g cow's milk in 0.2 ml alum, all 9  
25 pups received 1  $\mu$ g rice extract in 0.2 ml alum SQ in the right and left axilla, respectively.

A  $^{125}\text{I}$ -labelled rabbit anti-canine IgE serum (Frick, O.L. *et al.* (1983) *Am. J. Vet. Res.* 44: 440-445) was used in a RAST assay. Cyanogen-bromide activated filter paper discs were reacted with 100  $\mu\text{g}$  of either soy, cow's milk or rice antigen, as in standard RAST protocol (Wide, L. *et al.* (1967) *Lancet* 2:1105-1107). A pool of newborn canine cord sera from non-atopic mongrel pups was used as a negative control or baseline cpm.

The pups were nursed for 6 weeks and weaned onto regular Puppy Chow (Ralston-Purina Company, St. Louis, MO) which included the sensitizing proteins, soybean meal, dried whey, and rice hulls; they were fed once/day and given water ad lib, under veterinary care and supervision at the University of California, Davis, Animal Resources Services, School of Veterinary Medicine. After 6 months, they were fed regular Field & Farm Dog Chow.

Between 3 and 4 months of age, when it was found that the pups were making IgE antibodies to a particular food antigen, all 9 pups in the litter were given double-blinded, 240 ml of either soy or cow's milk infant formula, tofu, rice gruel, or vanilla-flavored Vivonex protein hydrolysate (Norwich-Eaton Co., Norwich, CT) in the early morning. Abdominal girth was measured at umbilical level before the challenge and at 2 hours intervals during the day. They were observed closely by a veterinary technician for clinical signs of itching or rash, vomiting and frequency and character of stools, and for cough or respiratory distress and nasal discharge. They were monitored for such signs of reaction for 4 days. The next challenge feeding was given 7 days later.

All 9 pups in the litter gained weight and developed normally with no medical problems. They had no diarrhea or other gastrointestinal signs unless they were challenged with the food they had been sensitized to. Also

no skin or respiratory abnormalities occurred. There were no untoward reactions to the vaccinations or immunizations.

Canine IgE-RASTs to the 3 food proteins were followed at fortnightly intervals with venous blood sampling.

- 5 Significantly more IgE-anti-food antibodies were produced by the corresponding milk ( $p < .05$  when challenged at 4 months) and soy ( $p < .0005$  for the first 3 months) immunized animals than by controls. The average titer for IgE-anti-rice-antibodies rose at 5-10 week, plateaued and then rose again sharply after 20-30 weeks of age. Also statistically significant reactions  
10 of diarrhea and abdominal bloat occurred in the soy and milk immunized animals when they were respectively fed soy and milk in their chow.

- Another group of 8 dogs allergic to cow's milk, soy, wheat and beef were also developed in a manner similar to the method described above. Four littermate pups from another litter in the described in-bred colony of high  
15 IgE-producing spaniels were injected SQ in the right axilla with 1  $\mu$ g each of the stock cow's milk, soy, wheat and beef allergenic extracts described in Example 33. Four pups from the same litter acted as controls. All eight pups were also given 1  $\mu$ g of the rice extract in 0.2 ml alum SQ in left axilla. The pups were vaccinated as above and at two and nine days after each  
20 vaccination, were given the same food antigens in the same amount that they received as neonates. As above, they were fed foods which contained the appropriate sensitizing proteins (*i.e.*, cow's milk, soy, beef and wheat) in a similar schedule. As with the previous soy and milk allergic animals, these immunized dogs produced significantly more anti-specific food IgE  
25 antibodies (including wheat and beef antibodies) than the controls. Again statistically significant reactions of diarrhea and abdominal bloat occurred in

the immunized animals when they were challenged with food containing wheat, beef, soy or cow's milk.

### **EXAMPLE 37**

#### **5            Skin Test Determinations of the Decrease of Allergenicity               in Food Allergens Treated with Reduced Thioredoxin**

Aliquots of 100 microliters of each dilution of the thioredoxin treated food allergen dilutions described in Example 33 were injected intradermally on the abdominal skin of the appropriate sensitized dogs described in Example 36 (*e.g.*, cow's milk sensitized dogs were injected with the cow's milk  
10 dilutions). Prior to the allergen dilution injections, the dogs' forelimb veins were injected with 4 ml of 0.5% Evans blue dye. Dogs exhibiting an allergenic reaction developed blue colored wheals in the area of the allergen injection. After 10 minutes of development the size of the wheal (length and width) were measured. The observed size of the wheal and the dilution end  
15 point following injection of a concentration range of each allergen preparation were used as the allergenicity indicator. It was found that thioredoxin treatment gave 50% protection with  $1 \times 10^5$  dilution of soy, full protection with a  $3 \times 10^4$  dilution of milk, full protection with a  $1 \times 10^6$  dilution of wheat and at least partial protection with a  $1 \times 10^5$  dilution of beef  
20 (see Figs. 1, 2, 3, 4 and 5 respectively).

### **EXAMPLE 38**

#### **Feeding Test Determinations of the Allergenicity of Food Allergens Treated with Reduced Thioredoxin**

Approximately one week prior to feeding, dogs sensitized in the manner  
25 described in Example 36 were skin tested intradermally as in Example 36 with the appropriate food allergen using the commercial allergenic extracts described in the previous examples. Based on these results, animals were

- separated into "control" and "thioredoxin-treated" groups. Each group was made up of representatives with complementary sensitivities - *i.e.*, an equal number of strong, medium and weak reactors was selected for each group. Unless indicated otherwise, the groups were comprised of 3 animals (6  
5 animals per experiment). For 3 days before and 5 days after the feeding challenge, dogs were maintained on a Hill's Prescription Diet Canine d/d diet (Hill's Division of Colgate Palmolive Co., Topeka, Kansas). Dogs were observed throughout this period for clinical symptoms such as retching and vomiting. In addition, stools were monitored and scored as an indicator of  
10 the allergenic response to the food being tested. The dogs' stools were counted for 3 days before and 3 days after the allergenic diet challenge and their consistency was indicated by a number: 1 = firm, 2 = soft, and 3 = runny. The stool score was then calculated by multiplying the number of stools time the consistency factor. As an indicator of the allergenic response  
15 to the food allergens being tested, the final net average stool score per day for each group ("control", or "thioredoxin treated") was calculated by subtracting the average stool score per day before from that after the allergenic diet challenge. A higher net average stool score per day represents a stronger allergenic response.
- 20 The procedures used for preparing and administering the diets is given below. Unless indicated otherwise, reactions were carried out at room temperature.

### Soy

- Commercial soy formula (Isomil supplemented with iron, Ross Laboratories, Columbus, Ohio), 1.026 kg, was dissolved in 3 l of water. The solution was  
25 separated into two lots of 1.925 l, one used as the control and the other treated with the NADP/thioredoxin system (NTS) as follows. A mixture containing 45 micromoles NADPH, 564 micrograms NADP-thioredoxin reductase (NTR) and 1.125 mg thioredoxin (all dissolved in 30 mM Tris-HCl



buffer, pH 7.9) was preincubated for 5 min and then added to one of the 1.925 l formula lots (henceforth the "thioredoxin-treated" lot). After adding the thioredoxin system, incubation was continued for an additional hour with constant stirring. An equal amount of buffer was added to the control lot.

- 5 After incubation, 600 ml of formula was fed to each of the 3 dogs of the assigned group. The portions fed to the animals were equivalent to 25.0 gm of soy protein prior to incubation.

### **Wheat**

- Unbleached flour, 1.5 kg, was added to 3 l of water, previously heated to 10 37°C, in a gallon-size Waring blender. After 1 min blending, the flour suspension was divided equally into two lots, one was used as the control and the other treated with the NADP/thioredoxin system as follows. A mixture containing 45 micromoles NADPH, 564 micrograms (NTR) and 1.125 mg thioredoxin, all dissolved in 30 mM Tris-HCl buffer, pH 7.9 was 15 preincubated for 5 min and then added to one lot (henceforth the "thioredoxin-treated" lot). An equivalent amount of buffer was added to the control lot. Both preparations were incubated 1 hr at 37°C with frequent stirring. A volume of 600 ml was removed from each preparation, mixed with one can (15-3/4 oz) d/d, wheat free, rice/lamb based, dog food and fed 20 to the 3 dogs of the assigned lot. Again these portions were equivalent to 25 gm of wheat protein. In the experiment with 8 dogs, the flour was increased to 2.0 kg and the procedure scaled up accordingly.

### **Milk**

- Preparations of dried CARNATION that had been reconstituted with water 25 were similarly incubated with the NADP/thioredoxin system. As before 3 of the dogs received untreated and 3 received the treated milk. The final portions that the dogs received were equivalent to 10 gm of milk protein.

## Results

The levels of components of the NADP/thioredoxin systems that were used were significantly higher than in the dough studies described above in Example 15. In this feeding study, preliminary trials indicated that higher levels of these compounds were required to reduce the allergenic proteins as determined *in vitro* by the mBBR/SDS-polyacrylamide gel procedure. The amounts of each component of the NADP/thioredoxin system used for each dog per gm protein in the feeding trials is indicated below relative to the amounts used in the baking tests:

10		Wheat Flour, Milk, <u>Soy Formula*</u>
	Thioredoxin	5-X
	NTR	5-X
	NADPH	2-X
15	<hr/>	

\*Amounts indicated are relative to those used in the above described baking tests in which 3 micrograms thioredoxin, 1.5 micrograms NTR and 0.3 micromoles NADPH were added per gram of flour protein. In the baking tests, loaves were baked with approximately 200 g flour or approximately 20 gm of flour protein. For the feeding experiments, food preparations were incubated with components of the NADP/thioredoxin system for one hour either at room temperature (milk and soy) or 37°C (wheat).

Based on "bedside symptoms" (vomiting and retching) as well as "stool score" (see Fig. 6), thioredoxin treatment was found to decrease the allergenic response of the dogs to the soy and wheat formulas. The allergenicity of the milk formula may also be decreased by treatment with the NTS. It should be noted that while the thioredoxin and NTR used were from *E. coli*, thioredoxins from other sources such as yeast and thioredoxin h and m may also be used.

### EXAMPLE 39

#### **Determination of the Increased Digestibility and Decreased Allergenicity of Thioredoxin Treated Milk Proteins and Raw Cow's Milk**

Milk allergy is caused by several proteins -- $\alpha$ -lactalbumin, serum albumin,  
5 caseins, and particularly  $\beta$ -lactoglobulin (BLG).

This example showed that the disulfide bonds of milk proteins and allergens were selectively and specifically reduced by thioredoxin. Once reduced, the most active of these allergens (BLG) showed not only a decrease in allergenicity, but also a striking increase in digestibility. The susceptibility of  
10 other milk disulfide proteins to pepsin also increased to some extent following reduction by thioredoxin.

#### **Materials and Methods**

##### **Allergen source**

Raw cows milk was obtained from the experimental farm, University of  
15 California at Davis. Pure  $\beta$ -Lactoglobulin A and B and an 80% mixture of the two forms were purchased from Sigma Chemical Co., St. Louis, MO.

##### **Animals**

Dogs obtained from the same colony of inbred, high IgE-producing atopic dogs used in Examples 36-38 were sensitized and maintained at the Animal  
20 Resources Service, School of Veterinary Medicine, University of California, Davis. These dogs, sensitized at birth, have been selected for a genetic predisposition to allergy and have a 15-year history of food and pollen hypersensitivity.

##### **Chemicals and enzymes**

25 Reagents for sodium dodecyl sulfate/polyacrylamide (SDS/PAGE) were obtained from Sigma, Boehringer Mannheim, Indianapolis, IN and Biorad

Laboratories, Hercules, CA. DTT was purchased from Boehringer Mannheim and monobromobimane (mBB<sub>r</sub>), from Calbiochem, San Diego, CA. Thioredoxin and NTR were purified from *E.coli* strains overexpressing the proteins. Glutathione reductase was purified from spinach leaves by the  
5 same procedure used for spinach NTR. Pepsin from porcine stomach mucosa, NADPH and other biochemical reagents were purchased from Sigma.

#### **Food sensitization of atopic dogs**

Newborn pups from two litters of the atopic dog colony, designated CGB and GCB, were injected subcutaneously at day 1 with 1  $\mu$ g each of wheat, cows  
10 milk and beef extract (Miles, Inc., Elkhart, Indiana, described in Example 33) in 0.2 ml of alum. A third litter, designated CBB, was injected with a soy extract (Miles, Inc. described in Example 33) in addition to these same allergens. Procedures for the sensitization, testing and maintenance of the pups were substantially the same as those described in Example 36.

#### **15 Skin tests**

About 3 min prior to skin testing, each dog received 4-5 ml of filtered 0.5 % Evans blue dye solution (equivalent to 0.2 ml of 0.5 % Evans blue dye per kg of weight) through a cephalic vein to enhance assessment of the cutaneous IgE antibodies. Serial dilutions of 100  $\mu$ l of each sample were injected  
20 intradermally on the abdominal skin to establish the titer. After 15-20 min, the allergic response was determined by measuring the size of the blue wheal reaction (maximum length and width). An appropriate negative control (buffer diluted in physiological saline) was included for each animal tested. Repeated tests with thioredoxin, NTR, NADPH and pepsin alone were  
25 consistently negative.

### Protein assay

Protein concentration was determined by the Bradford method using bovine gamma globulin as the standard. Concentration of pure BLG was determined by its absorbance at 278 nm using a calculated molar extinction coefficient of  
5 16800 (Gill, S.C. *et al.* (1989), *Anal. Biochem.* 182:319-326).

### Protein modeling

A model of BLG structure, determined by Brownlow *et al.* at 1.8 Å resolution, was provided by the Protein Data Bank, Brookhaven National Laboratory (Brownlow, S. *et al.* (1997), "Bovine beta-Lactoglobulin at 1.8  
10 Resolution-Still an Enigmatic Lipocalcin", *Structure* 5:481-495). A model of the protein with single mutated C160S (partly reduced) and double mutated C160S- C106S (fully reduced), was built by the Swiss-Model program (Peitsch, M.C. (1996), "ProMod and Swiss-Model: Internet-based tools for automated comparative protein modelling", *Biochem. Soc. Trans.* 24:274-  
15 279). A three dimension model of BLG was visualized by the RasMol program v2.6.

### Protein reduction

Reduction of the protein disulfide bonds was performed, in a volume of 100 µl with either: (i) the NADP/thioredoxin system (NTS), consisting of 5 µl of  
20 25 mM NADPH, 8 µl of 0.3 mg/ml *E. coli* thioredoxin (*i.e.*, 2.4 µg total thioredoxin) and 7 µl of 0.3 mg/ml *E. coli* NTR; or (ii) the NADP/glutathione system (NGS), composed of 5 µl of 25 mM NADPH, 10 µl of 30 mM reduced glutathione (GSH) and 15 µl of 0.1 mg/ml glutathione reductase. Reactions were carried out in physiological buffered saline  
25 solution (PBS; *i.e.*, 10 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.8 mM KH<sub>2</sub>PO<sub>4</sub>, 2.7 mM KCl and 137 mM NaCl, pH 7.4) containing either 10 µg of pure target protein or 50 µg of raw milk. The reaction mixtures were incubated at 4°C overnight or at 37°C and 55°C for 45 min. For complete reduction, samples were incubated

in PBS containing 5  $\mu$ l of 100 mM DTT and boiled 5 min. The reduced proteins were visualized on gels by mBBR labeling and gel electrophoresis as described below. The extent of reduction was determined by scanning the gels.

## 5 Pepsin assay

BLG, 640  $\mu$ g, or milk, 1 mg protein (*i.e.*, about 30  $\mu$ l), was incubated with or without the thioredoxin system (NTS) at either 4°C (to yield fully reduced form) or 55°C (to yield partly reduced form), under the conditions described above. The BLG, 320  $\mu$ g, or milk, 500  $\mu$ g protein, was then treated in 200  
10  $\mu$ l of simulated gastric fluid (SGF) as described by Astwood, J.D. *et al.* (1996), "Stability of food allergens to digestion in vitro", *Nature Biotechnol.* 14:1269-1273. SGF consists of 0.32% pepsin (w/v) and 30 mM NaCl adjusted at pH 1.2 with HCl (Board of Trustees (ed.) 1995, Simulated Gastric Fluid, TS., pp. 2053 in the *United States Pharmacopeia* 23, The  
15 National Formulary 18. United States Pharmacopeial Convention, Inc. Rockville, MD). The reaction mixture was incubated at 37°C and stopped by adding 0.375-fold volume of 160 mM Na<sub>2</sub>CO<sub>3</sub> (ca. pH 7) after 0, 0.25, 1, 2, 4, 8, 15, 30 and 60 min incubation. The protein mixture was then subjected to SDS-PAGE (15% gels) and stained for protein with Coomassie  
20 blue as described below. As indicated, the allergenicity of the digested samples was determined by skin test analysis.

## Feeding challenges

Reduction of 80% pure BLG (a mixture of the A and B forms) was carried out for each dog in 100 ml of water. Each gram of BLG was treated by  
25 adding an aqueous mixture of 104 mg of NADPH, 1 mg of *E. coli* thioredoxin and 1 mg of *E. coli* NTR. The reactions occurred in a shaker at 125 rpm at 37°C for 45 min. Samples were stored overnight at 4°C. The following day untreated (control) or treated BLG, 2.5 gm was mixed with a

12 oz. can of P/D food (Hills) and fed to a dog. Unchallenged animals received dog food alone without BLG while control animals received dog food with untreated BLG. Dogs were initially fed 1/4 can of untreated food to initiate gastric flow. After 15 min, they were then fed at 3 intervals  
5 separated by 15 min 1/3 of a can of food that had been mixed with 2.5 g of either untreated or thioredoxin treated BLG. During these intervals dogs were monitored and their GI response assessed. Dogs were then observed and their response recorded at hourly intervals for the next 5 hr.

### **Data analysis**

10 The digestion response of the food challenged dogs described above was assessed by assigning numbers to the timing, volume and fluidity of vomit induced by feeding. Fluidity: no vomit=0, solid vomit=1, liquid vomit=2, liquid with blood or bile vomit=3. Volume: no vomit=0, small vomit=1 and large vomit=2. Timing: delayed vomit=1, immediate vomit=2.

### **15 mBBR Labeling and Analysis of Proteins**

Sulfhydryl groups were visualized as their fluorescent mBBR derivatives. mBBR, 10  $\mu$ l of a 100 mM solution, was added to each protein sample. After 20 min of incubation, the reaction was stopped by adding 10  $\mu$ l of 100 mM 2-mercaptoethanol, 10  $\mu$ l of 20% SDS and 20  $\mu$ l of SDS/PAGE sample  
20 buffer containing 80 % (v/v) glycerol and 0.005 % bromophenol blue. Proteins were then separated by SDS/PAGE (10-20% acrylamide gradient) as described below. After electrophoresis, gels were placed in 12 % trichloroacetic acid for 1 hr for fixation and then soaked overnight or longer in 40% (v/v) methanol and 10% (v/v) acetic acid with several changes to  
25 remove excess mBBR. The destained gels were then placed under 365 nm UV light to visualize fluorescent bands. Pictures were taken by either (i) Polaroid photographs (Positive/Negative Landfilm, type 55) through a yellow

Wratten gelatin filter no. 8 with an exposure time of 45 s at  $f4.5$ , or (ii) the Nucleovision system of NucleoTech Corporation.

### SDS/PAGE

Gels (10-20% acrylamide gradient, 1.5 mm thickness or 15% acrylamide, 5 0.75 mm thickness) were prepared according to Laemmli, U.K. (1970), "Cleavage of structural proteins during the assembly of the head of bacteriophage T4", *Nature* 227:680-685. After electrophoresis, gels were stained with 0.01% Coomassie brilliant blue R-250 in 40% methanol and 10% acetic acid for 1 hr and destained overnight with a solution of 20% 10 ethanol and 10% acid acetic. Pictures were taken after destaining by either (i) Polaroid photograph (exposure time 1/15 s at  $f7$ ) or (ii) the Nucleovision system of NucleoTech Corporation.

### Sequence Analyses

The different forms of BLG (oxidized, partly and fully reduced, 10 ug) were 15 separated as mBBR derivatives by SDS-PAGE (10-20% acrylamide gradient, 1.5 mm thickness or 15% acrylamide, 0.75 mm thickness). An in-gel digestion method was used to obtain peptides containing cys residues that would allow structural characterization (Hwang, B.J. *et al.* (1996), "Internal sequence analysis of proteins eluted from polyacrylamide gels", *J. 20 Chromatogr. B. Biomed. Appl.* 686:165-175). The dried gel used as starting material was placed in water to allow swelling and the cellophane layer was removed. The appropriate BLG bands were then cut out from the reconstituted gel. Each protein band was diced into 1 to 2 mm pieces. In brief, gel pieces were dehydrated in a speedvac, rehydrated in 0.1 M Tris 25 buffer, pH 9 and 0.05% SDS, pH 9.0, containing 0.03 to 0.05  $\mu$ g Lys-C endopeptidase (Wako) and incubated overnight at 30°C. Peptides were eluted by extracting the gel twice for 2 hr with water and twice for 2 hr with 70% acetonitrile/0.1% trifluoroacetic acid (TFA). The pooled extracts were



dried, dissolved in a minimal volume of 6 M guanidine HCl-Tris, pH 8.2. Extracted peptides were then reduced with dithiothreitol (DTT) and alkylated with iodoacetamide. After the reaction, the mixture was diluted 4-X with water and the reduced guanidinium dodecylsulfate precipitate was removed  
5 by centrifugation. In-solution digestion was performed with trypsin in the diluted reaction mixture for complete digestion. Peptides were purified using a C18 microbore column (1 mm x 15 cm, VYDAC) using an Applied Biosystems 172 HPLC system. After injection of sample, the column was washed with 95 % solvent A (0.1 % TFA in water)/5 % solvent B (70  
10 acetonitrile/0/0.75 % TFA) for 10 min using a flow rate of 80 ul/min. Peptides were eluted with a gradient of 5 to 70 % Solvent B for 90 min. Purified peptides were sequenced using either an ABI 477 or 470A sequencer with on-line HPLC identification of phenylthiohydrazide (PTH) amino acids. Peptides containing either Cys 160 and Cys 119 were isolated as indicated  
15 below. The disulfides of the fully oxidized protein correspond to Cys 106-Cys 119 and Cys 60-Cys 160. To identify the disulfide bonds involved in the partly and fully reduced forms, trypsin peptide (s) unique to each were isolated. The partly and fully reduced forms both showed the peptide:  
-Leu<sub>149</sub>-Ser-Phe-Asn-Pro-Thr-Gln- Leu-Glu-Glu-Gln-Cys<sub>160</sub>-His-Ile<sub>162</sub>. The  
20 fully reduced protein showed in addition the following peptide:  
-Tyr<sub>102</sub>-Leu-Leu-Phe-Cys<sub>106</sub>-Met-Glu-Asn-  
Ser-Ala-Glu-Pro-Glu-Gln-Ser-Leu-Ala-Cys<sub>119</sub>-Gln-Cys<sub>121</sub>-Leu-Val-.

## Results

### Reduction of milk proteins by the thioredoxin system.

25 Thioredoxin was effective in the reduction of the most prevalent disulfide protein of milk, namely BLG. To this end, the redox state of pure BLG (A and B forms) was monitored following treatment with the NADP/thioredoxin system (NTS), consisting of NADPH, NTR and thioredoxin. As described above, samples were incubated with the NTS system and then analyzed using

the monobromobimane (mBBr)/SDS- polyacrylamide gel electrophoresis (PAGE) procedure. Here, as previously stated, the reduced (-SH) form of a target protein derivatized with mBBr and separated by SDS-PAGE, appears as a fluorescent band when viewed in ultraviolet light. It was found that, as  
5 seen in the previous Examples with a spectrum of proteins containing intramolecular disulfide bonds, the A and B forms of pure BLG were actively reduced by thioredoxin. When applied to milk, thioredoxin not only reduced BLG but also several other proteins, including  $\alpha$ -lactalbumin as determined by the SDS/PAGE/mBBr labeling procedure used for the gel in Fig. 7. For  
10 this gel 50  $\mu$ g of raw milk in PBS was applied to all lanes. The other variables for each lane are: Lane 1, control at 4°C, no addition; Lane 2, NTS, 55°C; Lane 3, NTS, 4°C; Lane 4, NGS, 55°C; Lane 5, NGS, 4°C; Lane 6, 5 mM DTT, 100°C, 5 min; Lane 7 was the same as Lane 6 except it was stained with Coomassie blue. Note that the excess of NADPH, NTR  
15 and thioredoxin maintained the target milk proteins in the reduced state throughout the experiment. The minor band traveling in front of the major (24 kDa) casein components ( $\alpha$  and  $\beta$ ) is  $\kappa$ -casein.

Temperature was found to have an interesting and useful effect on reduction. When treated at 55°C (45 min) as shown in Fig. 7, BLG was reduced but its  
20 mobility in the gel was only slightly changed. By contrast, when incubated at 4°C (17 h), the mobility of the bulk of BLG was decreased significantly and a new form of the protein appeared (Lane 3, Fig. 7). An assessment of the extent of reduction by gel scanning revealed that BLG is partly reduced at 55°C and fully reduced at 4°C. Comparison of the amino acid sequences of  
25 the tryptic peptides of the partly and fully reduced forms with the known structure of BLG gave further information on the nature of the reduction.

BLG is known to contain two disulfides, both intramolecular (See, Fig. 8): one clearly accessible on the surface and located close to the C-terminus (Cys

66-Cys 160) and the other, close to the core, located between two  $\beta$ -sheets (Cys 106-Cys 119 or possibly Cys 106-Cys 121). Sequence analysis confirmed that the exposed disulfide (Cys 66-Cys 160) was reduced at 55°C (partly reduced form) and that this as well as the hidden disulfide (Cys 5 106-Cys 119 or Cys 106-Cys 121) were both reduced at 4°C (fully reduced form) (see Materials and Methods). Differential reduction of BLG could also be achieved by altering pH. Reduction at pH 6.8 yielded only the partly reduced form whereas a mixture of the partly and fully reduced forms was observed at pH 8.0 (both at 37°C) (data not shown). The results confirm the 10 findings of others who showed that BLG was unstable at pH values above neutrality (Tanford, C. *et al.* (1959), "Transformation of  $\beta$ -Lactoglobulin at pH 7.5", *Biochemistry* 81:4032-4036) and undergoes conformational transitions at temperatures above 40°C (Qi, X.L. *et al.* (1997), "Effect of temperature on the secondary structure of beta-lactoglobulin at pH 6.7, as 15 determined by CD and IR spectroscopy: a test of the molten globule hypothesis", *Biochem. J.* 324:341-346). Significantly, the same reduction results for BLG as shown in Fig. 7 were obtained when only the components of the NADP/thioredoxin system were added to raw milk without buffer. Furthermore, the BLG in this treated milk without buffer remained reduced 20 for 2 days when stored at 4°C in air (data not shown).

As seen in Lanes 4 and 5 of the gel in Fig. 7, the monothiol glutathione maintained in the reduced state by NADPH and glutathione reductase also reduced BLG and to some extent other milk proteins, but less effectively than NTS. Other disulfide reductants, dithiotreitol and lipoic acid, were effective 25 but only when combined with thioredoxin as described in previous Examples for venom neurotoxins (data not shown).

### Digestion of milk proteins by pepsin and trypsin.

As shown in Examples 19 and 29, the trypsin sensitivity of small proteins containing intramolecular disulfide bonds (*e.g.*, trypsin inhibitors, venom neurotoxins) increases dramatically following reduction by thioredoxin.

- 5 Likewise, BLG was seen to follow this same pattern, that is, the thioredoxin-reduced BLG was highly sensitive to trypsin digestion whereas the oxidized BLG (*i.e.*, pure, untreated) was resistant (See, Example 34). Similar results were obtained in this Example with the pure BLG as well as with milk subjected to simulated gastric fluid as described by Astwood, J.D.
- 10 *et al.*, *supra*. When separated by SDS-PAGE and stained with Coomassie blue, BLG was found to be digested by pepsin but only after reduction by thioredoxin (See, Fig. 9). Fig. 9 shows the digestion of oxidized and reduced BLG as determined by mini SDS/PAGE and Coomassie blue dye. All incubations with SGF were at 37°C. 13.5  $\mu$ l of SGF mixture was applied
- 15 as indicated. The other variables for the gel in Fig. 9 were: Lane 1, BLG, SGF, 0 min; Lane 2, BLG, SGF, 60 min; Lane 3, reduced BLG by NTS at 55°C, SGF, 0 min; Lane 4, reduced BLG by NTS at 55°C, SGF, 60 min; Lane 5, reduced BLG at 4°C, SGF, 0 min; Lane 6, reduced BLG at 55°C, SGF, 60 min; Lane 7, SGF; Lane 8, BLG. As seen in Fig. 9, the difference
- 20 in sensitivity was striking.

- Oxidized BLG in milk was found to resist digestion for at least 60 min whereas the thioredoxin-reduced form was digested within 60 seconds (See, Figs. 10A-10C). Figs. 10A, 10B and 10C depict the effect of time on the digestion of milk buffered in PBS after thioredoxin reduction determined by
- 25 mini SDS/PAGE and Coomassie blue dye. Samples 2-10 contained 13.5  $\mu$ l of simulated gastric fluid mixture and samples 3-10 contained 50  $\mu$ g milk protein. After 0, 0.25, 1, 2, 4, 8, 15, 30, 60 min incubation the digestion was stopped by neutralization and aliquots were applied to the appropriate lanes of each of the gels in Figs. 10A, 10B and 10C. The other variables for

the gel in Fig. 10A were buffered milk control; for the gel in Fig. 10B: buffered milk, NTS, 55°C; for the gel in Fig. 10C: buffered milk, NTS, 4°C. The reduction of a single disulfide bond was sufficient. The partly (55°C) and fully (4°C) reduced forms of BLG showed no difference in pepsin sensitivity (compare Figs. 10B and 10C). Significantly, even though partially reduced by thioredoxin, the glutathione-treated sample was insensitive to digestion by simulated gastric fluid (data not shown). While the sensitivity of  $\alpha$ -lactalbumin and  $\kappa$ -casein were somewhat enhanced by thioredoxin reduction, BLG, either pure or in milk, was found to be the only protein not digested without reduction by thioredoxin (See, Figs. 7, 9 and 10A-10C).

#### **Allergy status of reduced and digested milk proteins.**

A comparison of the allergenicity of the major proteins of milk (casein,  $\alpha$ -lactalbumin, BSA, BLG) using dog model skin tests revealed that BLG is the major allergen of milk, accounting for 80% of the total wheal-inducing activity. Furthermore, when treated with the thioredoxin system, both milk and BLG showed a decreased ability to elicit an allergic response. Thus, similar to the results in the previous Examples testing allergenicity, the allergenicity of raw milk was decreased by a factor of 10 to 300, depending on the sensitivity of the dog tested (See, Figs. 11A and 11B). The graphs in Figs. 11A and 11B compare the thioredoxin-linked mitigation of skin test response to raw milk allergens in two dogs of differing sensitivity. The type I hypersensitivity reaction determined by the wheal area (mm<sup>2</sup>) was induced by 100  $\mu$ l intradermal injections of milk solution in PBS as indicated in Figs. 11A and 11B. The solutions were either pretreated with the NADP/thioredoxin system (NTS, 4°C) or untreated (control). The response of the two dogs is shown to illustrate that despite differences in sensitivity, the thioredoxin treatment mitigated allergenicity in both cases (*i.e.*, for mildly (Fig. 11B) and highly (Fig. 11A) sensitive animals). A PBS/glycerol

control and NTS were found to be negative for each dog. Also, tests with a number of dogs showing different sensitivity revealed no consistent difference in the allergenicity of the partly and fully reduced forms of BLG either pure or in milk (data not shown). The skin test data thus show that reduction by  
5   thioredoxin alters epitope accessibility such that the allergenicity of BLG and possibly other milk proteins is decreased.

Pure BLG showed an allergenic response similar to milk (compare oxidized and reduced zero time samples in Figs. 12A and 12B). In Figs. 12A and 12B, the oxidized "0" min and reduced "0" min bars represent the samples  
10   before digestion treatment, the oxidized 60 min and reduced 60 min bars represent samples treated with pepsin containing SGF. The type I hypersensitivity reaction observed by the skin wheal area ( $\text{mm}^2$ ) was induced by 100  $\mu\text{l}$  intradermal injections of digested neutralized BLG (Fig. 12A) or milk (Fig. 12B) either pretreated with the NADP/thioredoxin system (NTS)  
15   (reduced) or untreated (oxidized control). A neutralized SGF control was found to be negative for all tested dogs. Furthermore, skin tests showed no difference between the pure A and B forms of BLG with respect to the effect of thioredoxin on allergenicity (data not shown).

Skin tests carried out with both BLG and milk revealed that peptic digestion  
20   nearly completely eliminated the allergenicity of both preparations (See Figs. 12A and 12B). Based on skin tests, the allergy response is decreased by 300 to 1000 when digestion is coupled to reduction. Furthermore, the allergenicity of the digested samples was decreased to marginal levels, in both highly and mildly sensitive dogs.

25   These results are consistent with the conclusion that reduction by thioredoxin (1) alters the accessibility of the epitopes of intact proteins, so that allergenicity is decreased in most animals, and (2) renders stable allergens

such as BLG susceptible to pepsin digestion with the consequent almost complete loss of allergenic properties (see Figs. 13 and 14).

Finally, it was also found that  $\alpha$ -lactalbumin and BLG are digested by trypsin when reduced by the NTS system (See, Example 34). These results, coupled with the results obtained in this example with pepsin, confirm the ability of thioredoxin to render BLG protease sensitive.

#### **Feeding trials with atopic dogs.**

Upsets in the gastrointestinal tract leading to vomiting and diarrhea are symptoms that accompany the ingestion of food allergens but also the indigestibility of food proteins. Furthermore, the severity of these symptoms provides a measure of the strength of allergens that complements skin tests. To obtain evidence on the gastrointestinal response, the food of sensitive dogs was supplemented with BLG. As shown from the results set forth in Table VIII below, dogs consuming food containing 2.5 gm of reduced BLG (2.5 gm of BLG corresponds to 3/4 liter of milk) showed a significantly reduced extent of the vomiting. Repeated feeding trails indicated that, on average, about 70% of the gastric reflux disappeared. Significantly, this pattern was observed in a single set of dogs fed either treated or untreated BLG. A similar alleviation of the associated gastrointestinal upset response was observed by decreasing the BLG from the 2.5 gm used in Table VIII to 1.25 gm per can of food fed to each dog (data not shown). These observations complement the skin test results discussed above in showing that allergen recognition by the mucosal lymphatic tissue is mitigated by treatment with thioredoxin.

**Table XVIII**  
**Allergic Response of Dogs Alternately Fed**  
**Untreated or Thioredoxin-Treated  $\beta$ -Lactoglobulin**

5	Dog	Upper GI Index*			
		Exp. A		Exp. B	
		Control	Treated	Control	Treated
	6GCB3	7	--	--	0
	6GCB7	6	--	--	3
	6GCB1	--	0	5	--
10	6GCB4	--	6	9	--
	6GCB6	--	0	8	--

\* Measure of quantity and fluidity of induced vomit

Upper GI index response was measured using  
thioredoxin treated or untreated (control) BLG,  
2.5 gm, mixed with one can of dog food. Index  
calculations were performed after two separate  
feeding experiments, A and B. Upper GI index  
quotations were: no vomit=0, small vomit=1,  
large vomit=2, solid vomit=1, liquid  
vomit=2, liquid vomit with blood or bile=3,  
delayed vomit=1 and immediate vomit=2.

## Discussion

In this Example, thioredoxin was found to reduce specifically the  
intramolecular disulfide bonds present in BLG, a major milk allergen.  
Depending on conditions, thioredoxin, reduced by NADPH and NTR reduced  
either one or both of the disulfide bonds of BLG whether analyzed with pure  
protein or milk. The change in epitope distribution revealed experimentally  
by skin tests and feeding challenges was seen at the molecular level in  
structural models (Peitsch, M.C. (1996), *supra.*). When the exposed  
disulfide bond of BLG (Cys66-Cys160) was disrupted by site-directed  
mutagenesis using a computer model, the <sup>125</sup>Thr-<sup>135</sup>Lys epitope (Kaminogawa,  
S. *et al.* (1989), "Monoclonal antibodies as probes for monitoring the  
denaturation process of bovine  $\beta$ -lactoglobulin", *Biochemica et Biophysica*



*Acta* 998:50-56), which is nearby changed its position significantly whereas the epitope which is distant from both disulfides (<sup>8</sup>Lys-<sup>19</sup>Trp) did not (See, Figs. 13 and 14 ). Mutagenesis of the buried disulfide (Cys106-Cys119 or possibly Cys106-Cys121) led to no further change in the position of the  
5 epitope. A similar observation was made with a model prepared with human BLG epitopes (Ball, G. *et al.* (1994), "A major continuous allergenic epitope of bovine beta-lactoglobulin recognized by human IgE binding", *Clinical and Experimental Allergy* 24:758-764).

The present results suggest that reduction by thioredoxin lowers allergenicity  
10 of a target protein allergen in two ways. Reduction effects a change in protein structure that restricts epitope accessibility and enhances digestibility. In this way, the strength of the allergen is decreased and, in addition, should be more rapidly eliminated in the gastrointestinal tract.

#### EXAMPLE 40

##### 15                    **Digestion of Reduced Gliadins by Pepsin and Trypsin**

The protease susceptibility of the gliadin fraction isolated by ethanol extraction of wheat flour as described in Examples 9 and 10 was investigated. As shown in previous Example 10, gliadins contain intramolecular disulfide bonds that are specifically reduced by thioredoxin. Gliadins are also a major  
20 food allergen in wheat. Buchanan, B.B. *et al.* (1997), "Thioredoxin-linked mitigation of allergic responses to wheat", *Proc. Natl. Acad. Sci. USA* 94:5372-5377. The thioredoxin itself can be reduced either enzymatically by NADPH and NADP-thioredoxin reductase or chemically by dithiothreitol (DTT). For this Example, aliquots (50  $\mu$ l) of isolated gliadin (1.04 mg/ml)  
25 were incubated with or without thioredoxin (1.0  $\mu$ g ) which was reduced in the presence or absence of 1 mM DTT for an hour at room temperature at pH 7.5 in Tris-HCl buffer. At the end of the incubation, 5  $\mu$ l of trypsin from

Sigma (1 mg/ml) was added to each sample. The samples were subjected to digestion for another hour at 30°C. Digestion was terminated by PMSF (phenylmethylsulfonyl fluoride (2 µl of 100 mM) in each sample. After mixing with SDS (10 µl of 10%) and glycerol (15 µl of 50%), samples were  
5 analyzed by SDS-PAGE. After electrophoresis the 15% gel was stained with Coomassie blue and destained by methanol and acetic acid. Analysis of the protein bands demonstrated that the gliadins were stable to digestion in the oxidized (untreated) state but were degraded to lower molecular weight components following reduction by thioredoxin. Similar results were  
10 obtained when the gliadin aliquots were treated and digested with pepsin instead of trypsin as in Example 39.

This Example and Examples 34 and 39 show that a consequence of allergen reduction by thioredoxin is the striking increase in sensitivity to proteases. Thus, whereas the oxidized forms of gliadins and BLG were resistant to  
15 pepsin, the reduced forms were highly susceptible and were readily digested. The average concentration of innate thioredoxin in wheat flour is about 0.01% (Johnson, T.C. *et al.* (1987), "Reduction of purothionin by the wheat seed thioredoxin system and potential function as a secondary thiol messenger in redox control", *Plant Physiol.* 85:446-451). This is about 100 mg to  
20 about 200 mg per kilogram of flour or in bread, about 2 mg of thioredoxin per 100 gm of flour. However, the upper limit of naturally occurring thioredoxin in flour could be about 2 mg per gram of food protein. Bread, for example, that contained higher concentrations of thioredoxin treated with a thioredoxin reductant would be a less allergenic and more digestible bread.  
25 The increase in the sensitivity to pepsin by food proteins treated with thioredoxin and a thioredoxin reductant, seen also at the molecular level through modeling for BLG, would likely lead to a more rapid processing of ingested BLG in the gastrointestinal tract. If extended to the whole animal, treatment of milk and wheat and other food products with the thioredoxin

system would be expected to provide relief from the long-term effects of allergenicity and indigestibility -- notably edema and diarrhea.

Commercialization of the thioredoxin technology with respect to milk could be achieved by treating milk with the NTS before or after pasteurization (*i.e.*,  
5 at 55°C for 45 min or at 4°C for at least 10 hr). Such applications might include the following:

1. Applying the thioredoxin system in liquid form to raw milk or whey and coupling reduction to the pasteurization process.
- 10 2. Passing raw or pasteurized milk through a column of bound reduced thioredoxin. The thioredoxin could be reduced with NADPH and NTR prior to application of the milk. The thioredoxin could also possibly be reduced with dithiothreitol which could be removed from the column by washing prior to application of the milk
- 15 3. Storing NTS treated milk under non-oxidizing conditions to increase shelf life by using for example full or evacuated containers.
4. Adding thioredoxin and NTR to milk after pasteurization and adding NADPH in solid form just prior to use.
- 20 5. After treatment with thioredoxin, subjecting milk to limited proteolytic digestion with enzymes such as trypsin. Such product could be used to induce tolerance in milk-sensitive individuals.

#### EXAMPLE 41

##### **Determination of the Decreased Allergenicity and Increased Digestibility of Thioredoxin Treated Ragweed Allergen Proteins**

**Reduction of pollen proteins by the thioredoxin system.** Giant ragweed allergen extract, purchased from Bayer Inc., was analyzed for protein by the Bradford assay using bovine  $\gamma$ -globulin as standard (Wong, J.H. *et al.* (1995) "Thioredoxin and seed proteins" *Methods Enzymol* 252:228-240). The protein, 100  $\mu$ g, was reduced with 1.25 mM NADPH, 1.7  $\mu$ g of *E. coli* NADP/thioredoxin reductase (NTR) and 1.7  $\mu$ g of *E. coli* thioredoxin (Trx) by incubation for 45 min at 37°C in 30 mM physiological buffered saline (PBS) (10 mM  $\text{Na}_2\text{HPO}_4$ , 1.8 mM  $\text{KH}_2\text{PO}_4$  2.7 mM KCl and 137 mM NaCl, pH 7.4) resulting in a final volume of 100  $\mu$ l. The extent of reduction was determined by the SDS-PAGE/monobromobimane procedure as previously described above and also as described by Wong *et al.* (*Ibid.*). Gels were visualized under U.V.

**Skin tests.** Procedures to measure the type I hypersensitivity reaction by skin tests in allergic doses were substantially the same to those previously described above (see, Examples 36, 37 and 39). In brief, Evans blue dye 0.5% (0.2 ml/kg) was injected intravenously 5 minutes prior to skin testing. Aliquots of 0.1 ml of pollen allergen extract were injected intradermally on ventral abdominal skin in half-log dilutions. Skin tests were read blindly by the same experienced blinded observer scoring two perpendicular diameters for each blue spot. Appropriate negative controls (diluted in PBS) were included for each animal tested.

**Digestion of pollen proteins by simulated gastric fluid.** Pollen protein, 650  $\mu$ g, or pure Amb t V, 100  $\mu$ g, was incubated in 100  $\mu$ l PBS buffer at 37°C for 45 min, with or without a mixture of 2.4  $\mu$ g Trx, 4.2  $\mu$ g NTR and

- 2.5 mM NADPH. Then, 50 µl of each reaction mixture, containing 325 µg pollen protein or 50 µg Amb t V, was digested in 100 µl simulated gastric fluid (SGF) as described by Astwood et al. (Astwood, J.D. *et al.* (1996), "Stability of food allergens to digestion in vitro" *Nature Biotechnology* 14:1269-1273. SGF is composed of 0.32% porcine pepsin (w/v) and 30 mM NaCl adjusted to pH 1.2 with HCl. The digestion mixtures were incubated at 37°C and stopped at 0, 0.25, 1, 2, 4, 8, 15 or 60 min with a 0.375 volume of 160 mM Na<sub>2</sub>CO<sub>3</sub> to give a neutral pH. Proteins were analyzed by SDS-PAGE and, as indicated, by skin tests.
- 10 **Amb t V purification.** Purification was achieved by adaptation of the method of Roebber et al. 1985 (Roebber, M. *et al.* (1985), "Immunochemical and genetic studies of Amb.t. V (Ra5G), an Ra5 homologue from giant ragweed pollen" *J. Immunol.* 134:3062-9). In brief, 100 g of non-defatted pollen (Greer Laboratories) was suspended in 1 liter of
- 15 buffer [50 mM Tris-HCl pH 7.4 containing 1 µM phenylmethylsulfonyl fluoride (PMSF) and 1 mM EDTA-Na] and stirred gently for 30 min at room temperature. The mixture was then centrifuged for 15 min at 3,840 xg, 4°C. The collected supernatant fraction was filtered once through glass wool and then twice through Whatman quantitative filters. The pellet containing the
- 20 pollen grains was not further used. The high quantity of lipids in the filtered supernatant fraction was removed by extraction with an equal volume of petroleum ether and centrifuged at 11,300 xg for 10 min, 4°C. The petroleum ether step was repeated 4 times. The resulting clarified solution was concentrated by passage through an Amicon YM-3 membrane and
- 25 separated on a Sephadex G-50F gel filtration column (2.1 x 90 cm) equilibrated and eluted with 20 mM Tris-HCl pH 7.5 supplemented with 200 mM NaCl. The fractions containing 5 kDa proteins were analyzed by 15% SDS-PAGE, combined and concentrated again with a YM-3 membrane. Ammonium sulfate was added to the concentrated proteins to a final

concentration of 2.6 M. The protein mixture then was fractionated on a 1 ml HiTrap Phenyl Sepharose column (Pharmacia) equilibrated with 200 mM phosphate buffer pH 7.0 and eluted with a 50 ml decreasing gradient ranging from 2.5 to 0 M ammonium sulfate in this same buffer. The pure Amb t V  
5 was recovered in a single peak at approximately 0.8 M. Finally, the pure protein was dialyzed against 5 mM potassium phosphate buffer, pH 7.0 and stored at -70°C for further experiments. Protein content was quantified using a molar extinction coefficient of 5800.

**Data analysis.** The statistical significance of the skin test results showing the  
10 thioedoxin-linked mitigation of pollen allergens was determined by paired one-tailed, *t*-tests. The null hypothesis, which assumes no difference in wheal area induced by untreated vs. thioedoxin-treated pollen proteins, was tested against the alternative hypothesis that the treatment resulted in mitigation of allergic response. The *t*-tests were completed for each dilution  
15 series (0.07 to 219 ng allergen) at 0.05 level of significance on all sensitive dogs (df=8).

## Results and Discussion

**Reduction of the pollen proteins.** As shown in Figure 15, several proteins were actively reduced by thioedoxin, namely Amb t V at 5 kDa, and  
20 unidentified proteins at 12, 14 and 35 kDa. Furthermore, the extent of reduction varied with the reaction temperature. As shown in Figure 15, reduction was most efficient at 37°C or higher. In the case of Amb t V, the results indicate that multiple disulfide bonds (possibly all 4) are reduced at 37°C and 55°C, whereas a lower number is reduced at 4°C. In parallel, it  
25 was also observed that reduction by thioedoxin strikingly affected the high thermostability described for Amb t V (Baer, H. *et al.* (1980), "The heat stability of short ragweed pollen extract and the importance of individual allergens in skin reactivity" *J. Allergy Clin. Immunol.* 66:281-5).

Precipitation of protein was found to occur at 55°C after reduction by thioredoxin (data not shown).

**Mitigation of the allergenicity by thioredoxin reduction.** As shown in Table XIX below, reduction by thioredoxin effectively mitigated the skin test response. While less pronounced than found with milk (see, Example 39), the thioredoxin-linked decrease in the allergenicity of pollen crude extract ranged between 3- and 33-fold. Furthermore, based on t test analysis, the mitigation was statistically significant.

**Table XIX**

10 Statistical evaluation of the mitigation by thioredoxin of the skin test response to pollen allergy. Paired one tail t test two samples for mean where P value is equal to or less than 0.05.

	<u>Allergen, ng</u>	<u>Observation, n</u>	<u>t Critical</u>	<u>t test value</u>	<u>P value</u>	<u>Mitigation</u>
15	219	20	1.729	0.974	0.1711	Not Significant
	66	27	1.706	2.326	0.0140	Significant
	22	27	1.706	3.045	0.026	Significant
	6.6	27	1.706	3.253	0.0016	Significant
	2.2	27	1.706	2.185	0.0191	Significant
	0.66	27	1.706	2.805	0.0047	Significant
20	0.22	27	1.706	1.545	0.672	Not Significant
	0.07	10	1.833	1.471	0.0877	Not Significant

**Effect of thioredoxin reduction on the sensitivity of pollen proteins to pepsin.** To determine whether reduction by thioredoxin increases the

digestibility of pollen allergens, the oxidized and reduced forms of purified Amb t V protein were treated with SGF (Astwood, J.D. *et al.* (1996), "Stability of food allergens to digestion in vitro" *Nature Biotechnology* 14:1269-1273. 12). First, it was found that, like beta-lactoglobulin, Amb t V .  
5 was digested only when reduced by the thioredoxin system. The oxidized protein was resistant to pepsin for up to 60 min, consistent with its allergenic potential as shown in Figure 16A (Astwood, J.D. (1996), *supra*). By contrast, the thioredoxin-reduced protein almost completely disappeared after 2 minutes as shown in Figure 16B. This sensitivity to pepsin was  
10 substantiated by skin test data obtained with crude pollen extracts tested with 8 dogs. It was observed that the skin test reaction elicited by the oxidized pollen extract was retained after digestion in 4 of these dogs (typified by the dog designated 6CGB1, see Example 39, "Food sensitization of atopic dogs"), indicating that the allergens were pepsin resistant as shown in Figure  
15 17A. By contrast, when the preparation was reduced by thioredoxin, the allergic response declined markedly in these dogs. This confirmed that allergens were disulfide proteins and consistent with the gel data that the proteins had then been digested. The 4 other dogs (typified by the dog designated 5CBB3 in Figure 17B) showed a pronounced decrease in reaction  
20 when the oxidized (untreated) proteins were digested by SGF -- i.e., the allergens most active were digested from the outset. This finding indicated that the second group is less sensitive to disulfide proteins.

The data in Figure 18 provide evidence that this differential response to untreated and thioredoxin-treated preparations resides in the sensitivity of a  
25 particular animal to disulfide protein allergens. As seen for pure Amb t V in Figures 17A and 17B, the disulfide proteins in crude preparations (including Amb t V) were strongly resistant to pepsin unless reduced by the NADP-thioredoxin system (Figures 18A and 18B). Furthermore, the reduction temperature is important; digestion increased progressively as the reduction



temperature increased from 4°C to 37°C to 55°C. Without reduction, the proteins in the 5 to 20 kDa range were not digested even after 60 incubation with SGF.

- The above results are in accord with the conclusion that, in contrast to milk in which the major allergen-- $\beta$ -lactoglobulin--is a disulfide protein (del Val, G. *et al.* (1999), "Thioredoxin Treatment Increases Digestibility and Lowers Allergenicity of Milk" *J. of Allergy Clin. Immunol.* In press), ragweed pollen contains a complex allergen mixture consisting of proteins both with and without disulfide bonds.
- 10 These results show that the NADP/thioredoxin system alleviates the allergic response to a pollen. It also shows that the importance of active disulfide proteins on complex mixtures of allergens can be assessed by comparing the pepsin sensitivity of oxidized and thioredoxin-reduced samples. The more important that disulfide proteins are to the allergic response, the more
- 15 effective thioredoxin is in the alleviation of that response.

## EXAMPLE 42

### Use of Thioredoxin Treated Ragweed Pollen Allergen for Immunotherapy

- An animal that exhibits sneezing and coughing (bronchospasms) upon inhaling a specific amount of an aerosol of ragweed pollen protein Amb t V is the subject of this investigation. The Amb t V protein in the aerosol has not been treated with any reducing agents. The animal is subcutaneously injected with increasing doses of a clinical solution containing
- 20 67micrograms/cc of Amb t V which has been purified and incubated with
- 25 thioredoxin, NADPH and NTR as described in Example 41. These subcutaneous injections are to determine the closest dose or end point at

which a local allergic reaction to the treated Amb t V occurs. A wheal of about 5mm is observed to occur at the injection site with 10 nanograms (ng) in 0.15cc. With 10 ng established as the end point, the animal is then injected subcutaneously three days later with an amount of Amb t V one log  
5 below the end point or, in this case, 1 ng. This dosage corresponds to the first dose in the injection schedule set forth in the table below.

Table XX

	<u>Dilution IV (blue)</u>	1:10,000	<u>Dilution II (yellow)</u>	1:100
	First dose - 0.15cc		11th dose - 0.15cc	
10	2nd dose - 0.30cc		12th dose - 0.25cc	
	3rd dose - 0.60cc		13th dose - 0.35cc	
	4th dose - 1.00cc		14th dose - 0.50cc	
			15th dose - 0.75cc	
	<u>Dilution III (green)</u>	1:1,000	16th dose - 1.00cc	
15	5th dose - 0.15cc		<u>Dilution I (red) conc</u>	1:10
	6th dose - 0.25cc		17th dose - 0.10cc + 0.10cc saline	
	7th dose - 0.35cc		18th dose - 0.15cc + 0.15cc saline	
	8th dose - 0.50cc		19th dose - 0.20cc + 0.20cc saline	
	9th dose - 0.75cc		20th dose - 0.25cc + 0.25cc saline	
20	10th dose - 1.00cc		21st dose - 0.30cc + 0.30cc saline	

The first dose therefore is a 0.15cc injection of a 1:10,000 dilution of the Amb t V 67micrograms/cc solution. The subject is observed for a local reaction. Since no reaction is observed upon injection of the first dose, the  
25 animal is given the second injection of 0.30cc of the 1:10,000 dilution or 2ng. After three days, the third dose of 0.60cc of the 1:10,000 dilution is subcutaneously administered. The injection schedule is then followed with injections being given every three days until the top dose tolerated by the subject is reached. The top dose is indicated by a large local skin reaction,  
30 *i.e.*, a wheal larger than the size of a silver dollar, and/or systemic symptoms such as urticaria (hives), sneezing, vomiting or a fall in blood pressure. The top dose with this animal is the 21st dose but with another subject it could

have been the 11th, the 15th, or the 18th, *etc.* Upon observing the top dose, the dosage is decreased one or two doses and this reduced dosage is held as the new top dose. The animal is subsequently injected with the new top dose every three to four weeks. Approximately six months after receiving the new  
5 top dose at three to four week intervals, the subject animal inhales the same amount of the aerosol of Amb t V that previously caused sneezing. No, or limited, sneezing or coughing is observed as a result of this inhalation. The animal continues to receive the new top dose at the regular intervals and remains free or comparatively free of sneezing and any other allergic  
10 reactions upon further inhalation of non-reduced Amb t V.

#### **EXAMPLE 43**

##### **Comparison Between Thioredoxin-Treated Pollen Protein and Untreated Pollen Protein for Immunotherapy Effectiveness**

A group of 10 animals, all of the same species, that exhibited sneezing and  
15 other allergic reactions upon inhalation of a specific amount of non-reduced, allergen Amb t V are the subjects of the investigation. To determine an allergic end point, the animals are divided into two groups of five each. The animals are tested for antibody levels and Group A is then injected with increasing doses of non-reduced Amb t V, and Group B is injected with  
20 increasing doses of thioredoxin, NADPH and NTR incubated Amb t V as described in Example 42. The doses based on mg/kg of body weight for all 10 animals is the same even though the absolute dosage given each animal may differ. The end point dose for each animal in the group is determined. The average end point dose based on mg/kg of body weight for Group A is  
25 lower than for Group B. The animals are then injected according to the method and injection schedule set forth in Example 42. The top dose for each animal is determined. The top doses for the animals in Group B are consistently higher with lesser local or systemic symptoms being observed

- than with the top doses for the animals in Group A. Then in the manner described in Example 42, the animals are assigned a new top dose and are subsequently injected every three to four weeks with this new top dose as in Example 42. Following approximately 6 months of such injections, the
- 5 animals of Group A show a somewhat diminished allergic response upon subsequent inhalation of the specific amount of non-reduced allergen, while animals of Group B exhibit very limited or no significant allergic response. Furthermore, animals of Group B are able to tolerate much higher doses of the allergen, in some cases up to 10 to 100 times. A test to determine the
- 10 antibody levels in the treated animals shows that the IgG antibody titer in the Group B animals is higher following immunotherapy while the titer of IgE is lower over time. The IgG antibody titer is also somewhat higher in Group A following immunotherapy, but much lower on the average than the IgG titer of the Group B animals.
- 15 These results show that
- Thioredoxin reduction of ragweed pollen mitigates the allergic response.
  - Thioredoxin reduction markedly enhances the digestibility of disulfide pollen allergens.
  - 20 · Thioredoxin (and potentially other disulfide reductants such as lipoic acid) is useful in a hypoallergenic pollen extract for immunotherapy of pollen allergic patients by enhancing the production of IgG instead of IgE antibodies and improving the safety of the therapy.
- 25 Embodiments of the invention could include treating pollen allergies with eye drops or nose spray containing:
- lipoic acid,

- the NADP/thioredoxin system,
- lipoic acid and the NADP/thioredoxin system.

This invention can be used to treat and prevent many types of allergies (*e.g.*, food, pollen, dust mite) due to disulfide proteins by pretreating these proteins  
5 with the NADP/thioredoxin or lipoic acid as indicated above.

### Stabilization of Reduced Proteins

Two sulfhydryl (SH) groups can often spontaneously react to form an S-S bond in the presence of oxygen. This property suggested that the effects of thioredoxin (Trx) (*i.e.* the NADP-thioredoxin system (NTS)) and other  
10 reductants, such as dithiothreitol (DTT), to decrease allergenicity and increase digestibility by pepsin may not be permanent. For example, the reduced form of  $\beta$ -lactoglobulin (BLG), a major allergen of cow's milk, is fully reoxidized after 3 days in air whether reduced by either NTS or DTT. This invention provides for the stabilization of proteins, including allergenic  
15 proteins reduced by the NTS at 4°C and by DTT at 55°C. Other physiological reductants such as lipoic acid may also be used to initially reduce the proteins. In general, the reduction reactions were conducted as described in previous examples. The proteins are stabilized in that the SH groups of these proteins and allergens are blocked and prevented from  
20 reoxidizing to form S-S bonds. This stabilization is maintained for at least three months, preferably at least six months and more preferably for longer than one year. Reduced BLG was stabilized by reaction with the physiological disulfides, cystamine and oxidized glutathione (GSSG) and also by heat. Other physiological disulfides such as homoglutathione and some  
25 microbial disulfides would also be suitable. Besides BLG, examples are provided for milk, peanut and pollen extracts. However, other reduced proteins such as those of egg, soy, rice, wheat and barley may also be

stabilized by reaction with the same physiological disulfides and by treatment with heat. The allergenic and digestibility properties of such derivatives is described below.

#### EXAMPLE 44

##### 5                      Stabilization of Reduced BLG and Milk

###### *Experimental Procedures*

*Chemicals and Proteins.* Bovine  $\beta$ -lactoglobulin (BLG) and porcine pepsin were obtained from Sigma. Raw cow's milk was from the University of California Davis. Thioredoxin and NADP-thioredoxin reductase (NTR) were  
10 purified from *Escherichia coli* strains overexpressing the proteins (Jiao et al., 1992). Monobromobimane (mBBBr, trade name Thiolyte MB) was purchased from Calbiochem. Cystamine, glutathione oxidized form, NADPH, and DTT were from Sigma. Other chemicals and reagents were from sources previously identified in Example 39 and as described in Buchanan *et al.*,  
15 1997, herein incorporated by reference.

*Reduction and Blocking of Protein Disulfide Bond.* BLG was reduced by 2.5 mM DTT in phosphate buffered saline (PBS) for 1 hr at 55°C or by the NADP/thioredoxin system (NTS) for 17 hr at 4°C as described in Example 39 and in del Val *et al.*, 1999, herein incorporated by reference. Following  
20 gel filtration with a Sephadex G-25 column (Amersham Pharmacia) to remove reductants, the reduced protein (less than 0.25 mg/ml) was incubated either without addition or with 10 mM oxidized glutathione (GSSG) or 10 mM cystamine for 3 days on ice. The mixtures were then subjected to a second gel filtration on a Sephadex G-25 column to remove excess GSSG and  
25 cystamine. If necessary, the resultant preparations were concentrated by ultrafiltration with Centricon YM-3 filter (Millipore). Heat treatment of NTS-reduced BLG was carried out for 1 hour at 55°C after reduction at 4°C overnight. Raw cow's milk (47 mg protein/ml) was reduced with 50 mM

DTT and blocked with cystamine and GSSG as described above for BLG. Heated milk was prepared by incubation for 1 hour at 55°C and gel filtration with A Sephadex G-25 column.

*Monobromobimane Labeling and Analysis of Proteins.* Ten µg of BLG or 50  
5 µg milk protein was incubated with 4 mM mBBBr for 20 min as described in previous Examples and also as described in Wong *et al.*, 1995, herein incorporated by reference, and then subjected to non-reducing SDS-PAGE (10 to 20% gradient gel) to separate the native, partly reduced and fully reduced forms. The labeled proteins were stored at -80°C, if not  
10 immediately loaded on the gel. After rinse with two changes first in 12% acetic acid and then in 20% methanol/10% acetic acid to fix protein and remove excess mBBBr, the fluorescent image was taken with a Gel Doc (BioRad). The gel was then stained with Coomassie brilliant blue R-250. After destaining, the image was recorded as described above. The intensity  
15 of fluorescence and staining of each band were quantified by Multi-Analyst ver. 1.0.2 (BioRad).

*Skin Tests.* The background of the dog colony, immunization schedule and procedure for skin testing have been previously described in Examples 36-39 and also in del Val *et al.*, 1999, Buchanan *et al.*, 1997, Ermel *et al.*, 1997,  
20 Frick and Brooks, 1983, herein incorporated by reference. Here, 3 litters of the 7th generation designated 7FA, 7FB and 7FC were immunized with commercial preparations (Bayer) of cow's milk, soybean and other food extracts (wheat or barley and walnut, Brazil nut and peanut) and also ragweed pollen. BLG samples for testing were prepared in half-log dilutions with  
25 PBS; 100 µl of each dilution was injected intradermally on the dog's abdomen.

*Estimation of Relative Allergenicity.* Wheal area of each spot was plotted by the Lineweaver-Burk double-reciprocal plot to obtain the maximum wheal area. The log of the dose of BLG that effects a half-maximum of wheal area with the control sample, termed Effective Dose 50 (ED50), was calculated and used for statistical analysis. For simplification, ratio of the reciprocal of ED50 to that of the control sample is presented below as relative allergenicity.

*Assay for Stability to Pepsin.* BLG (1 mg/ml) or milk (15 mg/ml) were mixed with simulated gastric fluid (SGF) to give 0.32% pepsin and incubated for the indicated times at 37°C as described in Example 39 and Astwood *et al.*, 1996, herein incorporated by reference. The digested proteins were analyzed by reducing SDS-PAGE (10 to 20 % gradient gel) also as described in Laemmli, 1970, herein incorporated by reference.

*Quantification of Protein.* BLG was quantified by absorbance at 280 nm using an extinction coefficient of  $16740 \text{ M}^{-1}\text{cm}^{-1}$ . Milk protein was estimated according to Bradford, M. M. "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding." *Anal. Biochem.* 72:248-254 (1976)., herein incorporated by reference, using bovine  $\gamma$ -globulin as a standard.

*Peptide Mapping.* One hundred  $\mu\text{g}$  of the BLG sample was labeled with mBBR as described above. The labeled protein was precipitated with 12% trichloroacetic acid to remove excess mBBR. The protein precipitate was rinsed three times with acetone. The precipitate was dissolved in 100  $\mu\text{l}$  of 0.1 M  $\text{NH}_4\text{HCO}_3$  after drying by Vacufuge (Eppendorf). The labeled BLG was digested with 20  $\mu\text{g}/\text{ml}$  trypsin for 30 min at 37°C. The reaction was stopped by addition of 1  $\mu\text{l}$  of trifluoroacetic acid (TFA). The resultant solution was diluted 10 times with 0.1M  $\text{NH}_4\text{HCO}_3$  and applied to Sephasil



Peptide C18 column (Amersham Pharmacia) using the Biocad perfusion chromatography system (PerSeptive Biosystems). The tryptic peptides were eluted with a 0-50% gradient of acetonitrile containing 0.1% TFA. The separation was monitored by absorbance at 215nm and fluorescence at 480nm  
5 using excitation light of 374nm. The molecular mass of the fluorescent peptides was determined by mass spectroscopy.

### **Results**

*Cystamine- and Oxidized Glutathione-Derivatized  $\beta$ -Lactoglobulin.* As shown in Example 39, DTT reduced BLG, either pure or in cow's milk, was  
10 characterized by decreased allergenicity and increased susceptibility to digestion by pepsin. However, the reduced form of BLG was not stable and was fully reoxidized when left in air for 3 days. In an attempt to prevent the reoxidation and preserve the hypoallergenic and hyperdigestible form of the protein, the SH groups of reduced BLG were blocked with the physiological  
15 disulfides, cystamine and GSSG.

BLG was first reduced by 2.5 mM DTT rather than the NTS to simplify the reaction. Reduction by DTT was facilitated by incubation at the increased temperature of 55°C. After removing the DTT, the reduced BLG (less than 0.25 mg/ml) was mixed with cystamine or GSSG. Then the proteins were  
20 labeled with the fluorescent sulfhydryl probe, Monobromobimane (mBBBr) and the products were subjected to non-reducing SDS-PAGE. The native form of BLG migrated the fastest (FIGS. 19A and 19B, Lane 4) and the fully reduced form the slowest (FIGS. 19A and 19B, Lane 1). [In FIGS. 19A and 19B, please refer first to the right (protein) panel (FIG. 19B) and then to the  
25 left (fluorescence) panel (FIG. 19A). The full legends for Figs. 19 through 27 are located at the end of Example 44 following the tables.] The partly reduced form was in between (FIGS. 19A and 19B, Lanes 2, 3 and 8). As found previously, the mobility of BLG in non-reducing gels depended on the

number of intramolecular S-S bonds in the protein. The fluorescence intensity reflected the number of SH groups. After DTT was removed, there was an increase in the partly reduced form that was accompanied by formation of a new species, probably a dimer of BLG, with a molecular mass of about 36 kDa (FIGS. 19A and 19B, Lane 3 compared to Lane 2). This finding suggested that some reoxidation of the protein occurred during gel filtration.

The experiment using various concentrations (0 to 10 mM) of cystamine showed an increase in the slow-migrating protein band and decrease in the fluorescence. As shown in FIGS. 19A and 19B, (Lane 5 in 19A) the fluorescence and protein corresponding to native BLG (Lane 5 in 19B) were negligible with 10 mM cystamine, indicating that essentially all of the SH groups had reacted with cystamine to form a cysteamine-derivatized form. The proposed identity of the three different forms recovered in cystamine-treated BLG is summarized below (FIGS. 19A and 19B, Lane 5).

- A feebly fluorescent major migrating like the partly reduced form ( $52.2\% \pm 5.3\%$  of the total protein): BLG with 1 intramolecular S-S bond and 3 cysteines bound to cysteamine.
- A non-fluorescent minor band ( $19.5 \pm 1.3\%$ ) migrating like the fully reduced form: BLG with 5 cysteines bound to cysteamine.
- A non-fluorescent diffuse band ( $28.3 \pm 5.4\%$ ) migrating around 36 kDa: BLG dimer.

GSSG also bound to BLG in a concentration dependent manner. However, a 10-fold greater concentration was required relative to cystamine, indicating a

less efficient reaction. The different forms observed in the GSSG-treated BLG sample are summarized below (FIGS.19A and 19B, Lane 6).

- A weakly fluorescent band corresponding to the native form ( $0.47 \pm 0.09$  moles SH/mole BLG) ( $35.2 \pm 4.2\%$  of total protein): BLG with 2 intramolecular S-S bonds formed during reoxidization. In addition, half of the BLG contained one SH per molecule and half contained one bound glutathione per molecule.
  - A weakly fluorescent main band between the fully and partly reduced forms ( $48.8 \pm 4.8\%$ ): BLG with 1 intramolecular S-S bond formed during reoxidization, 2 cysteines bound to glutathione and 1 cysteine as SH.
  - A non-fluorescent diffuse band ( $16.0 \pm 6.9\%$ ) migrating around 36 kDa: BLG dimer.
- In sum, incubation of DTT-reduced BLG with cystamine or GSSG resulted in S-S bond exchange to yield thiolated BLG that existed in several different forms.

*Allergenicity of BLG Derivatives with DTT as Reductant.* The allergenicity of the cystamine- and GSSG-treated BLG derivatives as well as the reduced and reoxidized forms was determined by skin tests with dogs sensitized to cow's milk. Results with a moderately sensitive dog are shown in FIG. 20 for a DTT experiment. Ten-times more of the reduced and GSSG-treated forms and 5-times more of cystamine-treated form were required to initiate an allergic response relative to the control. Interestingly, the reoxidized form of BLG showed essentially the same response as the reduced form and the

- derivatives. Similar results were obtained by including all of the 7 dogs tested in the analysis (Table XXI). The dose of BLG required to induce a half maximum of wheel area, ED<sub>50</sub>, indicated that, respectively, 10- and 20-times more cystamine (CYM)- and GSSG treated BLG were required to induce the same allergic response as native BLG, whereas 7-times more DTT-reduced BLG was required. These results indicated that the thiolated BLG derivatives showed lower allergenicity than the DTT-reduced sample. Furthermore, the reoxidized (REOXD) form was 20-times less allergenic than the control.
- 10 In an attempt to determine the basis for its lower allergenicity, the reoxidized sample was subjected to peptide analysis. There was no difference in the tryptic peptide map between the reoxidized and native forms of BLG. This finding suggests the absence of recombination of sulfhydryl groups to form unusual S-S bonds during reoxidation, other than the 12.2% of the reoxidized
- 15 BLG that formed an apparent dimer via intermolecular S-S bridge formation (FIGS. 19A and 19B, Lane 7). While the basis for lower allergenicity of the reoxidized form remains to be determined, the change could be attributed to an irreversible alteration of the structure of BLG during reduction at 55°C and subsequent oxidation.
- 20 *Allergenicity of BLG Derivatives Using NTS as Reductant.* Following reduction by NTS, BLG was found to react with cystamine and GSSG to the same extent as the DTT-reduced counterpart (FIGS. 21A and 21B). However, the extent of reduction of allergenicity was less than seen with DTT (Table XXII). Thus, the allergenicity of both the cystamine- and
- 25 GSSG-BLG derivatives was about 2.5-times less than that of native BLG. A similar decrease in allergenicity was obtained with the protein reoxidized following NTS-reduction. The NTS-reduction was performed at 4°C rather than 55°C. For reasons that are not yet clear, the litters of dogs used in the

study showed less mitigation of BLG allergenicity on reduction by NTS than the parent litters used previously in Example 39. Thus, while some of the dogs showed a lower skin test response to NTS-reduced BLG, this was not observed with the majority of dogs tested, unlike results with the previous  
5 litter in Example 39. Taking this finding into account, the skin test results in Table XXII indicate that, as found with DTT (cf. Table XXI), thiolation following reduction is more effective than reduction in mitigating allergenicity.

*Effect of Heat.* The basis here for the discrepancy in allergenicity between  
10 NTS- and DTT-reduced BLG is unclear. It is noted that there are differences between the DTT and NTS treatments. Thus, reduction by DTT was carried out without Trx at 55°C, while that by NTS is with Trx at 4°C. To compare these differences, BLG reduction experiments were performed using DTT plus Trx at both 25°C and 55°C for 1 hour. Reduction at 25°C  
15 failed to mitigate allergenicity consistently (again unlike earlier results), but reduction at 55°C effected an 8-fold decrease in allergenicity possibly due to the reasons discussed above (Table XXIII). Significantly, BLG was fully reduced at both temperatures (FIGS. 22A and 22B, Lane 2 vs. 4 and 5 vs. 7, both panels). Moreover, no effect on the allergenicity of BLG at 55°C  
20 without any other treatment (55°C, Control), indicated that reduction of the protein was essential. Similar effects were seen with NTS in which the sample heated for 1 hour at 55°C after full reduction of BLG by NTS was about 5-fold less allergenic than the control (Table XXIV). The unheated NTS-reduced sample showed much less of an effect. For comparison, the  
25 sample reduced with DTT at 55°C showed a corresponding improvement of about 5-times. Interestingly, the effect of heat did not apply to cystamine-treated BLG (Table XXIV). Thus, in this example, the allergenicity of BLG reduced by NTS at 4°C showed similar mitigation (3 times) with and without heat before cystamine treatment. The results indicate

that heat treatment improves the mitigation of allergenicity of BLG reduced either by DTT or NTS, but not of the NTS reduced cystamine derivative.

*Digestibility Studies with of BLG.* The test for digestibility developed by Astwood *et al.* (1996) using simulated gastric fluid (SGF) was used. As  
5 found previously in Example 39 and also in Astwood *et al.*, 1996; del Val *et al.*, 1999, native BLG was stable to digestion by SGF (FIG. 23A, Lane 3) with a half-life of about 2 hours (FIG. 24). On the other hand, the DTT-reduced and cystamine-treated BLG were both completely digested with pepsin in 2 minutes (FIG. 23A, Lanes 5 and 7). The digestion of  
10 GSSG-treated BLG occurred in two phases, a rapid phase resembling the DTT-reduced and cystamine-treated samples and a much slower phase resembling the BLG control (FIG. 24). The indigestible component constituted 17% of the total and migrated like native BLG on non-reducing SDS-PAGE (FIG. 23B, Lane 4), suggesting that the indigestible component  
15 was the reoxidized form. About 80% of reoxidized BLG was stable in SGF for 2 minutes (FIG. 23A, Lane 11) and, like the control, showed a half-life of about 2 hours in SGF (FIG. 24). The digestibility of the BLG derivatives using Trx as reductant was as with DTT (FIG. 23C and FIG. 24). Incubation with SGF without pepsin for 1 hour at 37°C had no effect on all BLG  
20 samples. The results with the reoxidized form indicate that enhanced digestibility does not always correlate with decreased allergenicity -- a factor of considerable interest in assessing the allergenicity of unknown proteins.

*Redox Studies with Cow's Milk.* The DTT experiments presented above for BLG were repeated using cow's milk. In these experiments, milk was treated  
25 with cystamine and GSSG following reduction by the procedure described above for BLG except that the DTT concentration was 50 mM. With milk, aggregates were observed, which did not enter the gel in non-reducing SDS-PAGE with the cystamine- and GSSG-treated and reoxidized samples

(FIGS. 25A and 25B, Lanes 4, 5 and 6). Unlike observations made with pure BLG, there was no obvious protein band migrating at the location corresponding to BLG, especially in the GSSG-treated milk sample (FIGS. 25A and 25B, Lanes 4, 5 and 6 compared to FIGS. 19A and 19B, Lanes 5, 6 and 7). By contrast, profiles of the derivatized milk samples on reducing (by 2-mercaptoethanol) SDS-PAGE were the same as with raw milk, indicating that BLG was not degraded and was not removed by gel-filtration (FIG. 26, Lane 1 in panels A to E). The aggregates are, therefore, likely due to intermolecular S-S bridges that formed between BLG itself and possibly other milk proteins. The aggregates were fluorescent, indicating that they contained SH groups (FIG. 25A, Lanes 4, 5 and 6).

*Allergenicity of Cow's Milk Derivatives.* The treated milk samples were subjected to skin tests using dogs sensitized to cow's milk as above. The results were quite similar to those obtained with pure BLG for both the moderately sensitive dog used above (FIG. 27 compared to FIG. 20) and with the group of 12 dogs tested (Table XXV compare to Table XXI). The end point and ED50 indicated that not only the DTT-reduced and cystamine- and GSSG-treated milk samples, but also the reoxidized milk sample were at least 10 times less allergenic than the control raw and heated milk samples.

*Digestibility studies with Cow's Milk.* Cystamine-treatment as well as DTT-reduction of milk rendered the BLG component highly susceptible to pepsin as observed above with pure BLG (FIG. 26, Lane 3 in Panels B and C). BLG in both raw and heated milk was stable in SGF (FIG. 26, Lane 3 in Panels A and F). As found with the pure protein, the GSSG-treated and reoxidized milk samples contained a form of BLG that, respectively, was resistant to digestion at 10% and 50% after 2 min (FIG. 26, Lane 3 in Panel D and E). Interestingly, cystamine-, GSSG-treatment and reoxidization did

not change the susceptibility of other milk proteins to pepsin digestion (FIG. 26, Panels C, D and E, Lane 3 vs Lanes 1 and 2).

### **Feeding Trials**

5 The atopic dogs are fed milk treated with the NTS and DTT which has been stabilized with either cystamine or GSSG. There is a reduction in the severity of vomiting and diarrhea with dogs fed the NTS-cystamine, NTS-GSSG stabilized milk.



Table XXI. Allergenicity of BLG following reduction by DTT at 55°C and derivatization with cystamine and GSSG.

Dog	Log (ED50)				
	Control*	DTT-reduced*	DTT/CYM*	DTT/GSSG*	DTT/REOXD*
7FA2	0.469	1.479	1.735	1.833	1.600
7FA8	0.322	1.318	1.695	2.301	2.559
7FB4	0.752	1.636	1.963	2.008	>1.50**
7FB5	0.403	>1.50**	1.439	1.427	1.362
7FB6	0.379	0.911	1.321	1.219	1.840
7FB9	0.400	1.006	1.095	2.079	0.868
7FC4	0.450	1.416	1.212	1.154	2.525
Mean±SD	0.454±0.140	1.294±0.282§	1.494±0.314§	1.717±0.451¶	1.792±0.664¶
Average of ED50 Relative Allergenicity‡	2.8 1.00	19.7 0.14	31.2 0.09	52.1 0.05	61.9 0.05

\* Control, untreated BLG (cf. Lane 4 in FIG. 4); DTT-reduced, BLG reduced with DTT followed by gel filtration to remove DTT (cf. Lane 3); DTT/CYM, BLG treated with cystamine following DTT-reduction (cf. Lane 5); DTT/GSSG, BLG treated with GSSG following DTT-reduction (cf. Lane 6); DTT/REOXD, BLG reoxidized after DTT-reduction (cf. Lane 7).

\*\* Only one positive reaction, whose wheal area was smaller than the half maximum wheal area, was detected, suggesting the ED50 was higher than the highest concentration injected.

§ P-value versus Control BLG was less than 0.0001.

¶ P-value versus Control BLG was less than 0.0005.

‡ Control value of ED50 divided by value obtained with each of the different treatments.

Table XXII. Allergenicity of BLG following reduction by NTS at 4°C and derivatization with cystamine and GSSG.

Dog	Log (ED50)					
	Control*	NTS-Reduced*	NTS/CYM*	NTS/GSSG*	NTS/REOXD*	DTT-Reduced*
7FA2	0.741	0.812	1.175	1.708	1.839	2.773
7FA8	0.179	-1.957	0.601	0.835	0.373	2.054
7FA9	0.822	1.864	2.148	0.970	0.753	3.292
7FB4	0.767	1.012	0.865	0.559	0.714	0.558
7FB5	0.988	1.089	0.919	0.669	2.774	2.156
7FB6	-0.081	0.183	0.243	0.418	0.090	0.996
7FB9	-0.521	-0.526	0.161	0.525	-0.351	0.412
7FC4	-0.216	-0.331	-0.056	0.230	0.127	0.522
Mean±SD	0.335±0.56 7	0.268±1.196	0.757±0.702 §	0.739±0.454¶	0.790±1.032‡	1.595±1.120†
Average of ED50 Relative Allergenicity✓	2.2 1.00	1.9 1.17	5.7 0.38	5.5 0.39	6.2 0.35	39.4 0.05

\* Control, untreated BLG (cf. Lane 1 in FIG. 3); NTS-reduced, BLG reduced by NTS at 4°C overnight (cf. Lane 2); NTS/CYM, BLG treated with cystamine following NTS-reduction (cf. Lane 3); NTS/GSSG, BLG treated with GSSG following NTS-reduction (cf. Lane 4); NTS/REOXD, BLG reoxidized after NTS-reduction (cf. Lane 5); DTT-reduced, BLG reduced with 2.5 mM DTT for 1 hour at 55°C followed by gel filtration to remove DTT (cf. Lane 6).

P-value versus Control BLG; § 0.0140, ¶ 0.0284, ‡ 0.0442, † 0.0020.

✓ Control value of ED50 divided by value obtained with each of the different treatments.

**Table XXIII** Effect of heat at 55°C on the alleviation of allergenicity of BLG by DTT/Trx reduction.

Dog	Log (ED50)				
	Control*	25°C DTT/Trx*	DTT*	Control*	55°C DTT/Trx*
7FA2	-0.001	-0.051	0.056	-0.186	1.507
7FA9	0.483	0.405	0.445	0.116	1.051
7FB4	0.262	0.531	0.477	0.578	1.352
7FB6	1.013	-0.147	0.093	0.794	1.406
7FB9	0.488	0.399	0.392	0.216	1.250
7FC8	-0.101	0.042	-0.055	0.285	1.065
Mean±SD	0.357±0.40 2	0.196±0.28 3	0.235±0.230	0.301±0.346	1.272±0.185§
Average of ED50 Relative Allergenicity‡	2.3 1.00	1.6 1.45	1.7 1.32	2.0 1.14	18.6 0.12

\* Control, BLG incubated without addition; DTT/Trx, BLG incubated with 2.5 mM DTT and 24 µg/ml Trx; DTT, BLG incubated with 2.5 mM DTT (cf. FIG. 4).

§ P-values: versus samples incubated at 25°C, 0.0015 (Control), 0.0005 (DTT/Trx) and 0.0002 (DTT); versus Control at 55°C, 0.0008.

¶ P-values: versus samples incubated at 25°C, 0.0028 (Control), 0.0004 (DTT/Trx) and 0.0003 (DTT); versus Control at 55°C, 0.0003.

‡ Control value of ED50 divided by value obtained with each of the different treatments.

**Table XXIV** Effect of heat after fully reduction by NTS on allergenicity of BLG.

Dog	Log (ED50)					
	Control*	NTS-reduced*	NTS-reduced & heated**	DTT-reduced*	NTS/CYM*	NTS & heated/CYM**
7FA2	0.746	0.482	0.757	1.048	1.024	0.854
7FA8	0.208	0.213	0.436	0.446	-0.140	0.670
7FA9	0.803	0.362	0.851	1.502	0.702	1.821
7FB4	0.623	1.605	2.927	2.554	2.772	1.758
7FB5	0.983	0.669	2.173	1.580	1.189	1.482
7FB6	0.625	0.734	1.383	0.696	1.030	0.480
7FB9	-0.063	0.377	0.599	0.796	0.657	0.171
Mean±SD	0.561±0.36 3	0.635±0.464	§1.304±0.926	1.232±0.715 ¶	1.033±0.883‡	1.034±0.653†
Average of ED50 Relative Allergenicity✓	3.6 1.00	4.3 0.84	20.1 0.18	17.1 0.21	10.8 0.34	10.8 0.34

\* The BLG samples were prepared as described in Tables 1 and 2.

\*\* BLG was heated for 1 hour at 55°C following NTS-reduction at 4°C overnight (NTS-reduced & heated). And then the reduced protein was incubated with 10 mM cystamine after removing NADPH by gel filtration. Excess cystamine was removed by second gel filtration (NTS & heated/CYM).

P-values versus Control and NTS-reduced BLG: § 0.0256 and 0.0081, ¶ 0.0144 and 0.0050, ‡ 0.0182 and 0.0641, † 0.0879 and 0.0289.

✓ Control value of ED50 divided by value obtained with each of the different treatments.

Table XXV Allergenicity of cow's milk following reduction by DTT at 55°C and derivatization with cystamine and GSSG.

Dog	Log (ED50)				
	Control*	Heated*	DTT-reduced*	DTT/CYM*	DTT/GSSG*
7FA2	-0.651	-0.317	1.086	1.256	0.670
7FA8	0.164	0.128	1.345	1.112	2.696
7FA9	0.287	0.393	0.351	>1.00**	>1.00**
7FB4	-0.008	0.085	2.753	>1.00**	>1.00**
7FB5	0.471	0.310	0.839	0.833	1.853
7FB6	0.212	-0.163	1.347	1.647	0.555
7FB9	-0.797	-0.726	0.672	1.073	0.423
7FC3	0.331	0.589	>1.00**	>1.00**	1.161
7FC4	-0.544	-0.498	0.485	2.032	0.116
7FC5	-0.146	-0.075	1.078	0.492	0.479
7FC6	0.111	-0.373	>1.50**	>1.50**	>1.50**
7FC8	-0.289	-0.148	>0.50**	>0.50**	>0.50**
Mean±SD	- 0.072±0.415	- 0.066±0.386	1.106±0.711 §	1.207±0.509§	0.994±0.872¶
Average of ED50	0.8	0.9	12.8	16.1	9.9
Relative allergenicity	1.00	0.99	0.07	0.05	0.09
					12.2
					0.07

\* Control, Raw cow's milk (cf. Lane 1 in FIG. 7). Remaining samples were incubated for 1 hour at 55°C and subjected to gel filtration. Heated, milk incubated with no addition (cf. Lane 2); DTT-reduced, milk incubated with DTT (cf. Lane 3); DTT/CYM, milk derivatized with cystamine after DTT-reduction (cf. Lane 4); DTT/GSSG, milk derivatized with GSSG after DTT-reduction (cf. Lane 5); DTT/REOXD, milk reoxidized after DTT-reduction (cf. Lane 6).

\*\* No spot or one positive spot, whose area was smaller than half maximum wheal area, was detected, suggesting the ED50 was higher than the highest concentration injected.

§ P-values versus raw and heated milk were less than 0.0001.

¶ P-values versus raw and heated milk were less than 0.0005.

‡ P-values versus raw and heated milk were 0.0030 and 0.0028, respectively.

### Figure Legends

FIGS. 19A and 19B. Change in redox state of BLG following reduction by DTT at 55°C and derivatization with cystamine and GSSG. BLG reduced with 2.5 mM DTT at 55°C (Lane 2) was subjected to gel filtration (Lane 3) and then was incubated with 10 mM cystamine (Lane 5) or 10 mM GSSG (Lane 6) or without addition (Lane 7). Fully reduced BLG (Lane 1) was prepared by boiling for 5 minutes with 2.5 mM DTT, and native BLG (Lane 4) was the control. The DTT-reduced BLG assayed with the skin test (FIG. 20 and Table 1 ) was labeled with mBBr after the test was completed (Lane 8). Ten µg of each BLG sample was subjected to mBBr-labeling/non-reducing SDS-PAGE. Left panel shows a profile of fluorescence of the protein-bound mBBr. Right panel shows a profile of protein stained with Coomassie brilliant blue R-250

FIG. 20. Response of a moderately sensitive dog to BLG following reduction with DTT at 55°C and derivatization with cystamine and GSSG. DTT/CYM, cystamine-treated BLG (cf. Lane 5 in FIG. 1); DTT/GSSG, GSSG-treated BLG (cf. Lane 6); DTT/REOXD, reoxidized BLG (cf. Lane 7); DTT-Reduced, BLG reduced with 2.5 mM DTT followed by gel filtration to remove DTT (cf. Lane 3); Control, native BLG (cf. Lane 4). The dog used was 7FA2.

FIGS. 21A and 21B. Change in redox state of BLG following reduction by NTS at 4°C and derivatization with cystamine and GSSG. BLG reduced by NTS (Lane 2) was incubated with 10 mM cystamine (Lane 3) or 10 mM GSSG (Lane 4) or without addition (Lane 5) after remove NADPH by gel filtration. Excess disulfide compounds were removed by second gel filtration after 3 days. NTS-reduced BLG was incubated for 1 hour at 55°C (Lane 7). Lane 1, control (native) BLG; Lane 6, DTT-reduced BLG; Lane 8, fully reduced BLG. Other details are given in FIG. 1.

FIGS. 22A and 22B. Redox state of BLG reduced with DTT plus Trx at 25 and 55°C. Lane 1, fully reduced BLG control. BLG was incubated at 25

°C without addition (Lane 2) or with DTT (Lane 3) or DTT plus Trx (Lane 4). BLG was incubated at 55°C without addition (Lane 5) or with DTT (Lane 6) or DTT plus Trx (Lane 7). Other details are given in FIG. 1.

FIGS. 23A, 23B and 23C. Susceptibility of reduced and derivatized BLG to digestion by simulated gastric fluid. Each BLG sample was incubated in SGF at 37°C for 2 minutes. The reaction was stopped by addition of 160 mM  $\text{Na}_2\text{CO}_3$  (stop solution). SGF was premixed with the stop solution and added to each BLG sample (0 minute). "-", native BLG (not treated with SGF). Panel A: Reducing SDS-PAGE analysis of the digested samples of the BLG derivatives following DTT-reduction at 55°C. Lanes 1 to 3, native BLG (Control); Lanes 4 and 5, BLG reduced with DTT at 55°C followed by Sephadex G-25 gel filtration (DTT-reduced); Lanes 6 and 7, cystamine-treated BLG (DTT/CYM); Lanes 8 and 9, GSSG-treated BLG (DTT/GSSG); Lanes 10 and 11, reoxidized BLG (DTT/REOXD). Panel B: Non-reducing SDS-PAGE analysis of the digested samples of DTT-reduced and GSSG-treated BLG. Lane 1, DTT-reduced BLG; Lane 2, native BLG; Lanes 3 and 4, GSSG-treated BLG (DTT/GSSG). Panel C: Reducing SDS-PAGE analysis of the digested samples of BLG derivatives following NTS-reduction at 4°C. Lanes are referred to Panel A except that NTS (at 4°C) was used for reduction.

FIG. 24. Time course of digestion of reduced and derivatized BLG by simulated gastric fluid. Each BLG sample was incubated in SGF at 37°C. After 2, 15, 30 and 60 minutes, the reaction was stopped by addition of 160 mM  $\text{Na}_2\text{CO}_3$  (stop solution). SGF was premixed with the stop solution and added to the each BLG sample (0 minute). The digested sample was subjected to reducing SDS-PAGE and undigested BLG was quantified by Multi-Analyst ver. 1.0.2. Open symbols: BLG derivatives, which were reduced by DTT. Closed symbols: BLG derivatives, which were reduced by NTS. Diamond, reduced; circle, cystamine-treated; square, GSSG-treated; triangle, reoxidized; X, native BLG.

FIGS. 25A and 25B . Change in redox state of milk proteins following reduction by DTT at 55°C and derivatization with cystamine and GSSG. Cow's milk reduced with 50 mM DTT at 55°C (Lane 8) was subjected to Sephadex G-25 gel filtration (Lane 3) and then incubated with 10 mM cystamine (Lane 4) or 10 mM GSSG (Lane 5) or without addition (Lane 6). Raw milk (Lane 1), heated milk (Lane 2) and fully reduced milk (Lane 9) served as controls. The reduced milk sample used in the skin test (FIG. 9 and Table 5) was labeled with mBBBr after the test was completed (Lane 7). Fifty µg of the milk sample was subjected to mBBBr-labeling/non-reducing SDS-PAGE. Left panel shows a profile of fluorescence of the protein-bound mBBBr. Right panel shows a profile of protein stained with CBB.

FIG. 26. Stability of reduced and derivatized milk proteins in simulated gastric fluid. Each milk sample was incubated in SGF without pepsin for 60 minutes at 37°C (Lane 1), mixed with complete SGF which was premixed with the stop solution (Lane 2) and incubated in complete SGF for 2 minutes at 37°C (Lane 3). After the incubation, 100 µg of protein of each sample was analyzed by reducing SDS-PAGE. A, Raw milk (Control); B, milk reduced with DTT followed by Sephadex G-25 gel filtration (Red); C, cystamine-treated milk (CYM); D, GSSG-treated milk (GSSG); E, reoxidized milk (REOXD), F, heated milk (Heated).

FIG. 27. Response of a moderately sensitive dog to cow's milk following reduction with DTT and derivatization with cystamine and GSSG. DTT/CYM, cystamine-treated milk (cf. Lane 4 in FIG. 7); DTT/GSSG, GSSG-treated milk (cf. Lane 5); DTT/REOXD, reoxidized milk (cf. Lane 6); DTT-Reduced, milk reduced with 50 mM DTT followed by gel filtration to remove DTT (cf. Lane 3); Heated, milk incubated for 1 hour at 55°C followed by gel filtration (cf. Lane 2); Control, raw milk (cf. Lane 1). The dog used was 7FA2.



## EXAMPLE 45

### Stabilization of Reduced Peanut Extract

#### *Experimental Procedures*

*Chemicals and Proteins.* Peanut allergen extract containing Ara H1-6 was obtained from Meridian Biomedical. The other chemicals and reagents were the same as those used in Example 44.

*Reduction and Blocking of Protein Disulfide Bond.* Peanut extract, 1mg/ml, was reduced by 2.5mM DTT in phosphate buffered saline (PBS) for 1 hr at 55°C or by the NADP/thioredoxin system (NTS) for 5 hr at 37°C.

Following gel filtration with a Sephadex G-25 column (Amersham Pharmacia) to remove reductants, the reduced protein was incubated either without addition or with 10mM oxidized glutathione (GSSG) or 10mM cystamine for 3 days on ice. The mixtures were then subjected to a second gel filtration on a Sephadex G-25 column to remove excess GSSG and cystamine. If necessary, the resultant preparations were concentrated by ultrafiltration with Centricon YM-3 filter (Millipore). Heat treatment of NTS-reduced peanut extract was carried out for 1 hour at 55°C after reduction at 37°C overnight. Heated peanut extract was prepared by incubation for 1 hour at 55°C and gel filtration with A Sephadex G-25 column.

*Monobromobimane Labeling and Analysis of Proteins.* Fifty µg of peanut extract was incubated with 4mM mBBR for 20 min as described in previous Examples and also as described in Wong *et al.*, 1995, and then subjected to non-reducing SDS-PAGE (10 to 20% gradient gel) to separate the native, partly reduced and fully reduced forms. The labeled proteins were stored at -80°C, if not immediately loaded on the gel. The gel was prepared and treated as described in Example 44. The intensity of fluorescence and staining of each band was also quantified as in Example 44.

*Skin Tests.* The same colony of dogs as used in Example 44 are used for skin testing the allergenicity of the treated peanut extracts. The dogs differ from those in Example 44 in that they are all immunized with peanut extract.

*Estimation of Relative Allergenicity.* Relative allergenicity is determined using the same methods described in Example 44.

*Assay for Stability to Pepsin.* Peanut extract (0.2mg/ml) is mixed with simulated gastric fluid (SGF) to give 0.32% pepsin. Otherwise, the same methods as used to determine stability to pepsin in Example 44 are used in this example.

*Quantification of Protein.* Protein is analyzed by using the Bradford method described above.

## ***Results***

*Reduction by NTS and DTT.* Peanut extracts differed from BLG in reduction by NTS. Thus, while BLG was reduced by NTS at 25°C in 1 hr., peanut extracts required an incubation time of 5 hrs. at 37°C. With DTT, peanut proteins required 1 hr. at 55°C. Otherwise, the results with respect to mitigation of allergenicity and susceptibility to pepsin are similar to those obtained in Example 44 with BLG with the exception that, as in the examples other than 44, the NTS treated proteins definitely show a reduction in allergenicity. Here, peanut extract that is reduced by DTT exhibits reduced allergenicity and increased digestibility by pepsin.

Peanut extract that is reduced by the NTS exhibits increased digestibility by pepsin. Mitigation of allergenicity also occurs when the NTS or DTT reduced peanut extract is either heated or derivatized with cystamine or GSSG. Reduced allergenicity also occurs when the DTT reduced peanut extract is allowed to reoxidize. Increased digestibility by pepsin occurs when the peanut extract is reduced by DTT and heat or NTS or when the NTS or DTT reduced peanut extract is derivatized with cystamine. The increase in susceptibility to pepsin also occurs if the reduced peanut extract reduced by

DTT and heat or NTS is derivatized with GSSG but the increase in susceptibility does not occur to the same extent as derivatization with cystamine.

#### **EXAMPLE 46**

##### **Stabilization of Reduced Ragweed Allergen Proteins**

###### ***Experimental Procedures***

***Chemicals and Proteins.*** Giant ragweed pollen allergen extract is obtained from Bayer, Inc. The other chemicals and reagents are the same as those used in Example 44.

***Reduction and Blocking of Protein Disulfide Bond.*** Giant ragweed pollen extract 1 mg/ml is reduced by 2.5mM DTT in phosphate buffered saline (PBS) for 1 hr at 55°C or by the NADP/thioredoxin system (NTS) for 17 hr at 4°C as essentially described in Example 44. Following gel filtration with a Sephadex G-25 column (Amersham Pharmacia) to remove reductants, the reduced protein is incubated either without addition or with 10 mM oxidized glutathione (GSSG) or 10 mM cystamine for 3 days on ice. The mixtures are then subjected to a second gel filtration on a Sephadex G-25 column to remove excess GSSG and cystamine. If necessary, the resultant preparations are concentrated by ultrafiltration with Centricon YM-3 filter (Millipore). Heat treatment of NTS-reduced pollen extract is carried out for 1 hour at 55°C after reduction at 37°C for 45 min. Heated pollen extract is prepared by incubation for 1 hour at 55°C and gel filtration with A Sephadex G-25 column.

***Monobromobimane Labeling and Analysis of Proteins.*** Fifty µg of pollen extract is incubated with 4 mM mBBR for 20 min as described in previous Examples and also as described in Wong *et al.*, 1995, and then subjected to non-reducing SDS-PAGE (10 to 20% gradient gel) to separate the native, partly reduced and fully reduced forms. The labeled proteins are stored at -80°C, if not immediately loaded on the gel. The gel is prepared and treated

as described in Example 44. The intensity of fluorescence and staining of each band are also quantified as in Example 44.

*Skin Tests.* The same colony of dogs as used in Example 44 are used for skin testing the allergenicity of the treated pollen extracts. Again, the dogs differ from those in Example 44 in that they are from a different litter and are also all immunized with giant ragweed pollen extract.

*Estimation of Relative Allergenicity.* Relative allergenicity is determined using the same methods described in Example 44.

*Assay for Stability to Pepsin.* The pollen extract 0.2mg/ml is mixed with simulated gastric fluid (SGF) to give 0.32 % pepsin. Otherwise, the same methods as used to determine stability to pepsin in Example 44 are used in this example.

*Quantification of Protein.* Protein is analyzed by using the Bradford method as described in Example 44.

### ***Results***

The results with respect to mitigation of allergenicity and susceptibility to pepsin are similar to those obtained in Example 44 with BLG, but again, the NTS treated protein, Amb t V, definitively shows a reduction in allergenicity. Here, the pollen proteins that are reduced by DTT and heat exhibit reduced allergenicity and increased digestibility by pepsin.

Pollen extract, namely Amb t V, and other disulfide allergens that are reduced by the NTS exhibit increased digestibility by pepsin. Mitigation of allergenicity also occurs when the NTS or DTT reduced pollen extract is either heated or derivatized with cystamine or GSSG. Reduced allergenicity also occurs when the DTT reduced peanut extract is allowed to reoxidize. Increased digestibility by pepsin occurs when the peanut extract is reduced by

DTT and heat or NTS or when the NTS or DTT reduced peanut extract is derivatized with cystamine. The increase in susceptibility to pepsin also occurs if the reduced pollen extract reduced by DTT and heat or NTS is derivatized with GSSG but the increase in susceptibility does not occur to the same extent as derivatization with cystamine.

### **CONCLUDING REMARKS**

It can be seen from the foregoing general description of the invention and from the specific examples illustrating applications thereof, that the invention has manifold and far reaching consequences. The invention basically provides novel dough and dough mixtures and novel methods for creating new doughs and for improving the quality of dough and baked goods as well as novel methods for inactivating enzyme inhibitors in cereal products. The invention also provides a novel method for altering the biological activity and inactivity of animal toxins, and for eliminating or decreasing the allergenicity of several allergens, namely pollens and foods, namely wheat, egg, milk, whey, soy, nuts and beef. The invention further provides a method for increasing the proteolysis and digestibility of pollen and food allergens, particularly milk, whey and their products and wheat products. Also, the invention provides for stabilizing hypoallergenic and hyperdigestible previously reduced food and pollen proteins by physiological disulfides. It also provides for stabilizing hypoallergenic previously reduced proteins by heat and reoxidation. In addition, the invention provides a novel protein that is a pullulanase inhibitor and a method for its inactivation.

While the invention has described in connection with certain specific embodiments thereof, it should be realized that various modifications as may be apparent to those of skill in the art to which the invention pertains also fall within the scope of the invention as defined by the appended claims.

## References

- Lehrer, S. B., Horner, W. E. and Reese, G., "Why are some proteins allergenic? Implications for biotechnology." *Crit. Rev. Food Sci. Nutr.* 36:553-564 (1996).
- Buchanan, B. B., Adamidi, C., Lozano, R. M., Yee, B. C., Momma, M., Kobrehel, K., Ermel, R. and Frick, O. L. "Thioredoxin-linked mitigation of allergic responses to wheat." *Proc. Natl. Acad. Sci. USA* 94:5372-5377 (1997).
- del Val, G., Yee, B. C., Lozano, R. M., Buchanan, B. B., Ermel, R. W., Lee, Y.-M and Frick, O. L. "Thioredoxin treatment increases digestibility and lowers allergenicity of milk." *J. Allergy Clin. Immunol.* 103:690-697 (1999).
- Jiao, J., Yee, B. C., Kobrehel, K. and Buchanan, B. B. "Effect of thioredoxin-linked reduction on the activity and stability of the Kunitz and Bowman-Birk soybean trypsin inhibitor proteins." *J. Agric. Food Chem.* 40:2333-2336 (1992).
- Wong, J., II, Kobrehel, K. and Buchanan, B. B. "Thioredoxin and seed proteins." *Methods Enzymol.* 252:228-240 (1995).
- Ermel, R. W., Kock, M., Griffey, S. M., Reinhart, G.A. and Frick, O. L. "The atopic dog: a model for food allergy." *Lab. Anim. Sci.* 47:40-40 (1997).
- Frick, O. L and Brooks, D. L. "Immunoglobulin E antibodies to pollens augmented in dogs by virus vaccines." *Am. J. Vet. Res.* 44:440-445 (1983).
- Astwood, J. D., Leach, J. N. and Fuchs, R. L. "Stability of food allergens to digestion *in vitro*." *Nature Biotechnol.* 14:1269-1273 (1996).

Laemmli, U. K. "Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685 (1970).

Bradford, M. M. "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding." *Anal. Biochem.* 72:248-254 (1976).